Expression levels of domestic cDNA cassettes integrated in the nuclear
genomes of various *Chlamydomonas reinhardtii* strains

Fantao Kong,¹ Tomohito Yamasaki,¹ and Takeshi Ohama^{1, 2*}

¹*School of Environmental Science and Engineering, Kochi University of Technology
(KUT), Tosayamada, Kochi 782-8502 and* ² *Japan Science and Technology Agency,
CREST, 5, Sanbancho, Chiyoda-ku, Tokyo, 102-0075, Japan.*

Keywords:

cDNA expression cassettes, *Chlamydomonas reinhardtii*, nuclear transformation, UVM
strains

Corresponding author:

Takeshi Ohama

E-mail: ohama.takeshi@kochi-tech.ac.jp.

Tel.: +81-887-57-2512

Fax: +81-887-57-2520

Running title: Expression of domestic cDNA cassettes in *Chlamydomonas*

Abstract

We attempted to overexpress three types of expression cassettes, each of which contained a different open reading frame (ORF) of domestic *Chlamydomonas* cDNAs. Each ORF was strongly driven by an artificial hybrid promoter. We used two wild-type *Chlamydomonas* strains (i.e., CC-124 and CC-125) and two mutant strains [i.e., UV-mutated (UVM) 4 and UVM11] that have been reported to have a high potency for expressing nondomestic nuclear transgenes. We found that the 1-deoxy-D-xylulose-5-phosphatesynthase (*DXSI*), 1-deoxy-D-xylulose 5-phosphate reductoisomerase (*DXRI*), and squalene synthase (*SQS*) cassettes were not readily overexpressed in the wild-type strains at levels where the products were clearly detectable by Western blotting using a monoclonal antibody. In contrast, Western blot-positive *SQS* cassette transformants were frequently detected in the UVM4 and UVM11 strains, i.e., at an approximately 4.5 times higher frequency than that in the CC-124 wild-type strain. Moreover, transformants that accumulated large amounts of the *SQS* protein were obtained frequently in the UVM4 and UVM11 strains, i.e., the frequency was approximately 2.2 times higher than that in the CC-124 strain. However, a position effect of the integrated expression cassette was obviously detected not only in the wild-type but also in UVM strains. This suggests that the epigenetic repression mechanism of transgenic genes was not completely knocked out, even in the UVM strains. Further improved *Chlamydomonas* strains are essential to facilitate high-throughput screening of transformants that express nuclear transgenes at a high level.

1 **Introduction**

2 The unicellular green alga, *Chlamydomonas reinhardtii*, is expected to be a
3 potentially low-cost host for generating various products, e.g., biofuel, pharmaceutical
4 terpenoid, and vaccine, because of its rapid phototrophic growth potency and the
5 availability of several advanced molecular techniques (1). However, the nuclear
6 expression levels of nondomestic cDNA genes are disappointingly poor (2, 3).
7 Unsuitable codon usage in the open reading frame (ORF) is one of the proven causes of
8 this problem (4). The random integration of transgenes in *Chlamydomonas reinhardtii*
9 has been analyzed using the RESDA-PCR method (5).

10 Moreover, the integration position of the construct dramatically affects the expression
11 level (1, 6), which is referred to as the “position effect.” Recently, Neupert *et al.* (7)
12 developed *C. reinhardtii* UV-mutated (UVM) strains to overcome the inhibited
13 expression of foreign genes. These cell wall-deficient strains, UVM4 and UVM11,
14 harbor unknown and unmapped mutation(s), and it has been suggested that the
15 epigenetic transgene suppression mechanisms have been successfully knocked out in
16 these mutants. To the best of our knowledge, high expression in these UVM strains has
17 only been demonstrated for green fluorescent protein (*GFP*) and yellow fluorescent
18 protein (*YFP*) (7). Moreover, all of the reported genes are nondomestic genes, and their
19 codon usage was not optimized for *C. reinhardtii*. This suggests that we might readily
20 obtain transformants that accumulate high levels of target proteins if endogenous genes
21 could be introduced into the UVM strains, because their codon usage would not
22 interfere with their translation.

23 1-Deoxy-D-xylulose-5-phosphatesynthase (DXS1) and 1-deoxy-D-xylulose
24 5-phosphate reductoisomerase (DXR1) proteins are expected to be key enzymes

involved with isoprene synthesis in the 2-C-methyl-D-erythritol 4-phosphate (MEP) pathway (8), while squalene synthase (SQS) protein catalyzes the first enzymatic step in the central isoprenoid pathway during sterol and triterpenoid biosynthesis (9). These three genes are unique and essential and contain a limited number of rare codons, which probably have no harmful effects on their translation levels.

In this study, we examined the expression levels of the cDNA cassettes of DXS1 and SQS in the wild-type strain CC-124, and that of DXR1 in the wild-type strain CC-125. We also introduced the *SQS* expression cassette into the UVM4 and UVM11 strains to compare the expression levels of the wild-type strain and the UVM mutants.

Materials and Methods

Construction of transformation vectors

The cassettes of *Hsp70A/RbcS2* promoter (containing the first intron of *RbcS2*) and *RbcS2* terminator were cloned into pUC18 (without the *Bam*HI restriction enzyme site) as a *Sac*I/*Kpn*I fragment to generate the pHsp70A/RbcS2-plasmid. This artificial promoter also contains the first intron of *RbcS2* because of its enhancer activity (10). A gp64-tag sequence (11) was integrated into the respective constructs for analysis with Western blotting. The fragments containing the gp64-tag sequences were as follows:

gp64 tag 1:

5'-GATGCTCGAGTCCTGGAAGGACGCGTCCGGCTGGTCCCATATGTAATAAAG

TACTACTAGTGGATCCCCGCT-3' and gp64 tag 2:

5'-GATGCTCGAGAGATCTACACCGACTAGTTCCTGGAAGGACGCGTCCGGCT

GGTCCTAATAAGGATCCCCGCT-3'. These sequences were synthesized and cloned

into the *XhoI/BamHI* site of pHsp70A/RbcS2 plasmid to construct the pHsp70A/RbcS2-gp64 tag 1 and pHsp70A/RbcS2-gp64 tag 2 plasmids, respectively.

DXS1, *DXR1*, and *SQS* cDNA were amplified using polymerase chain reaction (PCR) with high-fidelity PrimeSTAR HS DNA polymerase (Takara, Japan), according to the manufacturer's instructions. The specific sets of primers used for each amplification reaction were as follows: DXS1fw (5'-AAGAGATCTATGCTGCGTGGTGCTG-3') and DXS1rev (5'-AAGACTAGTCGCTTGCAGCGCTGA-3') for *DXS1*, which were generated by introducing *BglII* and *SpeI* sites at the 5' ends, respectively (the restriction endonuclease sites are underlined); DXR1fw (5'-AAGCTCGAGATGCAGTCCGCGCAGC-3') and DXR1rev (5'-AAGACTAGTGGCCATGGCCAGCACC-3') for *DXR1*, which were generated by introducing *XhoI* and *SpeI* sites at the 5' ends, respectively; and SQSfw (5'-CATATGGGAAAGTTAGGGGAGCTACTCTC-3') and SQSrev (5'-GGATCCTCACGCACGCCGCA-3') for *SQS*, which were generated by introducing *NdeI* and *BamHI* sites at the 5' ends, respectively. The amplification conditions were as follows: 98°C for 1 min, followed by 35 cycles of denaturation at 98°C for 10 s, annealing at 54°C for 15 s, and extension at 72°C for 2 min using a Thermal Cycler 2720 (Applied Biosystems, Foster City, CA, USA).

The *DXS1* cDNA fragment (~2.2 kb) was inserted into the pHsp70A/RbcS2-gp64 tag 1 plasmid as a *BglII/SpeI* fragment, generating the recombinant plasmid pHsp70A/RbcS2-gp64-DXS1. The *DXR1* cDNA fragment (~1.4 kb) was inserted into the pHsp70A/RbcS2-gp64 tag 1 plasmid as a *XhoI/SpeI* fragment, generating the plasmid pHsp70A/RbcS2-gp64-DXR1. The *SQS* cDNA fragment (~1.4 Kb) was inserted into the pHsp70A/RbcS2-gp64 tag 2 plasmid as a *NdeI/BamHI* fragment,

generating the plasmid Hsp70A/RbcS2-gp64-SQS. The pHyg3 plasmid, which contained *aphVII* and conferred hygromycin resistance (12), was double digested with *HindIII* and subcloned into the Hsp70A/RbcS2-gp64-DXS1 plasmid in the sense orientation to generate the *DXS1* overexpression construct. The pSI103 plasmid, which contained *aphVIII* and conferred paromomycin resistance (13), was double digested with *XbaI* and *KpnI* and then subcloned into the Hsp70A/RbcS2-gp64-DXR1 plasmid in the sense orientation to generate the *DXR1* overexpression construct. The pSP124S plasmid, which contained *ble* and conferred zeocine resistance (14), was double digested with *XbaI* and *KpnI* and then subcloned into the Hsp70A/RbcS2-gp64-SQS plasmid in the sense orientation to generate the *SQS* overexpression construct. The DNA sequences of all the constructs were confirmed by direct sequencing using the dideoxy chain termination method (15). The final *DXS1*, *DXR1*, and *SQS* overexpression constructs are shown schematically in Supplementary Fig. S1.

Strains and transformation conditions

C. reinhardtii CC-124 (wild-type, *mt*⁻) and CC-125 (wild-type, *mt*⁺) were provided by the *Chlamydomonas* Resource Center (Duke University, USA), while the UVM4 and UVM 11 strains (7) were kindly provided by Dr. R. Bock (MPI-MP, Germany). Unless stated otherwise, cells were cultivated mixotrophically at 25°C in Tris-acetate phosphate (TAP) medium (16) under constant cool white fluorescent light (84 $\mu\text{mol photons m}^{-2} \text{s}^{-1}$) with gentle shaking. Before transformation, vectors were linearized with a single restriction enzyme and purified using the Wizard SV Gel and PCR Clean-Up System (Promega, USA), according to the manufacturer's protocols. Approximately 300 ng linearized plasmid DNA was used during each transformation

experiment by electroporation, as described previously (17). Transformants of the *DXS1*, *DXR1*, and *SQS* overexpression constructs were selected directly on 1.5% TAP agar plates containing hygromycin B (30 mg L⁻¹), paromomycin sulfate (20 mg L⁻¹), or zeocin (15 mg L⁻¹), respectively, and the plates were incubated under continuous fluorescent light (20 μmol m⁻² s⁻¹) at 25°C.

PCR screening of cotransformants

Transformants with antibiotic resistance were screened to detect the presence of the expression cassettes using a promoter-specific forward primer and a gene-specific reverse primer, the locations of which are shown in Supplementary Fig. S1. Genomic DNA was isolated from independent transformants using the CTAB method (18). The specific sets of primers used to examine construct integration were as follows: forward, 5'-AAAATGGCCAGGTGAGTCGACG-3' and reverse, 5'-GATTGACCGAAGCCTCCGCG-3' for *DXS1*; forward, 5'-CGTTTCCATTTGCAGGCTCGAGA-3' and reverse, 5'-AGCAGCTTCACGTTGGAGCC-3' for *DXR1*; and forward, 5'-TTGCAGGATGCTCGAGTCCTG-3' and reverse, 5'-CCACGTCCAGTGCCATGTCA-3' for *SQS*. The amplification conditions were as follows: 95°C for 5 min; followed by 30 cycles at 95°C for 15 s, 54°C for 30 s, and 72°C for 30 s; and a final step at 72°C for 7 min using the Thermal Cycler 2720 (Applied Biosystems).

Western blot screening of transformants

Total soluble protein extracts were prepared by resuspending pelleted 5.0×10^6 *Chlamydomonas* cells in 50 μ L of 2X SDS sample buffer (100 mM Tris, pH 6.8; 4% SDS; 1 mM DTT; and 30% glycerol). The samples were denatured at 95°C for 7 min. Subsequently, 5- μ L aliquots of the supernatants from the denatured cell lysates were separated using 10% SDS-PAGE, transferred using a Trans-Blot Electrophoretic Transfer Cell (Bio-Rad, USA) to polyvinylidene difluoride (PVDF) membranes (Hybond P; GE Healthcare, UK), and incubated overnight with standard transfer buffer (25 mM Tris, 20% methanol, 0.01% SDS, and 192 mM glycine). Immunobiochemical protein detection was performed using a monoclonal anti-BV envelope gp64 primary antibody (1/5,000) (Santa Cruz, CA, USA) and ECL peroxidase-labeled anti-mouse secondary antibody (1/20,000) (Sigma-Aldrich, Canada) with an ECL detection system (GE Healthcare).

Results and Discussion

All of the vectors used for transformation contained one of the three expression cassettes with a marker gene cassette in the same strand (Supplementary Fig. S1). The gp64 epitope tag for analysis with Western blotting was introduced at the N-terminal end in *SQS*, whereas it was introduced at the C-terminal end in *DXS1* and *DXR1* (Supplementary Fig. S1), given that the N-terminal transit peptide region of *DXS1* and *DXR1* proteins are removed post-transnationally to localize these proteins in plastids.

The linearized plasmids for *DXS1* or *SQS* were introduced into the CC-124 strain (*mt*⁻) to allow overexpression, whereas the linearized plasmid for *DXR1* was introduced into the CC-125 strain (*mt*⁺). The antibiotic resistance characteristics used for selection guaranteed the integration of the marker gene in the transformants, but this does not

1 guarantee the cointegration of the cDNA expression cassette in the transformants. To
2 exclude transformants without the cointegrated expression cassette of the target gene,
3 we performed PCR to confirm the integration of the promoter region. PCR was
4 performed using a set of primers, which were specific to 3' end of the *Hsp70A/RbcS2*
5 promoter and the N-terminal region of the cDNA expression cassette. For the
6 PCR-positive transformants, we assumed that the intact cDNA cassette was integrated
7 into the genome with the marker gene cassette; therefore, we tentatively counted these
8 transformants as being successfully cotransformed (Table 1). The PCR results showed
9 that the cotransformation ratios (number of PCR-positive transformants/number of
10 transformants analyzed) for *DXS1* with *aphVII* (introduced into the CC-124 strain),
11 *DXR1* with *aphVIII* (introduced into the CC-125 strain), and *SQS* with *ble* (introduced
12 into the CC-124 strain) were approximately 42.4% (112/264), 21.8% (55/252), and 51.9%
13 (109/210), respectively (Table 1). The relatively lower cotransformation ratio of the
14 *DXR1* cassette was probably because of the characteristics of the CC-125 strain.
15 Recently, it was reported that the optimal transformation efficiency, using the
16 electroporation method (19), for the CC-125 strain was approximately 66% of that for the
17 CC-124 strain.

18 Western blotting was performed to detect the protein accumulation levels of the
19 PCR-positive transformants. The results showed that the Western blot-positive ratios
20 (number of Western blot-positive transformants/number of PCR-positive transformants)
21 for the *DXS1* and *SQS* transformants in the CC-124 strain were 0.89% (1/112) and 4.6%
22 (5/109), respectively (Table 1 and Fig. 1A). However, no *DXR1*-positive transformants
23 were found among 55 PCR-positive transformants in the CC-125 strain (Table 1). The
24 low expression levels of the target protein in the wild-type strains were in agreement

with previous reports (1-3). The sensitivity of Western blotting performed using a monoclonal antibody for gp64-tag was sufficiently high to detect the protein derived from gp64-tag-containing *aphVII* transformants (data not shown).

Encouraged by the successful overexpression of three nonendogenous fluorescence-related genes in the UVM4 and UVM11 strains (7, 20), we examined the overexpression of the *SQS* cassette in the UVM strains. The PCR results showed that the estimated cotransformation ratios for *SQS* in the UVM 4 and UVM 11 strains were approximately 59.8% (52/87) and 58.6% (51/87) (Table 1), respectively, which were only approximately 8% and 7% higher than those in the CC-124 strain. These results were not consistent with the results of Neupert *et al.* (7), who showed that the cotransformation ratios of the full-length *GFP* cassette in the genomes of the UVM4 and UVM11 strains were about five times higher than that of the control strain (CW-15, *mt⁺*, *arg7*).

However, the Western blotting results detected dramatic differences between the UVM and wild-type strains. The Western blot-positive ratios for *SQS* in the UVM 4 and UVM 11 strains were approximately 21.2% (11/52) and 17.6% (9/51) (Table 1 and Fig. 1B, 1C), respectively, which were about five and four times higher than the ratio in the CC-124 strain (4.6%, 5/109). Moreover, the high-protein accumulation ratios (number of high-protein accumulation transformants/number of Western blot-positive transformants) of the Western blot-positive clones were 45.5% (5/11) for the UVM4 strain and 44.4% (4/9) for the UVM11 strain (Table 1 and Fig. 1B, 1C). These ratios were approximately two times higher than those in the CC-124 strain (20.0%, 1/5) (Table 1 and Fig. 1A).

Unstable transgene expression has been observed frequently in *C. reinhardtii* (2). However, the *SQS* transformants generated from the UVM strains were cultured on antibiotic-containing plates for over 5 months, and they exhibited stable expression (data not shown). In addition, the growth rates of all the *SQS* protein-expressing transformants of the UVM4 and UVM11 strains did not differ significantly from the CC-124 strain (data not shown) in mixotrophic conditions (in TAP medium) with a 16/8-h light/dark cycle or continuous light. These results suggest that the UVM strains are highly suited to the endogenous cDNA transgene expression of moderate-molecular-weight proteins, and they are not restricted to the expression of low-molecular-weight proteins such as *GFP*.

The *SQS* cDNA cassette originated from domestic mRNA; therefore, its codon usage and substitution in the ORF must have been optimized by evolution. Thus, *SQS* may have fewer translational disadvantages than foreign genes. However, Fig. 1 shows that the *SQS* protein accumulation levels were not uniformly high even in the UVM strains, which was also the case in the CC-124 strain. It is reasonable to suspect that translational repression works uniformly against all *SQS* mRNAs, irrespective of the genomic locations of the *SQS* that generated the *SQS* mRNA. Therefore, the observed differences in the levels of accumulated *SQS* protein must have been because of the variable transcriptional levels of the *SQS* located in different genomic regions.

Neupert *et al.* (7) reported that all PCR-positive *GFP* transformants (9/9) had uniformly high protein accumulation levels. Based on this fact, they suggested the successful knock out of the transcriptional repression pathways of nuclear transgenes in the UVM strains. However, our results of *SQS* expression showed that epigenetic transcriptional repression was alleviated dramatically, although they were not

completely free of it even in the UVM strains. This might have been due to the fact that the mutated gene(s) had not lost their activity completely, because these UVM strains were generated by UV treatments to introduce nucleotide mutations. The observed inconsistent characteristics of the UVM strains mentioned above might have been due to the different characteristics of the promoters used to drive the cDNAs, i.e., *GFP* was driven by either *PsaD* or *RbcS2*, whereas *SQS* was transcribed by the *Hsp70A/RbcS2*-hybrid promoter (21).

Further improvements are required to obtain truly epigenetic repression-free *Chlamydomonas* strains. The development of such strains would make *Chlamydomonas* an attractive host for producing biofuel and biopharmaceuticals.

Acknowledgements

The authors would also like to thank Dr. R. Bock (MPI-MP, Germany) for kindly providing *C. reinhardtii* UVM4 and UVM 11 strains. This research was supported by Japan Science and Technology Agency, CREST.

References

1. Rasala, B., Lee, P., Shen Z., Briggs, S., Mendez, M., and Mayfield, S.: Robust expression and secretion of *Xylanase1* in *Chlamydomonas reinhardtii* by fusion to a selection gene and processing with the FMDV 2A peptide, PLoS ONE, **7**, e43349 (2012).
2. Fuhrmann, M., Oertel, W., and Hegemann, P.: A synthetic gene coding for the green fluorescent protein (GFP) is a versatile reporter in *Chlamydomonas reinhardtii*,

Plant J., **19**, 353-361 (1999).

3. **Schroda, M., Blocker, D., and Beck, C.:** The HSP70A promoter as a tool for the improved expression of transgenes in *Chlamydomonas*, Plant J., **21**, 121-131 (2000).

4. **Shao, N. and Bock, R.:** A codon-optimized luciferase from *Gaussia princeps* facilitates the in vivo monitoring of gene expression in the model alga *Chlamydomonas reinhardtii*, Curr. Genet., **53**, 381-388 (2008).

5. Gonzalez-Ballester, D., de Montaigu, A., Galvan, A. and Fernandez, E.: Restriction enzyme site-directed amplification PCR: a tool to identify regions flanking a marker DNA. Anal. Biochem., **340**, 330–335 (2005).

6. **Cerutti, H., Johnson, AM., Gillham, NW., and Boynton, J.:** Epigenetic silencing of a foreign gene in nuclear transformants of *Chlamydomonas*, Plant Cell, **9**, 925-945 (1997).

7. **Neupert, J., Karcher, D., and Bock, R.:** Generation of *Chlamydomonas* strains that efficiently express nuclear transgenes, Plant J., **57**, 1140-1150 (2009).

8. **Cordoba, E., Salmi, M., and Leon, P.:** Unravelling the regulatory mechanisms that modulate the MEP pathway in higher plants, Journal of Experimental Botany, **60**, 2933-2943 (2009).

9. **Lee M-H., Seo J-W., Shin C-G., Kim Y-S., In J-G., Yang D-C., Yi J-S., and Choi Y-E.:** Enhanced triterpene and phytosterol biosynthesis in *Panax ginseng* overexpressing squalene synthase gene, *Plant Cell Physiol.*, **45**, 976–984 (2004).
10. **Lumbreras, V., Stevens, D. R., and Purton, S.:** Efficient foreign gene expression in *Chlamydomonas reinhardtii* mediated by an endogenous intron, *Plant J.*, **4**, 441-447 (1998).
11. **Zhou, J. and Blissard, G.:** Identification of a GP64 subdomain involved in receptor binding by budded virions of the baculovirus *autographica californica* multicapsid nucleopolyhedrovirus, *J. Virol.*, **82**, 4449-4460 (2008).
12. **Berthold, P., Schmitt, R., and Mages, W.:** An engineered *Streptomyces hygroscopicus aph 7"* gene mediates dominant resistance against hygromycin B in *Chlamydomonas reinhardtii*, *Protist*, **153**, 401-412 (2002).
13. **Sizova, I., Fuhrmann, M., and Hegemann, P.:** A *Streptomyces rimosus aphVIII* gene coding for a new type phosphotransferase provides stable antibiotic resistance to *Chlamydomonas reinhardtii*, *Gene*, **277**, 221-229 (2001).
14. **Stevens, D., Roehaix, J-D., and Purton, S.:** The bacterial phleomycin resistance gene *ble* as a dominant selectable marker in *Chlamydomonas*, *Mol. Gen Genet.*, **251**, 23-30 (1996).

- 1 15. **Sanger, F., Nicklen, S., and Coulson, A.:** DNA sequencing with chain-terminating
2 inhibitors, *Proc. Natl. Acad. Sci. USA*, **74**, 5463–5467 (1977).
3
- 4 16. **Gorman D. and Levine, R.:** Cytochrome f and plastocyanin: their sequence in the
5 photosynthetic electron transport chain of *Chlamydomonas reinhardtii*, *Proc. Natl. Acad.*
6 *Sci. U S A*, **54**, 1665–1669 (1965).
7
- 8 17. **Shimogawara, K., Fujiwara, S., Grossman, A., and Usuda, H.:** High-efficiency
9 transformation of *Chlamydomonas reinhardtii* by electroporation, *Genetics*, **148**,
10 1821-1828 (1998).
11
- 12 18. **Schroda, M., Vallon, O., Whitelegge, J., Beck, C., and Wollman, F.:** The
13 chloroplastic GrpE homolog of *Chlamydomonas*: two isoforms generated by differential
14 splicing, *Plant Cell*, **13**, 2823-2839 (2001).
15
- 16 19. **Yamano, T., Iguchi, H., and Fukuzawa, H.:** Rapid transformation of
17 *Chlamydomonas reinhardtii* without cell-wall removal, *J. Biosci. Bioeng.*, **115**, 691-694
18 (2013).
19
- 20 20. **Lauersen, K., Berger, H., Mussnug, J., and Kruse, O.:** Efficient recombinant
21 protein production and secretion from nuclear transgenes in *Chlamydomonas reinhardtii*,
22 *Journal of Biotechnology*, **167**, 101-110 (2012).
23
- 24 21. **Heitzer, M. and Zschoernig, B.:** Construction of modular tandem expression

vectors for the green alga *Chlamydomonas reinhardtii* using the Cre/lox-system,
Bio.Techniques, **43**, 324-328 (2007).

Figure legends

Figure 1. Identification of DXS1 and SQS protein-positive transformants by Western blotting. The DXS1- and SQS-positive transformants in the CC-124 strain (A) and SQS-positive transformants in the UVM4 (B) and UVM11 (C) strains are shown. The bands were quantified using the Quantity One 1-D Analysis Program (Bio-Rad). The band intensities are shown as relative intensity units in their respective gels. Band intensity values greater than 11 were defined as strongly expressing transformants in this study. Asterisk: transformants that expressed the target proteins at high levels.

Figure %

Table 1. Expression of *SQS* cDNA cassettes in various *Chlamydomonas reinhardtii* strains.

Gene name	Host strain	Cotransformation ratio ^a	Western blot-positive ratio ^b	Highly expressing transformant ratio ^c relative to Western blot-positive ratio
<i>SQS</i>	CC-124 (WT)	109/210 (51.9%)	5/109 (4.6%)	1/5 (20.0%)
	UVM 4	52/87 (59.8%)	11/52 (21.2%)	5/11 (45.5%)
		51/87 (58.6%)		
	UVM 11		9/51 (17.6%)	4/9 (44.4%)

Asterisk: wild type.

^aNumber of PCR-positive transformants/number of transformants analyzed.

^bNumber of Western blot-positive transformants/number of PCR-positive transformants.

^cNumber of high-protein accumulation transformants/number of Western blot-positive transformants.