1 **ORIGINAL PAPER**

2
3 **Molecular Phylogeny of the Widely Distributed Marine Protists, Phaeodaria**
4 **(Rhizaria, Cercozoa)**

5
6 Yasuhide Nakamura^{a,1}, Ichiro Imai^a, Atsushi Yamaguchi^a, Akihiro Tuji^b, Fabrice Not^c,
7 and Noritoshi Suzuki^d

8
9 ^aPlankton Laboratory, Graduate School of Fisheries Sciences, Hokkaido University,
10 Hakodate, Hokkaido 041–8611, Japan

11 ^bDepartment of Botany, National Museum of Nature and Science, Tsukuba 305–0005,
12 Japan

13 ^cCNRS, UMR 7144 & Université Pierre et Marie Curie, Station Biologique de Roscoff,
14 Equipe EPPO - Evolution du Plancton et PaléoOcéans, Place Georges Teissier, 29682
15 Roscoff, France

16 ^dInstitute of Geology and Paleontology, Graduate School of Science, Tohoku University,
17 Sendai 980–8578, Japan

18
19
20 Submitted January 1, 2015; Accepted April 27, 2015

21 Monitoring Editor: David Moreira

22
23 **Running title:** Molecular Phylogeny of Phaeodaria

24
25
26
27 ¹Corresponding author; fax +81 138 40 5542
28 e-mail y.nakamura@fish.hokudai.ac.jp (Y. Nakamura).

Phaeodarians are a group of widely distributed marine cercozoans. These plankton organisms can exhibit a large biomass in the environment and are supposed to play an important role in marine ecosystems and in material cycles in the ocean. Accurate knowledge of phaeodarian classification is thus necessary to better understand marine biology, however, phylogenetic information on Phaeodaria is limited. The present study analyzed 18S rDNA sequences encompassing all existing phaeodarian orders, to clarify their phylogenetic relationships and improve their taxonomic classification. The monophyly of Phaeodaria was confirmed and strongly supported by phylogenetic analysis with a larger data set than in previous studies. The phaeodarian clade contained 11 subclades which generally did not correspond to the families and orders of the current classification system. Two families (Challengeriidae and Aulosphaeridae) and two orders (Phaeogromida and Phaeocalpida) are possibly polyphyletic or paraphyletic, and consequently the classification needs to be revised at both the family and order levels by integrative taxonomy approaches. Two morphological criteria, 1) the scleracoma type and 2) its surface structure, could be useful markers at the family level.

Key words: Phaeodaria; Cercozoa; Rhizaria; 18S rDNA; single-cell PCR.

Introduction

Phaeodarians are marine unicellular protists. This group is defined by 1) the presence of a “central capsule” containing one or several nuclei, 2) a mass of brown aggregated particles named “phaeodium”, and 3) a hollow siliceous skeleton called “scleracoma”, in most species (Howe et al. 2011; Nakamura and Suzuki 2015; Takahashi and Anderson 2002). The size of the scleracoma varies with the different families and ranges from several hundred micrometers (e.g. Challengeriidae) to a few millimeters (e.g. Tuscaroridae). The type of the scleracoma and its surface structure are important taxonomic characters to classify these organisms at the order and family levels. Phaeodarians have long been considered as a member of Radiolaria but based on molecular analyses (Nikolaev et al. 2004; Polet et al. 2004; Sierra et al. 2013), they are now classified as a subclass in the class Thecofilosea of the phylum Cercozoa (Adl et al. 2012; Howe et al. 2011).

Phaeodarians are heterotrophic plankton organisms, which are distributed in the world ocean from the surface to the deep sea (Nakamura and Suzuki 2015). Despite

1 their wide range of distribution, this plankton group has attracted little attention from
2 marine biologists (Nakamura et al. 2013). Their abundance has been very likely
3 underestimated, because their scleracoma is fragile and breaks easily when samples are
4 collected by plankton nets. Yet, careful examination of the zooplankton community
5 revealed that these protists can be numerous and their high abundance has occasionally
6 been reported. For instance, the vertical flux of phaeodarians was higher than that of
7 polycystine radiolarians in the Panama Basin (Takahashi and Honjo 1983) and in the
8 California coast (Gowing and Coale 1989). The combined biomass of the
9 Aulosphaeridae, Sagosphaeridae, Aulacanthidae and Coelodendridae was estimated to
10 contribute 2.7– 13.7% of the total metazoan biomass in the 150– 1000 m layer of the
11 western North Pacific (Steinberg et al. 2008). The proportion of an Aulacanthidae
12 species, *Aulographis japonica*, with respect to the total zooplankton biomass is 22.3%
13 in 250– 3000 m layer in the Sea of Japan, and this percentage is the second largest,
14 following that of copepods (Nakamura et al. 2013). Considering that they harbor a
15 siliceous skeleton and occur occasionally at high biomass, this plankton group can play
16 an important role in ecosystems locally and have a significant impact on the silica cycle
17 of the ocean. Their importance in the carbon cycle is also reported from the Northeast
18 Atlantic (Lampitt et al. 2009). Cercozoans, including phaeodarians, have recently been
19 considered important players in the material cycles and food webs of the ocean (Howe
20 et al. 2011). Research on environmental DNA revealed that Cercozoa represented a high
21 percentage of the 18S rDNA sequences obtained from sea-floor sediments of the Arctic
22 and the Southern Ocean (Pawlowski et al. 2011). Consequently, improving our
23 knowledge on phaeodarians will be important for the entire field of marine biology.
24 Accurate knowledge of phaeodarian classification is critical to better assess their genetic
25 diversity and ecology. In the comprehensive systematics framework established by
26 Haeckel (1887), phaeodarian species were grouped into families and orders according to
27 their morphological similarities. Since then, some authors amended this classification,
28 but the relationship between the families remains uncertain (Cachon and Cachon 1985;
29 Campbell 1954; Kling and Boltovskoy 1999; Takahashi and Anderson 2002) and has
30 never been examined from the molecular point of view. Comprehensive molecular
31 studies on phaeodarians are difficult to conduct. They cannot be cultured, and therefore
32 their DNA has to be extracted from individual specimens collected and sorted from the
33 environment. Some phaeodarian species, however, live in the deep sea, and it is hard to
34 obtain specimens. In addition, efficient phaeodarian-specific primers do not exist,
35 increasing the risks of amplifying and sequencing contaminating organisms attached to
36 or ingested by the Phaeodaria.

Yet, an integrative taxonomy approach merging both morphological and molecular information has recently been applied successively to other unicellular and shell-bearing rhizarian taxa such as Polycystines, Acantharia and Foraminifera (e.g. Decelle et al. 2012; Kunitomo et al. 2006; Pawlowski et al. 2013). Such an approach, highlighting match and mismatch between morphology and phylogenetic analysis, contributed to an overall better understanding of the classification and evolutionary patterns among these groups.

The information on phaeodarian 18S ribosomal DNA currently exists for only 7 out of ca. 200 species, 18 families and 7 orders recognized in the latest phaeodarian classification (Nakamura and Suzuki 2015; Takahashi and Anderson 2002), and discrepancies between morphological classification and analysis of these partial 18S rDNA were pointed out (Nakamura et al. 2013). Here, we designed new phaeodarian-specific primers and analyzed the whole 18S rDNA sequence of extant single-cell specimens of phaeodarian species, covering all orders recognized in the current classification.

Results

A total of 39 phaeodarian specimens belonging to 26 distinct taxa (essentially distinct species) were collected from 0 m to 3 000 m depth, and their sequences were obtained by single-cell PCR. Detailed information and micrographs of analyzed specimens are shown in Table 1 and Figure 1, respectively.

The sequences from all phaeodarian specimens formed a single clade far separated from other Thecofilosea taxa (Cryomonadida, Ebriacea and *Pseudodiffugia*), and the monophyly of this clade was strongly supported by bootstrap values and BPP (Fig. 2). The phaeodarian clade consisted of 11 subclades (A–K) defined based on support values and morphological features (Fig. 4). Subclade A contains the highest number of individuals (n = 18). Subclades A, E, F, G, H and K were composed of sequences from distinct sampling locations. For instance, subclade E contains the specimens from the Mediterranean Sea (MS), the Sea of Japan (SJ) and the eastern North Pacific (ENP).

The 11 subclades and the families classified by Takahashi and Anderson (2002) generally did not match well, and disagreement was seen for the families Aulosphaeridae, Sagosphaeridae, Challengeriidae and Aulacanthidae (Fig. 3). *Sagoscena* sp. 1 (ENP3) of Sagosphaeridae formed the subclade E along with two

Aulosphaeridae species, being nested one into each other. Challengeriidae and Aulacanthidae were scattered across the phylogenetic tree. Most of the challengeriid phaeodarians appeared in subclade A, but two specimens of *Challengeron willemoesii* Haeckel (ECS6 and ECS7) formed the subclade B clearly distinct from subclade A with high support values (1.0/100/100, Fig. 3). The sequences of *C. diodon* (Haeckel) (ECS4 and NA (AB218765)) belonged to subclade H, distant from subclade A, together with the *Medusetta parthenopaea* Borgert (MS2) of Medusettidae (Fig. 3). Two Aulacanthidae species, *Aulographis japonica* Nakamura, Tuji and Suzuki (SJ1, SJ4 and SJ (AB820365)) and *Auloceros arborescens* Haeckel (ENP6) formed a single subclade K, whereas *Aulacantha scolymantha* Haeckel (MS1 and MS (AY266294)), which also belongs to Aulacanthidae, composed subclade I whose exact position compared to other subclades is unclear.

The orders of the current classification did not correspond to the phylogenetic analysis overall (Fig. 3). The order Phaeogromida was scattered across subclades A, B and F, and the order Phaeocystida has representatives in subclades I and K. The order Phaeosphaerida, composed of two families Sagosphaeridae and Aulosphaeridae (subclade E), turned out to be nested within the order Phaeocalpida (subclades C and D).

Discussion

Comparison Between the Phylogenetic Tree and the Current Classification

Previous studies analyzed partial 18S rDNA of phaeodarian species (n=6) belonging to 5 distinct families and orders, and reported that their sequences formed a monophyletic clade outside of Radiolaria but within cercozoans (Polet et al. 2004; Yuasa et al. 2006). In this study, the monophyly of Phaeodaria within Cercozoa is further confirmed based on a larger data set, including 29 distinct species distributed over all 7 orders. Phaeodarians are now classified as a member of the class Thecofilosea, together with Cryomonadida, Ebriacea and *Pseudodiffugia* (Adl et al. 2012; Howe et al. 2011). However, these Thecofilosea taxa did not form a monophyletic group in the 18S rDNA tree (Fig. 2), and their phylogenetic relationship could not be confirmed in the present study. For better clarifying the phylogenetic relationship between phaeodarians and other cercozoans, it would be necessary to add more taxa whose 18S rDNA sequences are not yet available (e.g. Ventricleftida).

The 11 subclades in the 18S rDNA phylogenetic reconstruction did not well

correspond to the families and the orders in the current classification system (Fig. 3). The family Challengeriidae is possibly a polyphyletic group, and the family Aulosphaeridae would be a paraphyletic group. As for ordinal level categorization, Phaeogromida turned out to be polyphyletic. The branching pattern of subclades C, D and E suggests that the order Phaeocalpida is a paraphyletic group. Consequently, the current classification system is not supported by the 18S rDNA phylogeny and needs to be reconsidered.

Comparison Between the Phylogenetic Tree and the Morphological Characters

Previous studies classified phaeodarian families according to morphological characters, especially based on 1) the scleracoma type and 2) the surface structure of the test (Cachon and Cachon 1985; Campbell 1954; Haeckel 1887; Kling and Boltovskoy 1999; Takahashi and Anderson 2000). The 11 subclades were generally congruent with these two morphological criteria in this study (Fig. 3). The scleracoma of subclades A, B, C, D, F and H formed relatively firm “test”, but displaying distinction between each other: cape-shaped or ovate test possessing smooth or small-pored surface (A and B), spherical test with “tubular spines” and the surface of “trizonal meshwork” (C), test of smooth surface with short “oral spines” and long “aboral spines” (D), campanulate test possessing developed “peristome” with “arches” (F), and test composed of two “valves” with large-pored surface (H). The scleracoma of subclade E corresponds to a sphere of geometric meshwork called “lattice-shell”. The phaeodarians of subclade G possess scleracoma composed of “inner shell” and numerous branch-like structure called “styles”. The specimens included in subclades I and K have scleracoma called “spherical veil”. Specimens belonging to subclade J do not possess scleracoma (Figs 1, 3).

Thus, the scleracoma type and the surface structure of the test would be valuable characters for the family-level classification. Future studies should rearrange phaeodarian families and orders according to these two criteria and the branching pattern of the phylogenetic tree. The combination of both detailed morphological examination and molecular phylogeny helped identifying new characters for the taxonomic classification of Phaeodaria. Such an integrative approach has already been proven valuable for a number of taxa, including some among Rhizaria (e.g. Acantharia, Decelle et al. 2012).

Taxonomical Consideration on the Families Challengeriidae and Medusettidae

1 The subclade A contains 18 specimens belonging to 3 different challengeriid genera:
2 *Challengeron*, *Protocystis* and *Entocannula*. These 3 genera were erected according to
3 the difference in their morphological characteristics. Phaeodarians of these genera,
4 however, did not make subclades clearly distinct from each other, and there is a
5 possibility that the current classification within the family Challengeriidae is artificial.
6 In order to further clarify their intra-family relationship, analyses using a higher
7 resolution genetic marker regions such as the D1 and D2 regions of the 28S rDNA or
8 ITS are necessary.

9 The genus *Challengeron* contains ca. 13 species among which 5, including the
10 type species *C. bethelli* (Murray) (WNP6), were analyzed in this study (Campbell 1954).
11 The scleracoma of *C. willemoesii* (ECS6 and ECS7) forms a firm “test”, similar to other
12 challengeriid phaeodarians composing subclade A (Fig. 1). This species has a test with
13 the surface made of a regularly arranged amphora-shaped structure, which is common to
14 other challengeriid species, except *C. diodon* (ECS4) (Takahashi 1991). Consequently,
15 the surface structure does not distinguish this species from other challengeriiids
16 examined in this study. *Challengeron willemoesii* has a developed “peristome” with four
17 “oral spines” (Fig. 1). This complex peristome structure is not seen in other
18 challengeriiids belonging to subclade A, and this character may be a morphological
19 criterion distinguishing this species from other challengeriiids.

20 The fact that *C. diodon* (ECS4) appeared in subclade F together with *Medusetta*
21 *parthenopaea* (MS2) of the family Medusettidae suggests that the former is
22 phylogenetically closer to the medusettids than other challengeriiids. Morphological
23 differences of *C. diodon* from other challengeriiids have previously been noticed: 1) a
24 dimpled surface; 2) clusters of small pores associated with individual amphorae (inner
25 shell surface); and 3) thin and delicate amphorae (Takahashi 1991). This author also
26 mentioned that the overall appearance of this species was closer to that of medusettid.
27 The two species composing subclade F have one “aboral spine” and the developed
28 “peristome” with “arches” in common (Fig. 1). These structures could also be important
29 new characteristic features to distinguish this subgroup. Considering the morphological
30 features and the position in the phylogenetic tree, these two species would better to be
31 included in the family Medusettidae.

32 33 Wide Distribution of Phaeodarians

34 Specimens of phaeodarian were collected from both deep sea and surface (e.g. MS1–4,
35 Table 1), emphasizing their wide vertical distribution as previously reported (Nakamura
36 and Suzuki 2015). Based on sampling location, *C. diodon* is likely a cosmopolitan

species. The two sequences of *C. diodon* within subclade F (Fig. 3) have been collected at distinct locations, Sogndalsfjord, western Norway (NA [AB218765], Yuasa et al. 2006) and the East China Sea (ECS4, Table 1). This species was also found in the North Sea, the Labrador Sea, the South Atlantic and the Mediterranean Sea (Borgert 1901).

The present study provides a strong morphogenetic framework for phaeodarian classification, and will be a valuable tool for accurate interpretation of ongoing environmental diversity surveys based on DNA sequencing. This morphogenetic approach and the single-cell PCR method can further elucidate the intra- and inter-group relationships of phaeodarians and other protists by combining other biological information such as distribution or physiology.

Methods

Sampling and identification: Plankton samples were collected in 2012 and 2013 from 14 stations located in 5 seas across the northern hemisphere: the western and eastern North Pacific Ocean, the Sea of Japan, the East China Sea and the Mediterranean Sea (Supplementary Material Table 1, Supplementary Material Fig. S1). Phaeodarian individuals were isolated manually from the samples under a stereomicroscope or inverted microscope, as soon as possible after the sampling. Isolated specimens were put into wells of cell culture plates and incubated for several hours to allow self-cleaning. The specimens were carefully identified according to their morphological characters under an inverted stereomicroscope and photographed with a digital camera (DIGITAL SIGHT DS-Ri1, Nikon, Japan). For identification, all the documents concerning phaeodarian taxonomy were examined, and a taxonomic database of synonyms of Phaeodaria was constructed. The specimens were then individually preserved in tubes filled with approximately 2.0 mL of 99.9% ethanol. The tubes containing the specimens were stored at 4 °C. Each specimen was labeled according to its sampling station (Table 1 and Fig. 1).

Single-cell PCR: After confirming that there were no other organisms attached to the cell surface, each specimen was put into 50 µL of guanidine-containing extraction buffer (GITC buffer, Decelle et al. 2012) and preserved in -80 °C overnight. Some large specimens (e.g. Aulosphaeridae) were dissected by a sterilized scalpel, and only their “central capsules” (protoplasmic body with nucleus) were transferred into the GITC buffer. The specimens were heated at 70 °C for 20 min to break the wall of their central capsules. The extracted DNA was purified with a chemagic DNA Plant kit

(PerkinElmer chemagen, Germany). Single-cell PCR was conducted to amplify 18S rDNA with a total reaction volume of 25 μ L. A total of 9 primers were used, including 4 phaeodarian-specific primers designed in this study: Phaeo F3a, Phaeo11F, PhaeoF4-2 and Phaeo880F (Table 2). PCR reactions were performed using the following protocol: initial denaturation at 95 °C for 10 min, 40 cycles at 95 °C for 20 sec, 58 °C for 15 sec and 72 °C for 120 sec with a final extension at 72 °C for 7 min. The amplified PCR products were purified with AMPure XP Kit (Beckman Coulter, USA). Sequencing reactions were conducted using an ABI PRISM 3130xl Genetic Analyzer (ABI, U.S.A.).

Phylogenetic analysis: The sequences obtained were assembled using ChromasPro (Technelysium Pty Ltd, Australia), and the alignments were checked manually. The sequences of 6 phaeodarian species, 57 cercozoans were also retrieved from the Basic Local Alignment Search Tool (BLAST) at NCBI (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) (Supplementary Material Table 2), and all the sequences were then aligned using MEGA 5 (Tamura et al. 2011). Phylogenetic trees were inferred by maximum-likelihood (ML), neighbor-joining (NJ) and Bayesian analysis methods. For the ML and NJ trees, the General Time Reversible (GTR) model plus Gamma with the shape parameter for among-site rate variation (G) was selected based on the lowest Bayesian Information Criterion (BIC) score, and the substitution nucleotide matrix parameters were calculated. The Bootstrap values (Felsenstein 1985) were estimated based on 1000 pseudo-replicates. GTR model with invariant sites and the gamma distributed model (IG) were selected for the Bayesian analysis, and the Bayesian posterior probabilities (BPP) were calculated with Mr Bayes version 3.2.2. (Ronquist and Huelsenbeck 2003). The Markov Chain Monte Carlo chains ran for 3.0×10^7 generations, sampling every 1000 generations. The first 1.0×10^6 generations were discarded as burn-in, checking by a program Tracer version 1.6 (Rambaut and Drummond 2009). The remaining data reaching the steady state were used for building the consensus tree, and the tree was visualized by Fig Tree version 1.4.0. (Rambaut 2012).

Acknowledgements

We thank the captains and crews of the T/S *Oshoro-maru* (Hokkaido University), T/S *Ushio-maru* (Hokkaido University) and T/S *Toyoshio-maru* (Hiroshima University) for their assistance in sampling. We also thank the members of the oceanic plankton group

1 of Station Biologique de Roscoff for their kind technical advice. Appreciation is also
2 due to the members of Plankton laboratory of Hokkaido University for their help in
3 analysis. The comments of two anonymous reviewers and the editor also greatly
4 improved the manuscript. This study was financially supported by Grant-in-Aid for
5 JSPS Fellows (no. 26-2889, Y. Nakamura), by the Cooperative Research Project with
6 Centre National de la Recherche Scientifique (CNRS), “Morpho-molecular Diversity
7 Assessment of Ecologically, Evolutionally, and Geologically Relevant Marine Plankton
8 (Radiolaria)” by the Strategic International Research Cooperative Program hosted by
9 the Japan Science and Technology Agency (JST) (N. Suzuki and F. Not), and by
10 “Integrated Research on Biodiversity of Interspecies Relationships” of the National
11 Museum of Nature and Science (A. Tuji).

References

- Adl SM, Simpson AG, Lane CE, Lukeš J, Bass D, Bowser SS, Brown MW, Burki F, Dunthorn M, Hampl V, Heiss A, Hoppenrath M, Lara E, Le Gall L, Lynn DH, McManus H, Mitchell EA, Mozley-Stanridge SE, Parfrey LW, Pawlowski J, Rueckert S, Shadwick RS, Schoch CL, Smirnov A, Spiegel FW** (2012) The revised classification of eukaryotes. *J Eukaryot Microbiol* **59**:429–493
- Borgert A** (1901) Die nordischen Tripyleen-Arten. *Nord Plankton* **15**:1–52
- Cachon J, Cachon M** (1985) 2. Class Polycystinea. In Lee JJ, Hutner SH, Bovee EC (eds) *Illustrated Guide to the Protozoa*. Allen Press, Lawrence, pp 283–295
- Campbell AS** (1954) Radiolaria. In Moore RC (ed) *Treatise on Invertebrate Paleontology*, Part D, Protista 3. Protozoa (Chiefly Radiolaria and Tintinnina). Geological Society of America and University of Kansas Press, Kansas, pp D1–D163
- Decelle J, Suzuki N, Mahé F, de Vargas C, Not F** (2012) Molecular phylogeny and morphological evolution of the Acantharia (Radiolaria). *Protist* **163**:435–450
- Felsenstein J** (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**:783–791
- Gowing MM, Coale SL** (1989) Fluxes of living radiolarians and their skeletons along a northeast Pacific transect from coastal upwelling to open ocean waters. *Deep-Sea Res* **36**:550–565
- Haeckel E** (1887) Report on the Radiolaria collected by H.M.S. Challenger during the years 1873–1876. *Rep Sci Res Voyage HMS Challenger 1873–1876*, Zool **18**:1–1803
- Howe AT, Bass D, Scoble JM, Lewis R, Vickerman, K Arndt H, Cavalier-Smith T** (2011) Novel cultured protists identify deep-branching environmental DNA clades of Cercozoa: new genera *Tremula*, *Micrometopion*, *Minimassisteria*, *Nudifila*, *Peregrinia*. *Protist* **162**:332–372
- Kling SA, Boltovskoy D** (1999) Radiolaria Phaeodaria In Boltovskoy D (ed) *South Atlantic Zooplankton*. Backhuys Publishers, Leiden, pp 231–264

- Kunitomo Y, Sarashina I, Iijima M, Endo K, Sashida K** (2006) Molecular phylogeny of acantharian and polycystine radiolarians based on ribosomal DNA sequences, and some comparisons with data from the fossil record. *Eur J Protistol* **43**:143–153
- Lampitt RS, Salter I, Johns D** (2009) Radiolaria: Major exporters of organic carbon to the deep ocean. *Global Biogeochem Cycles* **23**, GB1010
- Nakamura Y, Suzuki N** (2015) Chapter 10 Phaeodaria: Diverse marine cercozoans of world-wide distribution. In: Ohtsuka S, Suzuki T, Horiguchi T, Suzuki N, Not F (eds.), *Marine Protists Diversity and Dynamics*, Springer-Verlag, Berlin (in press)
- Nakamura Y, Imai I, Yamaguchi A, Tuji A, Suzuki N** (2013) *Aulographis japonica* sp. nov. (Phaeodaria, Aulacanthida, Aulacanthidae), an abundant zooplankton in the deep sea of the Sea of Japan. *Plankton Benthos Res* **8**:107–115
- Nikolaev SI, Berney C, Fahrni JF, Bolivar I, Polet S, Mylnikov AP, Aleshin VV, Petrov NB, Pawlowski J** (2004) The twilight of Heliozoa and rise of Rhizaria, an emerging supergroup of amoeboid eukaryotes. *Proc Natl Acad Sci USA* **101**:8066–8071
- Pawlowski J, Holzmann M, Tyszka J** (2013) New supraordinal classification of Foraminifera: Molecules meet morphology. *Mar Micropaleontol* **100**:1–10
- Pawlowski J, Christen R, Lecroq B, Bachar D, Shahbazkia HR, Amaral-Zettler L, Guillou L** (2011) Eukaryotic richness in the abyss: insights from pyrotag sequencing. *PLoS ONE* **6**:e18169
- Polet S, Berney C, Fahrni J, Pawlowski J** (2004) Small-subunit ribosomal RNA gene sequences of Phaeodarea challenge the monophyly of Haeckel's Radiolaria. *Protist* **155**:53–63
- Rambaut A** (2012) FigTree version 1.4.0. <http://tree.bio.ed.ac.uk/software/figtree/>
- Rambaut A, Drummond AJ** (2009) Tracer version 1.6. <http://beast.bio.ed.ac.uk/Tracer>
- Ronquist F, Huelsenbeck JP** (2003) MrBayes version 3.0: Bayesian phylogenetic inference under mixed models. *Bioinformatics* **19**:1572–1574

- 1
2 **Sierra R, Matz MV, Aglyamova G, Pillet L, Decelle J, Not F, de Vargas C, Pawlowski J**
3 (2013) Deep relationships of Rhizaria revealed by phylogenomics: A farewell to Haeckel's
4 Radiolaria. *Mol Phylogenet Evol* **67**:53–59
5
6 **Steinberg DK, Cope JS, Wilson SE, Kobari T** (2008) A comparison of mesopelagic
7 mesozooplankton community structure in the subtropical and subarctic North Pacific Ocean.
8 *Deep-Sea Res II* **55**:1615–1635
9
10 **Takahashi K** (1991) Radiolaria: Flux, Ecology, and Taxonomy in the Pacific and Atlantic.
11 Ocean Biocoenosis Series 3. Woods Hole Oceanographic Institution Press, Woods Hole, 303 p
12
13 **Takahashi K, Anderson OR** (2002) Class Phaeodaria. In Lee JJ, Leedale GF and Brandbury P
14 (eds) *Illustrated Guide to the Protozoa*. 2nd ed, Society of Protozoologists, Kansas, pp 981–994
15
16 **Takahashi K, Honjo S** (1983) Radiolarian skeletons: size, weight, sinking speed, and residence
17 time in tropical pelagic oceans. *Deep-Sea Res* **30**:543–568
18
19 **Tamura K, Peterson D, Peterson N, Steker G, Nei M, Kumar S** (2011) MEGA5: Molecular
20 Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary Distance, and
21 Maximum Parsimony Methods. *Mol Biol Evol* **28**:2731–2739
22
23 **Yuasa T, Takahashi O, Dolven JK, Mayama S, Matsuoka A, Honda D, Bjørklund KR**
24 (2006) Phylogenetic position of the small solitary phaeodarians (Radiolaria) based on 18S
25 rDNA sequences by single cell PCR analysis. *Mar Micropaleontol* **59**:104–114
26

Figure legends

Figure 1. Light micrographs of phaeodarian specimens analyzed in this study. The information on each specimen is presented in Table 2. cc: central capsule; ph: phaeodium; sc: scleracoma.

Figure 2. Phylogenetic tree of phaeodarians and other cercozoans based on 18S rDNA alignments and the Maximum-likelihood (ML) method. Note that the phaeodarian clade is presented with 11 subclades (A–K) detected according to the support values and morphological features. The details of this clade are shown in Figure 3. Numbers at nodes indicate Bayesian posterior probabilities (BPP) and bootstrap support values of Neighbor Joining (NJ) and ML methods (BPP/NJ/ML). Only values higher than 0.5 pp and 50% are shown.

Figure 3. The phaeodarian clade in the phylogenetic tree inferred by the Maximum-likelihood (ML) method, based on 46 sequences of 18S rDNA. The outgroups are shown in Figure 2. Note that the 11 subclades (A–K) are detected based on the support values and morphological features. Each specimen is coded according to its sampling station as shown in Table 2, and species retrieved from NCBI database are presented with their accession numbers. Numbers at nodes indicate Bayesian posterior probabilities (BPP) and bootstrap support values of Neighbor Joining (NJ) and ML method (BPP/NJ/ML). Only values higher than 0.85 pp and 50% are shown. The families and orders are based on Takahashi and Anderson (2002).

Figure 1a

Figure 1b

Figure 2

Figure 3

Table 1. List of phaeodarian sequences used for the phylogenetic analysis in this study. the specimens are coded according to the sampling station: the western North Pacific (WNP); the eastern north Pacific (ENP); the East China Sea (ECS); the Sea of Japan (SJ); the Mediterranean Sea (MS).

Taxon	Accession no.	Specimen name	Sampling station	Depth range (m)	Sampling date
<i>Protocystis vicina</i>	AB998884	WNP1	WN1	250–500	Aug. 2013
<i>Protocystis vicina</i>	AB998885	WNP2	WN1	250–500	Aug. 2013
Challengeriidae sp. 1	AB998886	WNP3	WN2	100–150	Aug. 2013
<i>Porospathis holostoma</i>	AB998887	WNP4	WN2	150–250	Aug. 2013
Challengeriidae sp. 2	AB998888	WNP5	WN2	500–750	Aug. 2013
<i>Challengeron bethelli</i>	AB998889	WNP6	WN2	500–750	Aug. 2013
<i>Protocystis thomsoni</i>	AB998890	WNP7	WN2	500–750	Aug. 2013
<i>Challengeron tizardi</i>	AB998891	WNP8	WN2	750–1000	Aug. 2013
<i>Challengeron tizardi</i>	AB998892	WNP9	WN2	1500–2000	Aug. 2013
<i>Challengeron tizardi</i>	AB998893	WNP10	WN2	1500–2000	Aug. 2013
<i>Entocannula infundibulum</i>	AB998894	WNP11	WN2	1500–2000	Aug. 2013
<i>Porospathis holostoma</i>	AB998895	WNP12	WN2	1500–2000	Aug. 2013
<i>Protocystis harstoni</i>	AB998896	WNP13	WN2	1500–2000	Aug. 2013
<i>Protocystis murrayi</i>	AB998897	WNP14	WN2	1500–2000	Aug. 2013
<i>Tuscaretta belknapi</i>	AB998898	ENP1	EN1	0–300	Aug. 2012
<i>Coelodendrum furcatissimum</i>	AB998899	ENP2	EN2	0–1000	Aug. 2012
<i>Sagoscena</i> sp. 1	AB998900	ENP3	EN2	0–1000	Aug. 2012
<i>Tuscaretta belknapi</i>	AB998901	ENP4	EN3	0–1000	July 2012
<i>Tuscaretta</i> sp. 1	AB998902	ENP5	EN4	0–1000	July 2012
<i>Auloceros arborescens</i>	AB998903	ENP6	EN5	0–1000	July 2012
<i>Coelodendrum furcatissimum</i>	AB998904	ENP7	EN5	0–1000	July 2012
Conchariidae sp. 1	AB998905	ECS1	E1	0–150	May 2013
<i>Challengeron radians</i>	AB998906	ECS2	E2	200–500	May 2013
Conchariidae sp. 2	AB998907	ECS3	E2	200–500	May 2013
<i>Challengeron diodon</i>	AB998908	ECS4	E3	650–800	May 2012
<i>Challengeron radians</i>	AB998909	ECS5	E3	650–800	May 2012
<i>Challengeron willemoesii</i>	AB998910	ECS6	E3	650–800	May 2012
<i>Challengeron willemoesii</i>	AB998911	ECS7	E3	650–800	May 2012
<i>Protocystis xiphodon</i>	AB998912	ECS8	E4	0–150	May 2012
<i>Protocystis xiphodon</i>	AB998913	ECS9	E4	0–150	May 2012
<i>Aulographis japonica</i>	AB998914	SJ1	S1	250–750	Apr. 2013
<i>Aulosena</i> sp. 1	AB998915	SJ2	S1	250–750	Apr. 2013
<i>Protocystis vicina</i>	AB998916	SJ3	S1	250–750	Apr. 2013
<i>Aulographis japonica</i>	AB998917	SJ4	S2	250–750	Apr. 2012
<i>Aulosena</i> sp. 1	AB998918	SJ5	S3	500–750	Mar. 2013
<i>Aulacantha scolymantha</i>	AB998919	MS1	M1	0–40	Nov. 2012
<i>Medusetta parthenopaea</i>	AB998920	MS2	M1	0–40	Nov. 2012
<i>Phaeodina</i> sp. 1	AB998921	MS3	M1	0–40	Nov. 2012
<i>Phaeodina</i> sp. 1	AB998922	MS4	M1	0–40	Nov. 2012

Table 2. Sequences of the primers for 18S rDNA used in this study.

Primer name	Position*	Sequence 5'-3'	Direction	Specificity	Reference
Medlin A	5'-end	AAC CTG GTT GAT CCT GCC AGT	Forward	Universal	Medlin et al. (1988)
PhaeoF3a	288–319	TCA TTC AAA TTT CTG CCC TAT CAG CTW GAY GG	Forward	Phaeodaria	This study
NS2	546–566	GGC TGC TGG CAC CAG ACT TGC	Reverse	Universal	White et al. (1990)
Phaeo11F	559–583	CAG CAG CCG CKG TAA TTC CAG CTC C	Forward	Phaeodaria	This study
PhaeoF4-2	872–890	AKG GAT AGT TGG GGG TGC T	Forward	Phaeodaria	This study
Phaeo880F	872–893	AKG GAT RGT TGG GGG TGC TAG T	Forward	Phaeodaria	This study
Phaeo R	885–904	GGC CAY TRA ATA CTA GCA CC	Reverse	Phaeodaria	Nakamura et al. (2013)
SR6	1761–1777	TGT TAC GAC TTT TAC TT	Reverse	Universal	designed in Vilgalys lab, Duke University.
Medlin B	3'-end	CCT TCT GCA GGT TCA CCT AC	Reverse	Universal	Medlin et al. (1988)

* Position within 18S rDNA of *Protocystis xiphodon*.