Development and Assessment of Indices to Determine Stream Fish Vulnerability to Climate Change and Habitat Alteration

Nicholas A. Sievert^a, Craig P. Paukert^b, Yin-Phan Tsang^c, and Dana Infante^d

^a Corresponding Author

E-mail: NAS4tf@mail.missouri.edu

Missouri Cooperative Fish and Wildlife Research Unit

Department of Fisheries and Wildlife Sciences

302 Anheuser-Busch Natural Resources Building

University of Missouri

Columbia, MO 65211

United States

^b

E-mail: paukerc@missouri.edu

U.S. Geological Survey, Missouri Cooperative Fish and Wildlife Research Unit

Department of Fisheries and Wildlife Sciences

302 Anheuser-Busch Natural Resources Building

University of Missouri

Columbia, MO 65211

United States

^c

E-mail: tsangy@hawaii.edu

Department of Natural Resources and Environmental Management

University of Hawaii

Sherman 243

1910 East-West Road

Honolulu, HI 96822

United States

^d

E-mail: infanted@anr.msu.edu

Department of Fisheries and Wildlife

Michigan State University

Manly Miles Building, Suite 318

1405 South Harrison Road

East Lansing, MI 48823

United States

1 **ABSTRACT**

2 Understanding the future impacts of climate and land use change are critical for long-term biodiversity
3 conservation. We developed and compared two indices to assess the vulnerability of stream fish in Missouri, USA
4 based on species environmental tolerances, rarity, range size, dispersal ability and on the average connectivity of
5 the streams occupied by each species. These two indices differed in how environmental tolerance was classified
6 (i.e., vulnerability to habitat alteration, changes in stream temperature, and changes to flow regimes). Environmental
7 tolerance was classified based on measured species responses to habitat alteration, and extremes in stream
8 temperatures and flow conditions for one index, while environmental tolerance for the second index was based on
9 species' traits. The indices were compared to determine if vulnerability scores differed by index or state listing
10 status. We also evaluated the spatial distribution of species classified as vulnerable to habitat alteration, changes in
11 stream temperature, and change in flow regimes. Vulnerability scores were calculated for all 133 species with the
12 trait association index, while only 101 species were evaluated using the species response index, because 32 species
13 lacked data to analyze for a response. Scores from the trait association index were greater than the species response
14 index. This is likely due to the species response index's inability to evaluate many rare species, which generally had
15 high vulnerability scores for the trait association index. The indices were consistent in classifying vulnerability to
16 habitat alteration, but varied in their classification of vulnerability due to increases in stream temperature and
17 alterations to flow regimes, likely because extremes in current climate may not fully capture future conditions and
18 their influence on stream fish communities. Both indices showed higher mean vulnerability scores for listed species
19 than unlisted species, which provided a coarse measure of validation. Our indices classified species identified as
20 being in need of conservation by the state of Missouri as highly vulnerable. The distribution of vulnerable species in
21 Missouri showed consistent patterns between indices, with the more forest-dominated, groundwater fed streams in
22 the Ozark subregion generally having higher numbers and proportions of vulnerable species per site than
23 subregions that were agriculturally dominated with more overland flow. These results suggest that both indices will
24 identify similar habitats as conservation action targets even despite discrepancies in the classification of vulnerable
25 species. Our vulnerability assessment provides a framework that can be refined and used in other regions.

26

1
2
3
4
5
6
7

KEYWORDS

- 1) Climate Change
- 2) Vulnerability Assessment
- 3) Stream Fish
- 4) Traits
- 5) Freshwater Conservation

1. INTRODUCTION

Stream ecosystems have some of the most imperiled communities on Earth, and freshwater biodiversity is declining at a higher rate than most other taxa groups (Allan and Flecker 1993; Abell 2002; Dudgeon *et al.* 2006). Anthropogenic habitat alteration, changes in stream temperatures, and change in flow regimes will likely cause continuing declines in aquatic biota (Ricciardi and Rasmussen 1999; Dudgeon *et al.* 2006). Conservation actions to protect aquatic biodiversity are critical for preventing future biodiversity losses (Master *et al.* 1998). In order to plan for long-term biodiversity conservation, a better understanding of how impacts and threats, such as climate change and habitat alteration, affect aquatic species is needed. Conservation and management of stream fish species will require researchers and managers to identify which species are vulnerable, or likely to experience harm, under future conditions (Turner *et al.* 2003; Glick *et al.* 2011; Poff *et al.* 2012). An assessment of stream fish vulnerability requires the identification of species which are threatened by primary drivers of future stream fish declines such as habitat alteration, predicted changes in stream temperatures, and predicted change in flow regimes (Malmqvist and Rundle 2002; Poff *et al.* 2002).

Climate change, which is expected to increase stream temperatures and alter flow regimes, is one of the most significant threats facing stream fish (Eaton and Scheller 1996; Poff *et al.* 2002). In Missouri, USA, the average annual air temperature is expected to rise approximately 4° C by 2095 based on an intermediate (A1B) climate scenario (Girvetz *et al.* 2009). This is expected to increase stream temperature by an average of 3.6°C (Eaton and Scheller 1996), which may have a substantial effect on stream fishes. In Wisconsin, USA, a 3°C air temperature increase is predicted to result in loss of 343,034 km of stream habitat for cool- and cold-water fishes, which includes a species extirpation (Lyons *et al.* 2010). Studies have predicted substantial losses of cool- and cold-water stream habitats (15-50%) with a doubling of atmospheric CO₂ (Eaton and Scheller 1996; Mohseni *et al.* 2003). Warm-water streams in the Great Plains, USA, achieve maximum temperatures at or near the physiological limits of some resident fish species, and an increase in stream temperature of just a few degrees is predicted to result in local extirpation and extinction (Matthews and Zimmerman 1990). In addition, decreases in fitness based on changes in energetics and growth may be a non-lethal consequence of warming temperatures (Pease and Paukert 2014; Westhoff and Paukert 2014). These studies provide strong evidence that some stream fish species may decline or face local extirpations as stream temperatures warm.

Changes to flow regimes are also expected to have substantial impacts on stream fish (Poff *et al.* 2002). Predicted increased frequency and extremes of flood and droughts may lead to shifts in species composition and local species extirpations (Poff *et al.* 1997, 2002). In Missouri, mean annual precipitation is expected to increase only slightly (~10mm) between historic (2065-2100) and future (2080-2100) time periods, however this precipitation is expected to come in the form of heavier precipitation (~10mm increase on wet days) on fewer days (12 less wet days) (Girvetz *et al.* 2009). The variability of precipitation patterns is likely to increase, which may cause declines in species which exhibit equilibrium or periodic life history strategies (Poff *et al.* 2002; Olden and Kennard 2010; Mims and Olden 2012, 2013).

Habitat alteration caused by anthropogenic modifications including the conversion of land to agricultural or urban uses or the direct modification of streams and rivers from channelization, dredging, and damming often result in altered and degraded stream conditions and losses of aquatic biodiversity (Malmqvist and Rundle 2002; Allan 2004). Indices of biological integrity (IBIs) have identified many stream fish species as intolerant of habitat alteration (Esselman *et al.* 2011), and life history traits sensitive to the effects of habitat alterations such as lithophilic spawning, and benthic invertivory have been used as criteria for IBIs measuring habitat alteration (Berkman and Rabeni 1987; Barbour *et al.* 1999; Simon 1999).

A framework for the assessment of stream fish vulnerability (i.e., the extent to which a species is likely to be impacted by the cumulative effects of climate change and habitat alteration [Turner *et al.* 2003; Schnieder *et al.* 2007; Glick *et al.* 2011]) can be developed by associating the effects of climate change and habitat alteration with species or trait-specific impacts. Vulnerability is often determined using a framework that assesses a species' sensitivity, exposure, and adaptive capacity to threats (Turner *et al.* 2003; IPCC 2007; Glick *et al.* 2011; Poff *et al.* 2012; Staudinger *et al.* 2013). A number of vulnerability assessments have been developed using this framework (Bagne *et al.* 2011; Glick *et al.* 2011; Young *et al.* 2011). Two prominent vulnerability assessment tools (the System for Assessing Vulnerability of Species (SAVS); Bagne *et al.* 2011, and the NatureServe Climate Change Vulnerability Index (NSCCVI; Young *et al.* 2011) could not be applied to Missouri stream fish due to a lack of necessary data for evaluating many of the criteria these assessment techniques use. These assessments depend on models of current and future climate conditions which have not yet been developed for stream temperature and flow in Missouri. Additionally, these tools were designed for a wide array of taxa over large spatial scales, and they

1 depend on information which is largely unknown for stream fish species; examples include knowledge of a species'
2 reliance on interspecific interactions, measures of genetic variation, occurrence of bottlenecks in recent
3 evolutionary history, and phenological response to changing seasonal temperature and precipitation dynamics
4 (Bagne *et al.* 2011; Young *et al.* 2011).Currently available information is too limited to broadly apply these
5 assessment tools to stream fishes in Missouri.The inability to apply currently available vulnerability assessment
6 tools to determine stream fish vulnerability necessitates the development of a new methodology.

7 Poff *et al.*(2012) developed a framework for assessing the threat posed by climate change to freshwater
8 diversity.This framework is a function of three components: exposure to the flow and temperature conditions which
9 deviate from regional baselines; sensitivity of species based on intrinsic factors related to a species environmental
10 tolerance, dispersal ability, genetic adaptation, range, and population size; and habitat resilience or the level of
11 connectivity of habitat which provides opportunities for adaptation via dispersal.Although this framework deviates
12 from the definitions and structure presented by Glick *et al.*(2011), we believe that it provides a mechanism for
13 assessing stream fish vulnerability based on the information that is currently available, so we adapted it for our use.

14 Species environmental tolerances, which are often expressed as sensitivity in the vulnerability assessment
15 literature, specifically to habitat alteration, changes in stream temperature, and change to flow regimes, as well as
16 factors such as a species range, rarity, dispersal ability, and the hydrological connectivity of a species habitat can be
17 incorporated into this framework to create a method for assessing the vulnerability of stream fish species.Our
18 analysis of species vulnerability will focus on the sensitivity and habitat resilience components of the Poff *et*
19 *al.*(2012) framework as adequate information is not yet available to assess exposure in Missouri.

20 Species tolerance of habitat alteration, changes in flow regimes, and increasing stream temperatures have
21 been assessed using two different approaches; species trait associations (Angermeier 1995; Parent and Schriml
22 1995; Poff 1997; Olden *et al.* 2007, 2008; Culp *et al.* 2011; Mims and Olden 2013) and measured species responses
23 (Hering *et al.* 2006; Poff and Zimmerman 2010; Lyons *et al.* 2010).We developedtwo separate indices, one which
24 scores environmental tolerance based on traits and the other which scores based on species responses.The same
25 scoring framework was used for both indices, however the indices used different methods for classifying
26 environmental tolerance.The trait association approach to classifying environmental tolerance is based on traits
27 which have been linked to vulnerability to habitat alteration, changes to flow regimes, and increases in stream

1 temperature in peer-reviewed literature, and the species response approach is based on species-specific observations
2 of sensitivity to each of the environmental tolerance components.

3 Another aspect of the sensitivity component is species dispersal ability (Poff *et al.* 2012).The ability of a
4 fish species to disperse throughout a stream network is critical to the persistence of species faced by
5 threats(Albanese *et al.* 2009; Poff *et al.* 2012).Dispersal, a one-way movement from a site (Lidicker Jr. and
6 Stenseth 1992), may benefit species facing threats by allowing 1) for gene flow between subpopulations (Hanski
7 1998; Heggenes *et al.* 2006), 2) increased colonization of newly available habitats (Detenbeck *et al.* 1992; Albanese
8 *et al.* 2009), and 3) species to shift their ranges as environmental changes occur at the landscape scale (Matthews
9 and Zimmerman 1990; Tonn 1990).

10 Both range size and population size can influence the vulnerability of a species (Poff *et al.* 2012).Species
11 with smaller ranges are limited in their ability to withstand stochastic environmental and demographic fluctuations
12 and are therefore more vulnerable (Angermeier 1995).Species constrained to a small range are more likely to lack
13 access to refugia or a gradient of conditions necessary to cope with the impacts of threats.Rare species, even those
14 which are widely distributed across the landscape, may have increased vulnerability to both deterministic threats
15 such as habitat loss, as well as stochastic events such as invasions and epidemics (Mace *et al.* 2008).Therefore,
16 species with narrowly restricted distributions and rare species generally have high levels of vulnerability.

17 Habitat resilience, defined as the hydrological connectivity of the landscape, is another component of the
18 vulnerability assessment framework developed by Poff *et al.*(2012).A stream system with high levels of
19 connectivity provides species the opportunity to emigrate to suitable habitats as conditions change (Poff *et al.*
20 2012), while a system with high levels of fragmentation, often caused by the creation of barriers such as dams and
21 road crossings (Jackson and Marmulla 2001; Fagan *et al.* 2002), can limit the ability of species to cope with threats
22 causing populations to face an elevated risk of extinction (Fagan *et al.* 2002). Fragmentation caused by dams is
23 particularly problematic for lotic obligate species as the lentic habitat created upstream of dams is not accessible for
24 these species (Gillette *et al.* 2005).

25 Scientists and decision makers benefit from the ability to identify vulnerable species to make effective
26 decisions regarding the conservation of stream fish.The objectives of this study are to 1) classify species'
27 vulnerability based on environmental tolerance of habitat alteration, changing stream temperatures, and changes to

1 flow regimes, dispersal ability, restricted distribution, rarity, and habitat resilience, 2) compare the use of traits and
2 species responses to identify species environmental tolerance, 3) develop indices to measure stream fish
3 vulnerability using our species classifications, 4) determine whether our measures of vulnerability correlate with
4 listing status, and 5) analyze distribution patterns of vulnerable species in a large area (the ecologically diverse state
5 of Missouri USA). Our aim is to develop a framework for assessing stream fish vulnerability at coarse spatial scales
6 and demonstrate its applicability to Missouri's stream fish species.

7

8 **2. MATERIALS AND METHODS**

9 **2.1. Study area**

10 The state of Missouri is over 180,500 km² and is classified into three ecologically unique subregions: the
11 Central Plains, Ozarks, and Mississippi Alluvial Basin (MAB) (Figure 1; Pflieger 1970; Pflieger and Missouri
12 Department of Conservation 1989; Sowa *et al.* 2007). These subregions are characterized by major differences in
13 geology, landform, soils, land cover, and groundwater influence which create unique habitats and fish communities
14 in the streams of each area. The Central Plains, in northern Missouri, is dominated by open grassland and agriculture
15 (78%; Blodgett and Lea 2005) with streams associated with wide, gently sloping valleys which have a relatively
16 low gradient, high turbidity, and fine silt and sand substrates (Blodgett and Lea 2005; Sowa *et al.* 2007). The
17 Ozarks, in southern Missouri (except the southeastern corner), is a mix of forested areas (51.4%) and agriculture
18 (7%; Blodgett and Lea 2005; Sowa *et al.* 2007). Most of the region has rugged terrain with high relief and higher
19 stream gradients. Many streams in this region have substantial groundwater inputs. The substrate of these streams is
20 highly variable but often consists of gravel, cobble, or bedrock with sand and silt in slower moving areas and
21 pools. The MAB, in southeastern Missouri, is a broad flat plain with low stream gradients, most of which have been
22 channelized for agriculture, the predominant land cover in the region (83%; Blodgett and Lea 2005; Sowa *et al.*
23 2007). The substrate varies throughout the subregion but larger and slower moving streams generally have fine silt
24 while streams with faster flow are comprised of sand and small gravel substrates (Sowa *et al.* 2007). Among
25 subregions, agricultural and forest land cover vary substantially, however the percent urban land use ranges only
26 from 1.8% to 4.1% (Blodgett and Lea 2005).

2.2. Fish data

Our study used 1,499 fish samples which were collected using a statewide stratified random sampling design (Fischer and Combes 2003) of unique confluence to confluence stream segments based on the 1:100,000 stream network file developed by Sowa *et al.* (2007). Samples were collected between April and September, 2000 to 2011 from permanent wadeable streams (2nd-5th Strahler order) as a part of a wadeable stream monitoring program (Resource Assessment and Monitoring Program; RAM) through the Missouri Department of Conservation (Figure 1). All samples within the RAM database were collected using standardized methods; block nets enclosed a sample reach (length 40 times the mean wetted width, minimum 150 m) which was then thoroughly sampled via electrofishing and seining (Fischer and Combes 2003), with all individuals identified to species and enumerated.

Sample Locations

Figure 1: Map of sample locations used in this study for fishes collected from 2000 to 2011 in wadeable streams in Missouri.

2.3. Scoring system and classification of species vulnerability to threats

Species vulnerability scores were calculated using two different approaches: one based on species responses to measured threats and the other based on species traits (Figure 2). Both indices were classified and scored for species environmental tolerance, dispersal ability, restricted distribution, and rarity as well as the habitat resilience of their habitats. Species environmental tolerance scores were calculated by summing the scores for habitat condition, temperature, and flow vulnerability separately for each of the two indices, while all other criteria were scored the same way for both indices (Figure 2). Species sensitivity scores were calculated by multiplying rarity scores by the sum of environmental tolerance, dispersal ability, and range size scores, while the final species score is the sum of the sensitivity and habitat resilience scores (Figure 2). Scores were assigned to each species based on the classification of species vulnerability to each threat category. Rarity was incorporated as a multiplicative factor rather than an additive factor because rarity is expected to compound the impacts of the other components. Scores for each threat category were assigned as either discrete (0 or 1) for vulnerable (1) or not (0) classifications, continuous-additive (0 to 1) for vulnerability which was quantified along a gradient and incorporated into the vulnerability index as an additive component, or continuous-multiplicative (1 to 2) for vulnerability which was quantified along a gradient and incorporated into the vulnerability index as a multiplicative component. Multiplicative components were scaled from 1 to 2, rather than 0 to 1, because these components are only intended to have no effect or increase a species score, not reduce it.

Figure 2: Conceptual diagram of stepwise procedure for calculating species vulnerability scores using the Species Response Index and Trait Association Index. Inputs for dispersal, range, rarity and habitat resilience were the same for both indices; differences between index calculations occurred only for the environmental tolerance scoring criteria. Numbers in parenthesis indicate potential scoring range. For all “0 or 1” scoring, species meeting listed criteria receive a score of 1, while all others receive a score of 0.

Environmental tolerance was scored using the discrete value system by summing the vulnerable (1) or not (0) classifications for vulnerability to habitat alteration, change in stream temperatures, and change to flow regimes (Figure 2). Two separate environmental tolerance scores were calculated; one using a traits-based approach and another using a measured species response approach. Classifications for the traits-based approach were made from a review of literature that identified traits linked to vulnerability to each of the threats. Species which exhibit either lithophilic spawning or benthic invertivory were classified as vulnerable to habitat alteration as both have been linked to sensitivity to sedimentation (Berkman and Rabeni 1987; Mims *et al.* 2010). Cool- or cold-water adapted species have been identified as vulnerable to warming stream temperatures and were classified as such for our indices (Matthews and Zimmerman 1990; Eaton and Scheller 1996; Mohseni *et al.* 2003; Lyons *et al.* 2010; Mims *et al.* 2010). Species with periodic life history strategy (prefer predictable flow patterns), and the equilibrium life

1 history strategy (prefer low variability in flow patterns) are expected to be more vulnerable than species exhibiting
2 the opportunistic life history strategy (thrive in streams with unpredictable and variable flow patterns) (Winemiller
3 and Rose 1992; Olden and Kennard 2010; Mims and Olden 2012, 2013).

4 The species response index used species specific criteria to identify the environmental tolerance of species
5 to habitat alteration, changes in stream temperature, and change to flow regimes (Figure 2). Esselman *et al.* (2011)
6 conducted a literature review of fish based IBIs from across North America and compiled a list of intolerant
7 species. Species classified as intolerant by Esselman *et al.* (2011) within the World Wildlife Fund ecoregions (Abell
8 *et al.* 2008) which intersect Missouri were classified as vulnerable to habitat alteration for our species response
9 index. We determined species' associations with extremes in water temperature and stream flow metrics (i.e.,
10 representative of changes that could occur with changes in climate) using Indicator Species Analysis (ISA, Dufrene
11 and Legendre 1997). Fish data used for this determination were acquired from state and federal programs targeting
12 entire stream assemblages in the Temperate Plains (TPL) and Southern Appalachians (SAP) aggregated ecoregions
13 comprising Missouri (Daniel *et al.* 2015). To characterize water temperature and flow condition, daily stream flow
14 and temperature records were acquired from the USGS Water Data for the Nation (U.S. Geological Survey, 2010)
15 and summarized into 171 flow metrics (using the National Hydrologic Assessment Tool,
16 <https://www.sciencebase.gov/catalog/item/5387735ee4b0aa26cd7b5461>) and 169 temperature metrics (Tsang *et al.*
17 In Review). Locations of fish samples were associated with stream reaches of the NHDPlusV1 where temperature
18 and flow data were collected, and thus were presumed to characterize habitat conditions affecting fish species in
19 those stream reaches. To test for associations, ISA calculates indicator values for groupings of sites having high
20 species abundances with extreme values for a particular metric. High indicator values determined for a grouping of
21 sites with high species abundances at extremes in values for a particular metric (and corresponding low indicator
22 values for another grouping for more moderate metric values) support the idea that a species may be associated with
23 extremes in conditions. For this study, we used Threshold Indicator Taxa Analysis (TITAN; Baker and King, 2010)
24 to perform ISA iteratively, testing values of metrics that split species into two groupings across gradients
25 represented by each flow and temperature metric. TITAN calculates indicator value scores for all possible
26 groupings, and then identifies the point along the metric gradient that yields the group of sites with the highest
27 indicator value of all groupings tested (i.e., the change point). Significant indicator values for groups on either side

1 of a change point were used to suggest positive or negative association between species and flow metrics. We
2 identified significant associations between species and extremes in metrics using a p-value of 0.01, purity ≥ 0.95 ,
3 and reliability ≥ 0.9 , following Baker and King (2010). Relationships between temperature and flow metrics and
4 current climate metrics (Hostetler *et al.* 2011) were assessed using Pearson product-moment correlations. Any
5 correlation coefficient with an absolute value greater than 0.5 was considered to show strong associations. We used
6 data from GENMOM, a global climate model which was dynamically downscaled to North America, to determine
7 how climate would change between current and future conditions (five timesteps; 2020-2024, 2030-2034, 2040-
8 2044, 2065-2069, and 2085-2089). The GENMOM GCM was selected because it was the most conservative for late
9 century predictions of temperature and precipitation of the downscaled climate models available for our study
10 region (GENMOM, ECHAM5, and GFDL) and was demonstrated to perform similarly to other climatologies based
11 on PMIP2 and PMIP3 simulations, paleoclimate data-model analyses, and proxy reconstructions (Hostetler *et al.*
12 2011; Herb *et al.* 2014, Alder and Hostetler 2015). The directionality of the change in climate metrics between
13 current and future conditions was used to determine whether a species group (positive or negative association with
14 a flow or temperature metric) would shift towards increased or decreased abundance based on predicted changes in
15 climate. Species which were linked to a decrease in abundance by any temperature or flow metric were classified as
16 vulnerable to predicted changes in temperature or stream flow, respectively.

17 We classified species with low dispersal ability as vulnerable. Radinger *et al.* (2013) quantified the average
18 dispersal distance for the mobile component (dispersers) for nine fish families. Species from three families,
19 Fundulidae, Cottidae, and Percidae, had limited dispersal abilities, averaging less than 250 m (Radinger and Wolter
20 2013). Species in these families were classified as vulnerable and received scores of 1 (Figure 2), with the exception
21 of *Sander vitreus*, a Percid which is known to undertake annual spawning migrations of greater than 100 km
22 (Dupont *et al.* 2006). Species from all other families received scores of 0 for the dispersal ability component. Species
23 range size and rarity have been linked to vulnerability due to environmental change (Poff *et al.* 2012). Range size
24 was scored using the continuous-additive approach (Figure 2) and was quantified based on the number of
25 ecological drainage units (EDUs) in which a species has a known occurrence. Ecological drainage units generally
26 follow the boundaries of the major river basins in the state and were developed by identifying contiguous units
27 made up of multiple USGS 8-digit hydrologic units with relatively similar fish assemblages (Sowa *et al.* 2007).

EDUs are nested within subregion boundaries with a total of 5 EDUs in the Plains, 9 in the Ozarks, and 3 in the MAB (Sowa *et al.* 2007). These values were then scaled from 0 for species which occur in all 17 EDUs within Missouri to 1 for species which occur in a single EDU. Intermediate values were calculated along this scale (e.g., a species found in 25% of EDUs would receive a score of 0.75, a species found in 88% of drainages would receive a score of 0.12). Similarly, species rarity was calculated by using scaled values from 1 to 2. Species were considered common if they occurred at greater than 10% of sites (87 species; Figure 3). Using the continuous-multiplicative approach, species which were found at <10% of sites received scores scaled from 1 (10% of sites) to 2 (A single site, 0.01% of sites). Therefore a species found at seven percent of sites would receive a score of 1.3 while a species found at two percent of sites would receive a score of 1.8.

Figure 3: Histogram of species rarity for fish sampled from 1,499 Wadeable stream segments in Missouri from 2000 to 2011. The dashed line represents threshold used to classify rare species.

Habitat resilience scores were calculated as the ratio of the number of connected stream kilometers per occurrence scaled using the continuous additive approach from 0 for the maximum value (1,787 km/occurrence) to 1 for the minimum value (123 km/occurrence) (Figure 2). The amount of connected stream kilometers was calculated by partitioning out 2nd-6th order stream segments of the stream network GIS layer (Blodgett and Lea 2005). Stream segments classified as stream or stream/river (i.e., not reservoir or impoundment) were removed, which clipped the stream network at any location which flowed into an impoundment. The number of connected

stream kilometers for each intact stream segment was calculated with the Rivex tool (Hornby 2014), from which we quantified the average number of connected stream kilometers per occurrence for each species. Species with high average connected stream kilometers per occurrence inhabit areas with relatively high stream network connectivity, while species with relatively low values occur in fragmented habitats with low connectivity.

2.4. Index comparisons

Analyses were conducted to determine whether the species response and trait association approaches for classifying environmental tolerance achieved similar results, whether listed species received higher vulnerability scores than unlisted species, and whether there were patterns in the distribution of vulnerable species between the ecologically diverse aquatic subregions of Missouri. The percent agreement between the two approaches for classifications of vulnerability to habitat alteration, changes in stream temperatures, and change to flow regimes was used to determine the consistency of the approaches to classifying environmental tolerance. A Fischer's exact test was used to determine if the indices showed a positive odd's ratio ($\alpha < 0.01$) for vulnerable classifications of the environmental tolerance components of the two approaches. A two-way analysis of variance was used to determine if mean vulnerability score differed by listing status, (Missouri species of conservation concern including all federally threatened and endangered species [Missouri Natural Heritage Program 2012]), and index. Pearson's correlation analysis was used to determine if the two indices were related based on the number and proportion of vulnerable species at each of the study sites for each of the three components of environmental tolerance. An analysis of variance was used to determine whether the mean number and mean proportion of vulnerable species to each of the three environmental tolerance components differed by aquatic subregion. All statistical calculations were performed using R statistical software (R Development Core Team 2011).

3. RESULTS

3.1. Index development

The trait association approach to classifying environmental tolerance was used to calculate vulnerability scores for all 133 species with a mean vulnerability score of 3.83 ± 0.36 95% CI (Appendix A). Environmental tolerance was classified for 101 of the 133 species using the species response approach. This index produced a mean vulnerability score of 2.60 ± 0.30 95% CI (Appendix A). The remaining 32 species were not evaluated because we did not have a sufficient number of occurrences which could be associated with temperature and

flowmetrics. A total of three species declined in abundance with predicted changes in four temperature metrics (Appendix A), whereas a total of 38 species were shown to have declines in abundance with predicted changes in five flow metrics (Appendix B) and 43 species were identified as being vulnerable to habitat alteration (Appendix C). The trait association approach classified 71, 55, and 54 species vulnerable to habitat alteration, predicted changes in stream temperature, and predicted changes in flow regimes respectively (Appendix C). For each of the 133 species rarity, dispersal, range, and habitat resilience scores were calculated (Appendix C). The mean rarity score was 1.48 with 65.4% of species classified as rare (i.e., occurred in <10% of sites). Thirty three species (24.8%) were classified as vulnerable based on limited dispersal ability. The mean number of EDUs in which a species was present was 8 (of 17 total), which equates to a score of 0.53. The mean value for habitat resilience was 0.53 which equates to an average of 743 connected kilometers per occurrence.

3.2.Index comparisons

The species response and trait association approaches to classifying environmental tolerance showed similarities and differences in classification. The most consistent classification between methods was for vulnerability to habitat alteration (65.3%), followed by changing temperatures (60.4%) and changes to flow regimes (51.5%) (Table 1). If a species was classified as vulnerable to habitat alteration for one index, it was 3.60 times more likely to be classified as vulnerable to habitat alteration with the other index (Odd's ratio of 3.60, $p=0.002$). In contrast, no association was found between the trait and response based approaches to classifying vulnerability based on changing temperatures (Odd's ratio of 0.79, $p=0.999$) or changes to flow regimes (Odd's ratio of 1.05; $p=0.99$) (Table 1).

Table 1: Comparison of trait association and species response approaches for classifying species environmental tolerances based on percent agreement of species classifications, odd's ratio and p-value calculated using Fischer's exact test.

Threat	Percent Agreement	Odd's Ratio	P-Value
Habitat Alteration	65.3	3.60	0.003
Temperature	60.4	0.79	0.999
Flow	51.5	1.05	0.999

Mean vulnerability scores differed by both listing status ($p<0.001$), and index ($p<0.001$), with no interaction between listing status and index ($p=0.693$). For each of the indices listing status was significantly related

to score ($p<0.001$) with mean scores for listed species of 1.0 (response) to 1.4 (trait) units greater than unlisted species (Figure 4). In addition, rare species had higher vulnerability scores for both approaches (Response: Rare=3.05, Not =2.10, $P<0.001$; Trait: Rare=4.58, Not=2.47, $P<0.001$).

Environmental Tolerance Classification System
 Figure 4: Mean score (+/- 95% Confidence Interval) for listed and unlisted species using the species response and traits association approaches for classifying environmental tolerance.

The number of vulnerable species at each site was highly correlated between indices for habitat alteration ($r=0.94$), but much less for temperature ($r=0.37$), and flow alteration ($r=0.23$) (Figure 5). Similarly, the proportion of vulnerable species at each site had the highest correlation between indices for habitat alteration ($r=0.87$), but less so for temperature ($r=0.15$), and flow alteration ($r=-0.25$) (Figure 5). The mean number of vulnerable species per site was greatest in the forest-dominated Ozark subregion for both habitat alteration and temperature using both approaches of classifying environmental tolerances (Table 2). The greatest numbers of species vulnerable per site for flow alteration occurred in the MAB with the trait association approach, and the Plains with the species response approach (Table 2). The mean proportion of vulnerable species per site was highest in the Ozarks for habitat and temperature for the trait association approach and habitat for the species response approach (Table 2). The greatest proportion of flow vulnerable species per site was highest in the MAB for the trait association, and in the plains for the species response approach (Table 2).

Figure 5: Correlation between the trait and response based approaches for classifying environmental tolerance based on the number and proportion of vulnerable species at each of the study sites, r = Pearson's correlation coefficient.

Table 2: Mean number and proportion of vulnerable species at sites within each subregion by index and threat; highest value in bold. P-values tested whether the mean number or proportion of species differed by subregion using a one-way ANOVA. "MAB"=Mississippi Alluvial Basin.

Index	Plains			Ozarks		MAB		P-Value	
	Threat	Number	Proportion	Number	Proportion	Number	Proportion	Number	Proportion
Trait									
Association									
	Habitat	4.6	0.30	10.6	0.55	3.3	0.18	<0.001	<0.001
	Flow	5.9	0.41	8.0	0.42	10.1	0.28	<0.001	<0.001
	Temperature	3.5	0.24	8.0	0.42	5.1	0.28	<0.001	<0.001
Species									
Response									
	Habitat	3.0	0.19	9.2	0.51	2.4	0.16	<0.001	<0.001
	Flow	3.5	0.25	2.3	0.13	2.1	0.14	<0.001	<0.001
	Temperature	1.1	0.01	1.5	0.01	0.0	0.0	0.003	0.027

1 **4.DISCUSSION**

2 The results of this assessment provide a framework for assessing stream fish vulnerability to climate
3 change and habitat alteration.Stream communities worldwide are likely to be impacted by climate change and
4 habitat alteration(Eaton and Scheller 1996; Poff *et al.* 2002; Heino *et al.* 2009), and evaluating species'
5 vulnerability will be critical for conservation planning efforts (Poff *et al.* 2012).Our assessment provides a new
6 approach for stream fish vulnerability assessment which can be applied and modified for use in other regions. The
7 classification of species as vulnerable based on their environmental tolerances can be refined as new data becomes
8 available. Models of stream temperature and flow will allow exposure to change to be evaluated. Additionally,
9 these indices can be modified to fit the needs of specific management interests.

10 The trait association approach produced higher vulnerability scores, and exhibited a greater range of scores
11 than the species response approach.Both of these discrepancies between the indices largely stem from the lack of
12 data available to classify the environmental tolerances of rare species using the response-based approach.Of the 32
13 species for which environmental tolerance could not be classified based on measured species responses, 31 were
14 classified as rare.Rare species had significantly higher vulnerability scores than common species using both
15 approaches which likely resulted in higher average and maximum scores for the trait association index.Although
16 some regions appear to have sufficient information regarding the vulnerability of all species to specific metrics
17 (Moyle et al. 2013), many regions lack the stream temperature and flow data necessary to elicit the responses of
18 species to environmental threats necessary for a broadly applied vulnerability assessment (Deweber *et al.*
19 2014).Quantitative assessments of stream fish relationships with potential stressors are needed (Schlosser 1991) and
20 broad scale vulnerability assessments which rely on measured species responses to environmental threats will
21 depend on additional research to determine the impacts of threats on all species.Many of these rare species are
22 likely the most vulnerable, so a failure to include them in a vulnerability assessment would likely leave an
23 important data gap and present a biased assessment.

24 A traits based approach allowed for a more complete assessment of species vulnerability as traits have been
25 described for most North American stream fish (Frimpong and Angermeier 2009; Mims *et al.* 2010).The use of
26 traits has flourished for the assessment of stream fish communities, and is a useful tool in stream fish ecology and
27 conservation (Parent and Schriml 1995; Poff 1997; Olden and Kennard 2010; Culp *et al.* 2011; McManamay *et*

al.2014).Our results suggest a traits based approach allows for a more complete representation of species; however uncertainty still exists regarding our ability to link traits to species vulnerability.Species have complex relationships with the environment and stressors, and the use of single or even a suite of traits may not provide enough information to fully account for this complexity. More research linking specific traits to environmental threats would help decrease the uncertainty of these assessments and validate trait based measures of vulnerability.

Substantial discrepancies existed in the number of species classified as vulnerable to environmental tolerance components using the trait and response based approaches.For species which were assessed using both indices a similar number of species were classified as vulnerable to habitat alteration (50 versus 43 respectively) and changes to flow regime (47 versus 38), however the number of species classified as vulnerable to changing stream temperatures (39 versus 3)was considerably higher for the trait association approach. The lack of consensus between the use of traits versus measured responses for temperature and flow classifications may stem from the lack of available temperature and flow data available to link species responses to projected changes in climate. As additional stream temperature and flow data become available the relationships between species and these variables will be able to be evaluated more fully. As more data become available we will be able to evaluate a broader range of conditions and also have more replicates across this range. This information will aid in identifying causal, mechanistic explanations for the relationship between stream fish and stream temperature and flow.

Vulnerability to habitat alteration was classified relatively consistently by both approaches (65.7% agreement), suggesting that both indices were likely successful in capturing and representing species vulnerability to land use changes.Species classified as vulnerable to habitat alteration with the trait association index but not the measured response index were often those identified as tolerant of alteration but belong to families often viewed as intolerant (e.g. johnny darter (*Etheostoma nigrum*), orangethroat darter (*Etheostoma spectabile*), and golden redhorse(*Moxostoma erythrurum*); Karr 1981), and other common, tolerant species (e.g. orangespotted sunfish(*Lepomis humilis*) (Meador and Carlisle 2007), creek chub (*Semotilus atromaculatus*) (Miller *et al.* 1988), and black bullhead (*Ameiurus melas*) (Novomeská and Kováč 2009)).Species classified as vulnerable to habitat alteration by the species response index but not by the trait association index included a number of Centrarchids, (e.g. warmouth (*Lepomis gulosus*), longear sunfish (*Lepomis megalotis*), smallmouth bass (*Micropterus dolomeiu*), shadow bass (*Ambloplites ariommus*), Ozark bass (*Ambloplites constellatus*), and rock bass (*Ambloplites*

1 *rupestris*)), and some species of the genus *Cyprinella* (e.g. whitetail shiner (*Cyprinella galactura*), spotfin shiner
2 (*Cyprinella spiloptera*), and steelcolor shiner (*Cyprinella whipplei*)). Although these indices are relatively consistent
3 in classifying vulnerability to habitat alteration, there are still some vulnerable species which were excluded by
4 each approach. The relative consistency between the approaches of classifying habitat vulnerable species suggests
5 that the current understanding of vulnerability to habitat alteration may be sufficient for the purposes of a broad-
6 scale assessment using either approach.

7 In contrast, vulnerability to changes in stream temperature and changes to flow regimes showed major
8 discrepancies between the two approaches. Although only three Missouri stream fish species (white sucker
9 (*Catostomus commersonii*), smallmouth bass, and central stoneroller (*Camptostoma anomalum*)) have shown
10 decreased abundances with future predicted climatic conditions according to our response based index, many
11 additional species may be vulnerable. In studies completed in areas where future stream temperature models exist,
12 species which have yet to exhibit decreased abundance or range contractions are predicted to in the future, and the
13 same is likely for some Missouri species (Mohseni *et al.* 2003; Heino *et al.* 2009; Lyons *et al.* 2010). This could be
14 due to our inability to assess a number of species using the response based approach due to limited data availability
15 and because the current range of observable conditions does not fully reflect the range of future conditions (we are
16 unable to assess future extremes which have not yet occurred). The trait association approach allows decreased
17 abundances and range contractions to be extrapolated even though they may not yet have been observed, and a
18 large number of Cyprinids (14 of 33) and Percids (8 of 15) may be vulnerable to future warming based on this
19 approach, however uncertainty exists as to whether this approach is able to account for the complexities of the
20 impacts of temperature on fish. Species may appear vulnerable based on a certain trait, but may also have other
21 characteristics which allow them to be more or less resilient to increasing temperatures than we can infer based on
22 the simplistic, trait-based approach.

23 Substantial discrepancies also existed between the two indices for flow which suggests that there is a need
24 to resolve the conflicting results between vulnerability predictions based on life history strategy (Mims *et al.* 2010)
25 versus vulnerability predictions based on our measured species responses. The majority of Ictalurids (10/12), and
26 Centrarchids (15/16) were classified as vulnerable for the trait based approach, however very few of the species in
27 these families were classified as vulnerable based on measured responses (3/10), (3/14) respectively (Two species

1 from each of these families were not evaluated based on the species response approach). This lack of a measured
2 response for these equilibrium and periodic life history strategists suggests that other factors may complicate the
3 predicted relationship between these traits and decreases in abundance and occupancy associated with changes in
4 stream flow. These discrepancies suggest that a thoughtful comparison between the response and trait association
5 approaches should be used to select the most appropriate approach, and future research may consider rectifying
6 these differences. When selecting an approach for classifying environmental tolerances researchers and managers
7 may need to consider if sufficient information is available for target species, and which approach to classifying
8 environmental tolerance better fits the most current understanding of species vulnerability, particularly regarding
9 predicted changes in flow regimes and stream temperatures. The use of a traits based approach allows for a broader
10 species assessment, however it is limited in its ability to account for some of the complexity of species/environment
11 interactions. The response based approach provides more direct evidence of species vulnerability, however it is
12 limited in the number of species it can be used to assess based on data availability and limited to the current range
13 of observable environmental conditions, which likely doesn't fully encompass the range of future conditions.

14 Species vulnerability scores for both indices were greater for species listed as threatened, endangered, or of
15 conservation concern in Missouri suggesting that both of the indices achieve results that align with those of more
16 traditional methods of identifying vulnerable species. In Missouri, species of conservation concern were determined
17 through expert knowledge while also considering rarity, population trends, and threats following the NatureServe
18 Conservation Status Assessment Criteria (Missouri Natural Heritage Program 2012). All 23 species of conservation
19 concern were able to be evaluated using the trait association index, while only five species of conservation concern
20 were with the species response index. Our quantitative assessment of stream fish vulnerability presented here is
21 consistent with expert opinion and provides a general validation of our method. It also suggests that species of
22 conservation concern are likely to become increasingly vulnerable when faced with future threats.

23 Our analysis of the distribution of vulnerable species showed strong spatial trends. Both approaches to
24 classifying environmental tolerance identified the Ozarks as generally having both the highest numbers and
25 proportions of vulnerable species per site than in the other subregions. The exception to this was for flow alteration
26 vulnerability, for which the number and proportion of vulnerable species per site was highest in the MAB according
27 to the trait association approach and the Plains according to the species response approach. Streams in the Ozarks

1 have steeper stream gradients, more diverse substrates, cooler stream temperatures, and less human development
2 than Plains streams and are able to support more sensitive species (Sowa *et al.* 2007).The spatial consistency of the
3 counts and proportions of species classified as vulnerable to habitat alteration and changes in stream temperature
4 suggest that management for those threats can be focused on the Ozarks regardless of which approach to classifying
5 environmental tolerance is used.The inconsistency regarding flow vulnerability corroborates the previous
6 comparison of environmental tolerance classification by confirming that further research to determine species
7 vulnerability to predicted alterations to flow regimes is needed.Streams are predicted to experience greater variation
8 in flows (IPCC 2002; Poff *et al.* 2002), however the impact of increased variation in flow on stream fish is
9 unclear.Decision makers may need to consider which approach to classifying environmental tolerance best fits their
10 understanding of flow vulnerable species when making decisions regarding the identification and selection of areas
11 for management of these species.

12 Understanding the impact of uncertainty is crucial for any vulnerability assessment (Patt *et al.* 2005; Füssel
13 and Klein 2006) and many sources of uncertainty affect our understanding of stream fish vulnerability in
14 Missouri.Projected changes in climate have substantial uncertainty (Stainforth *et al.* 2007) so multiple scenarios
15 have been developed which predict a range of future temperature and precipitation conditions (IPCC
16 2013).Additional uncertainty surrounds how these predicted changes in climate will impact stream temperature and
17 flow, and currently Missouri does not have any models that make these predictions over a broad spatial scale
18 (Whitledge *et al.* 2006; Pease and Paukert 2014; Westhoff and Paukert 2014).Because of these limitations,
19 exposure, which is often used in vulnerability assessments (Glick *et al.* 2011; Staudinger *et al.* 2013), cannot be
20 used as a factor of discrimination in our species vulnerability analysis.Additionally, accounting for additional
21 threats which were not considered for in our assessment such as invasive species and pollution may shift the
22 relative vulnerability of some species.

23 There are clear tradeoffs between the two approaches of assessing environmental tolerance.Benefits of a
24 trait association approach include broad applicability to many species, consistency between studies/regions, and
25 ability to link to a causal mechanism (Parent and Schriml 1995; Poff 1997; Frimpong and Angermeier 2009; Culp
26 *et al.* 2011).Benefits of a measured species response approach include direct evidence of species responses to
27 threats, and the ability to account for complex interactions between species characteristics, their responses, and the

1 environment(Williams *et al.* 2008; Lyons *et al.* 2010).Disadvantages of a trait association approach include
2 difficulty representing variability in environmental tolerances, limited inference to known and quantified traits, and
3 difficulty accounting for complex species/environmental relationships.Disadvantages of a measured species
4 response approach include restriction to species which have data available, limitation of inference to threat impacts
5 which are observable in the environment or dependence on laboratory studies to simulate threat impacts, and
6 difficulty isolating the impact of a threat from other factors which may affect a species response. Additional
7 research which investigates species responses to threats could add to the feasibility of a response based assessment
8 of vulnerability.A more complete set of information regarding species temperature and flow tolerances,
9 bioenergetics, and population viability coupled with landscape models of stream temperatures and flow regimes
10 would assist in the development of response based models which could predict species specific vulnerability.

11 Many stream fish species are already threatened with local extirpation, and increases in habitat alteration,
12 changes in stream temperature, and changes to the flow regimes of streams will likely result in increases in the
13 number of species impacted as well as the magnitude of this imperilment (Poff *et al.* 2012).To conserve stream fish
14 more effectively, decision makers may need a better understanding of which species are most vulnerable to each of
15 the threats and overall.Our study created a quantifiable index of stream fish vulnerability based on the outline by
16 Poff *et al.*(2012) for 133 species in a Midwest US state.As more information is gathered for stream fishes the
17 application of more traditional vulnerability assessments following the criteria and framework established by Glick
18 *et al.*(2011) may be possible, however based on currently available information we believe this approach provides a
19 suitable alternative.Our assessment of vulnerability allows for identification of species vulnerable to specific threats
20 and also gives us a cumulative measure of a species vulnerability to a suite of threats.This study developed a
21 framework which can serve as a foundation for future vulnerability assessments and can be adapted to include new
22 information on species environmental tolerances, additional threats, and the weighting or scale of threats.Our hope
23 is that others can apply and modify this framework for assessment of vulnerable stream fish species in other
24 regions.

1 **5. ACKNOWLEDGEMENTS**

2 Funding for this project was provided by the U.S. Geological Survey National Climate Change and
3 Wildlife Science Center and the Northeast Climate Science Center. We would like to thank Matt Combes and Doug
4 Novinger of the Missouri Department of Conservation for sharing much of the data used in this study with us. We
5 would also like to thank Dr. Emmanuel Frimpong, Dr. Julian Olden, and Dr. Meryl Mims for sharing the trait
6 information which was used for this project. We also thank Dr. Jodi Whittier, Dr. Michelle Staudinger, and Dr. Tim
7 Matisziw as well as two anonymous reviewers for their feedback on the manuscript. Any use of trade, firm, or
8 product names is for descriptive purposes only and does not imply endorsement by the U.S. Government. The
9 Missouri Cooperative Fish and Wildlife Research Unit is jointly sponsored by the Missouri Department of
10 Conservation, the University of Missouri, the U.S. Geological Survey, the U.S. Fish and Wildlife Service, and the
11 Wildlife Management Institute.

12

13 **6. ROLL OF THE FUNDING SOURCE**

14 The U.S. Geological Survey National Climate Change and Wildlife Science Center and the Northeast
15 Climate Science Center (funding sources) played no role in determining the study design, the collection, analysis
16 and interpretation of data, the writing of the report, and in the decision to submit the article for publication.

17

7. REFERENCES

- Abell, R. (2002) Conservation Biology for the Biodiversity Crisis: a Freshwater Follow-up. *Conservation Biology***16**, 1435–1437.
- Abell, R., Thieme, M., Revenga, C., *et al.* (2008) Freshwater ecoregions of the world: a new map of biogeographic units for freshwater biodiversity conservation. *BioScience***58**, 403–414.
- Albanese, B., Angermeier, P. and Peterson, J. (2009) Does mobility explain variation in colonisation and population recovery among stream fishes? *Freshwater Biology***54**, 1444–1460.
- Alder, J. and Hostetler, S. (2015) Global climate simulations at 3000-year intervals for the last 21000 years with the GENMOM coupled atmosphere-ocean model. *Climate of the Past***11**, 449–471.
- Allan, J. (2004) Landscapes and Riverscapes: The Influence of Land Use on Stream Ecosystems. *Annual Review of Ecology, Evolution, and Systematics***35**, 257–284.
- Allan, J. and Flecker, A. (1993) Biodiversity Conservation in Running Waters. *BioScience***43**, 32–43.
- Angermeier, P. (1995) Ecological Attributes of Extinction-Prone Species: Loss of Freshwater Fishes of Virginia. *Conservation Biology***9**, 143–158.
- Bagne, K., Friggens, M. and Finch, D. (2011) *A System for Assessing Vulnerability of Species (SAVS) to Climate Change*. USDA Forest Service. Albuquerque, New Mexico.
- Baker, M., and King R. (2010) A new method for detecting and interpreting biodiversity and ecological community thresholds. *Methods in Ecology and Evolution***1**, 25–37.
- Barbour, M., Gerritsen, J., Snyder, B. and Stribling, J. (1999) Rapid bioassessment protocols for use in streams and wadeable rivers. *USEPA, Washington D.C.*
- Berkman, H. and Rabeni, C. (1987) Effect of siltation on stream fish communities. *Environmental Biology of Fishes***18**, 285–294.
- Blodgett, C. and Lea, R. (2005) *Missouri Landcover Project*. Missouri Resource Assessment Partnership. Columbia, Missouri.
- Culp, J., Armanini, D., Dunbar, M., *et al.* (2011) Incorporating traits in aquatic biomonitoring to enhance causal diagnosis and prediction. *Integrated Environmental Assessment and Management***7**, 187–197.
- Daniel, W., Infante D., Hughes R., Tsang Y., *et al.* (2015) Characterizing coal and mineral mines as a regional source of stress to stream fish assemblages. *Ecological Indicators***50**, 50–61.
- Detenbeck, N., DeVore, P., Niemi, G. and Lima, A. (1992) Recovery of temperate-stream fish communities from disturbance: a review of case studies and synthesis of theory. *Environmental Management***16**, 33–53.
- Deweber, J., Tsang, Y., Krueger, D., *et al.* (2014) Importance of understanding landscape biases in USGS gage locations: Implications and solutions for managers. *Fisheries* **39**, 155–163.
- Dudgeon, D., Arthington, A., Gessner, M., *et al.* (2006) Freshwater biodiversity: importance, threats, status and conservation challenges. *Biological Reviews***81**, 163–182.
- Dufrêne, M., and Legendre P (1997) Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecological monographs***67**, 345–366.
- Dupont, P., Bourret, V., and Bernatchez, L. Interplay between ecological, behavioral and historical factors in shaping the genetic structure of sympatric walleye populations (*Sander vitreus*). *Molecular Ecology***16**, 937–951.
- Eaton, J. and Scheller, R. (1996) Effects of Climate Warming on Fish Thermal Habitat in Streams of the United States. *Limnology and Oceanography***41**, 1109–1115.
- Esselman, P., Infante, D., Wang, L., Wu, D., Cooper, A. and Taylor, W. (2011) An index of cumulative disturbance to river fish habitats of the conterminous United States from landscape anthropogenic activities. *Ecological Restoration***29**, 133–151.
- Fagan, W., Unmack, P., Burgess, C. and Minckley, W. (2002) Rarity, fragmentation, and extinction risk in desert fishes. *Ecology***83**, 3250–3256.
- Fischer, S. and Combes, M. (2003) *Resource Assessment and Monitoring Program: Standard Operation Procedures - Fish Sampling*. Missouri Department of Conservation.
- Frimpong, E. and Angermeier, P. (2009) Fish traits: A database of ecological and life-history traits of freshwater fishes of the United States. *Fisheries***34**, 487–495.
- Füssel, H. and Klein, R. (2006) Climate Change Vulnerability Assessments: An Evolution of Conceptual Thinking. *Climatic Change***75**, 301–329.

- 1 Gillette, D., Tiemann, J., Edds, D., and Wildhaber, M. (2005) Spatiotemporal Patterns of Fish Assemblage
2 Structure in a River Impounded by Low-Head Dams. *Copeia***3** 539-549.
- 3 Girvetz, E., Zganjar, C., Raber, G., Maurer, E., Kareiva, P. and Lawler, J. (2009) Applied Climate-Change
4 Analysis: The Climate Wizard Tool. *PLoS ONE***4**, e8320.
- 5 Glick, P., Stein, B. and Edelson, N. (2011) *Scanning the conservation horizon: a guide to climate change*
6 *vulnerability assessment*. National Wildlife Federation Washington, DC, USA.
- 7 Hanski, I. (1998) Metapopulation dynamics. *Nature***396**, 41–49.
- 8 Heggenes, J., Qvenild, T., Stamford, M. and Taylor, E. (2006) Genetic structure in relation to movements in wild
9 European grayling (*Thymallus thymallus*) in three Norwegian rivers. *Canadian Journal of Fisheries and*
10 *Aquatic Sciences***63**, 1309–1319.
- 11 Heino, J., Virkkala, R. and Toivonen, H. (2009) Climate change and freshwater biodiversity: detected patterns,
12 future trends and adaptations in northern regions. *Biological Reviews***84**, 39–54.
- 13 Herb, W., Johnson, L., Jacobson, P., and Stefan, H. (2014) Projecting cold-water fish habitat in lakes of the glacial
14 lakes region under changing land use and climate regimes. *Canadian Journal of Fisheries and Aquatic*
15 *Sciences***71**, 1334-1348.
- 16 Hering, D., Johnson, R., Kramm, S., Schmutz, S., Szoszkiewicz, K. and Verdonshot, P. (2006) Assessment of
17 European streams with diatoms, macrophytes, macroinvertebrates and fish: a comparative metric-based
18 analysis of organism response to stress. *Freshwater Biology***51**, 1757–1785.
- 19 Hostetler, S., Alder, J., and Allan, A. (2011) Dynamically Downscaled Climate Simulations over North America:
20 Methods, Evaluation, and Supporting Documentation for Users. U.S. Geological Survey Open-File Report
21 2011-1238.
- 22 Hornby, D. (2014) *RivEX (Version 10.15) [Software]*. <http://rivex.co.uk/index.html>.
- 23 IPCC (2013) Climate Change 2013: The Physical Science Basis. Working Group 1 Contribution to the Fifth
24 Assessment Report of the Intergovernmental Panel on Climate Change. 1535 pp. Intergovernmental Panel
25 on Climate Change, Cambridge, United Kingdom and New York, NY.
- 26 IPCC (2007) *Climate Change 2007: Impacts, Adaptation and Vulnerability: Working Group II Contribution to the*
27 *Fourth Assessment Report of the IPCC Intergovernmental Panel on Climate Change*, (Vol. 4). Cambridge
28 University Press.
- 29 IPCC (2002) Climate Change and Biodiversity. IPCC Technical Paper V. Intergovernmental Panel on Climate
30 Change: Working Group II.
- 31 Jackson, D. and Marmulla, G. (2001) The influence of dams on river fisheries. *FAO Fisheries technical paper*, 1–
32 44.
- 33 Karr, J. (1981) Assessment of biotic integrity using fish communities. *Fisheries***6**, 21–27.
- 34 Lidicker Jr., W. and Stenseth, N. (1992) To disperse or not to disperse: who does it and why?. In: *Animal Dispersal*.
35 (eds N. Stenseth and W. Lidicker Jr.). Springer Netherlands, pp 21–36.
- 36 Lyons, J., Stewart, J. and Mitro, M. (2010) Predicted effects of climate warming on the distribution of 50 stream
37 fishes in Wisconsin, U.S.A. *Journal of Fish Biology***77**, 1867–1898.
- 38 Mace, G., Collar, N., Gaston, K., *et al.* (2008) Quantification of extinction risk: IUCN's system for classifying
39 threatened species. *Conservation Biology***22**, 1424–1442.
- 40 Malmqvist, B. and Rundle, S. (2002) Threats to the running water ecosystems of the world. *Environmental*
41 *conservation***29**, 134–153.
- 42 Master, L., Flack, S., and Stein, B. (1998) *Rivers of life: critical watersheds for protecting freshwater biodiversity*.
43 Nature Conservancy Arlington, Virginia.
- 44 Matthews, W. and Zimmerman, E. (1990) Potential effects of global warming on native fishes of the southern Great
45 Plains and the Southwest. *Fisheries***15**, 26–32.
- 46 McManamay, R., Bevelhimer, M. and Frimpong, E. (2014) Associations among hydrologic classifications and fish
47 traits to support environmental flow standards. *Ecohydrology***8**, 460-479.
- 48 Meador, M. and Carlisle, D. (2007) Quantifying tolerance indicator values for common stream fish species of the
49 United States. *Ecological Indicators***7**, 329–338.
- 50 Miller, D., Hughes, R., Karr, J., *et al.* (1988) Regional Applications of an Index of Biotic Integrity for Use in Water
51 Resource Management. *Fisheries***13**, 12–20.
- 52 Mims, M. and Olden, J. (2013) Fish assemblages respond to altered flow regimes via ecological filtering of life
53 history strategies. *Freshwater Biology***58**, 50–62.

- 1 Mims, M. and Olden, J. (2012) Life history theory predicts fish assemblage response to hydrologic regimes.
2 *Ecology***93**, 35–45.
- 3 Mims, M., Olden, J., Shattuck, Z. and Poff, N. (2010) Life history trait diversity of native freshwater fishes in North
4 America. *Ecology of Freshwater Fish***19**, 390–400.
- 5 Missouri Natural Heritage Program (2012) *Missouri Species and Communities of Conservation Concern Checklist*.
6 Missouri Department of Conservation.
- 7 Mohseni, O., Stefan, H. and Eaton, J. (2003) Global Warming and Potential Changes in Fish Habitat in U.S.
8 Streams. *Climatic Change***59**, 389–409.
- 9 Moyle, P., Kiernan, J., Crain, P. and Quinones, R. (2013) Climate change vulnerability of native and alien
10 freshwater fishes of California: a systematic assessment approach. *PloS one***8**, e63883.
- 11 Novomeská, A. and Kováč, V. (2009) Life-history traits of non-native black bullhead *Ameiurus melas* with
12 comments on its invasive potential. *Journal of Applied Ichthyology***25**, 79–84.
- 13 Olden, J. and Kennard, M. (2010) Intercontinental comparison of fish life history strategies along a gradient of
14 hydrologic variability. In: *American Fisheries Society Symposium*, Vol. 73. pp 83–107.
- 15 Olden, J., Poff, N. and Bestgen, K. (2008) Trait synergisms and the rarity, extirpation, and extinction risk of desert
16 fishes. *Ecology***89**, 847–856.
- 17 Olden, J., Hogan, Z. and Zanden, M. (2007) Small fish, big fish, red fish, blue fish: size-biased extinction risk of
18 the world's freshwater and marine fishes. *Global Ecology and Biogeography***16**, 694–701.
- 19 Parent, S. and Schriml, L. (1995) A model for the determination of fish species at risk based upon life-history traits
20 and ecological data. *Canadian Journal of Fisheries and Aquatic Sciences***52**, 1768–1781.
- 21 Patt, A., Klein, R. and de la Vega-Leinert, A. (2005) Taking the uncertainty in climate-change vulnerability
22 assessment seriously. *Comptes Rendus Geoscience***337**, 411–424.
- 23 Pease, A. and Paukert, C. (2014) Potential impacts of climate change on growth and prey consumption of stream-
24 dwelling smallmouth bass in the central United States. *Ecology of Freshwater Fish* **23**, 336–346.
- 25 Pflieger, W. and Missouri Department of Conservation (1989) *Aquatic community classification system for*
26 *Missouri*. Missouri Dept. of Conservation.
- 27 Pflieger, W. (1970) A distributional study of Missouri fishes. University of Kansas Publications, Museum of
28 Natural History. **20**, 225–570.
- 29 Poff, N. (1997) Landscape filters and species traits: towards mechanistic understanding and prediction in stream
30 ecology. *Journal of the North American Benthological Society*, 391–409.
- 31 Poff, N., Olden, J. and Strayer, D. (2012) Climate change and freshwater extinction risk. In: *Saving a Million*
32 *Species: Extinction Risk from Climate Change*. (ed L. Hannah). Island Press, pp 309–336.
- 33 Poff, N., and Zimmerman, J. (2012) Ecological responses to altered flow regimes: a literature review to inform the
34 science and management of environmental flows. *Freshwater Biology***55**, 194–205.
- 35 Poff, N., Brinson, M. and Day, J. (2002) Aquatic ecosystems and global climate change. *Pew Center on Global*
36 *Climate Change, Arlington, VA*, 44.
- 37 Poff, N., Allan, J., Bain, M., *et al.* (1997) The Natural Flow Regime. *BioScience***47**, 769–784.
- 38 Radinger, J. and Wolter, C. (2013) Patterns and predictors of fish dispersal in rivers. *Fish and Fisheries***15**, 456-
39 473.
- 40 R Development Core Team (2011) *R: A Language and Environment for Statistical Computing*. R Foundation for
41 Statistical Computing, Vienna, Austria.
- 42 Ricciardi, A. and Rasmussen, J. (1999) Extinction rates of North American freshwater fauna. *Conservation*
43 *Biology***13**, 1220–1222.
- 44 Schlosser, I. (1991) Stream Fish Ecology: A Landscape Perspective. *BioScience***41**, 704–712.
- 45 Schnieder, S., Semenov, S., Patwardhan, A., *et al.* (2007) Assessing key vulnerabilities and the risk from climate
46 change. *Climate change 2007: Impacts, adaptation and vulnerability: Contribution of working group II to*
47 *the fourth assessment report of the intergovernmental panel on climate change*, 779–810.
- 48 Simon, T. (1999) *Assessing the sustainability and biological integrity of water resources using fish communities*.
49 CRC Press.
- 50 Sowa, S., Annis, G., Morey, M. and Diamond, D. (2007) A Gap Analysis and Comprehensive Conservation
51 Strategy for Riverine Ecosystems of Missouri. *Ecological Monographs***77**, 301–334.

- 1 Stainforth, D., Allen, M., Tredger, E. and Smith, L. (2007) Confidence, uncertainty and decision-support relevance
2 in climate predictions. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and*
3 *Engineering Sciences***365**, 2145–2161.
- 4 Staudinger, M., Carter, S., Cross, M., *et al.* (2013) Biodiversity in a changing climate: a synthesis of current and
5 projected trends in the US. *Frontiers in Ecology and the Environment***11**, 465–473.
- 6 Tonn, W. (1990) Climate change and fish communities: a conceptual framework. *Transactions of the American*
7 *Fisheries Society***119**, 337–352.
- 8 Tsang, Y., Infante, D., Stewart, J., Wang, L., Tingley, D., Thornbrugh, D., Cooper, A., Daniel, W. (In Review)
9 StreamThermal: A software package for calculating thermal metrics from stream temperature data.
10 Fisheries.
- 11 Turner, B., Kasperson, R., Matson, P., *et al.* (2003) A framework for vulnerability analysis in sustainability science.
12 *Proceedings of the national academy of sciences***100**, 8074–8079.
- 13 U.S. Geological Survey (2010) National Water Information System data available on the World Wide Web (USGS
14 Water Data for the Nation), accessed [October 18, 2010], at URL [<http://waterdata.usgs.gov/nwis/>].
- 15 Wang, L., Infante, D., Esselman, P., Cooper, A., Wu, D., Taylor, W., Beard, D., Whelan, G., Ostroff, A. (2011) A
16 hierarchical spatial framework and database for the national river fish habitat condition assessment.
17 *Fisheries***36**, 436–449.
- 18 Westhoff, J. and Paukert, C. (2014) Climate Change Simulations Predict Altered Biotic Response in a Thermally
19 Heterogeneous Stream System. *PloS one***9**, e111438.
- 20 Whitley, G., Rabeni, C., Annis, G. and Sowa, S. (2006) Riparian Shading and Groundwater Enhance Growth
21 Potential for Smallmouth Bass in Ozark Streams. *Ecological Applications***16**, 1461–1473.
- 22 Williams, S., Shoo, L., Isaac, J., Hoffmann, A. and Langham, G. (2008) Towards an Integrated Framework for
23 Assessing the Vulnerability of Species to Climate Change. *PLoS Biol***6**, e325.
- 24 Winemiller, K. and Rose, K. (1992) Patterns of life-history diversification in North American fishes: implications
25 for population regulation. *Canadian Journal of Fisheries and Aquatic Sciences***49**, 2196–2218.
- 26 Young, B., Byers, E., Gravuer, K., Hall, K., Hammerson, G. and Redder, A. (2011) Guidelines for Using the
27 NatureServe Climate Change Vulnerability Index. NatureServe.
- 28

Appendix A: Species response categorization based on temperature metrics for species which showed decreases in abundance with predicted future conditions (species which did not show a decrease in abundance for any of the metrics were not included in the table). Signs indicate direction of future change based on correlation between temperature metrics and climate data. ‘X’ indicates the metric to which a given species is predicted to show declines in abundance with predicted future changes to temperature metrics.

Scientific Name	Temperate Plains Ecoregion	Southern Appalachian Ecoregion		
	Maximum of 7 day moving average of daily mean (+)	Spring average daily mean (+)	Summer average daily mean (+)	Average daily mean in winter (+)
CAMPOSTOMA PULLUM				X
CATOSTOMUS COMMERSONI		X		
MICROPTERUS DOLOMIEU				X

Appendix B: Species response categorization based on flow metrics for species which showed decreases in abundance with predicted future conditions (species which did not show a decrease in abundance for any of the metrics were not included in the table). Signs indicate direction of future change based on correlation between flow metrics and climate data. X indicates species which will show declines in abundance with predicted future changes to flow metrics.

Scientific Name	Temperate Plains Ecoregion			Southern Appalachian Ecoregion	
	Variability in annual flows divided by median annual flows, where variability is calculated as interquartile (+)	High flow duration above 25th median flow (-)	Variability in Julian date of annual minimum (-)	Coefficient of variation in mean monthly flows (+)	Variability in annual maxima of 90-day means of daily discharge (+)
AMBLOPLITES RUPESTRIS	X			X	X
AMEIURUS MELAS			X		
APLODINOTUS GRUNNIENS		X			
CAMPOSTOMA PULLUM	X				
CARPIODES CARPIO		X			
CARPIODES CYPRINUS		X			X
CATOSTOMUS COMMERSIONII			X	X	X
COTTUS BAIRDII	X				X
CYPRINELLA SPILOPTERA					X
CYPRINELLA WHIPPLEI	X				
ETHEOSTOMA BLENNIOIDES	X				X
ETHEOSTOMA CAERULEUM	X				
ETHEOSTOMA NIGRUM		X	X		
ETHEOSTOMA STIGMAEUM					X
ETHEOSTOMA ZONALE					X
HYBOGNATHUS HANKINSONI		X	X		
HYBOPSIS AMBLOPS	X				
HYPENTELIUM NIGRICANS	X				X
ICTALURUS PUNCTATUS		X			
LEPOMIS MEGALOTIS	X				
LUXILUS CHRYSOCEPHALUS	X				X
LUXILUS CORNUTUS		X	X		X
MICROPTERUS DOLOMIEU					X
MOXOSTOMA ANISURUM		X			

Scientific Name	Temperate Plains Ecoregion			Southern Appalachian Ecoregion	
	Variability in annual flows divided by median annual flows, where variability is calculated as interquartile (+)	High flow duration above 25th median flow (-)	Variability in Julian date of annual minimum (-)	Coefficient of variation in mean monthly flows (+)	Variability in annual maxima of 90-day means of daily discharge (+)
MOXOSTOMA CARINATUM	X				
MOXOSTOMA DUQUESNEI	X				
MOXOSTOMA ERYTHRURUM		X			X
MOXOSTOMA MACROLEPIDOTUM		X			
NOTROPIS ATHERINOIDES					X
NOTROPIS DORSALIS		X	X		
NOTROPIS STRAMINEUS		X			
NOTROPIS VOLUCELLUS					X
NOTURUS FLAVUS		X			
PERCINA CAPRODES	X				X
PERCINA MACULATA		X	X		
PIMEPHALES NOTATUS					X
PIMEPHALES PROMELAS		X	X		
SANDER VITREUS		X			

Appendix C: Species scores and classification criteria.

Scientific Name	Response	Trait	Response			Trait			Rarity	Range	Dispersal	Resilience
	Score	Score	Flow	Temp	Habitat	Flow	Temp	Habitat				
AMBLOPLITES ARIOMMUS	3.12	4.65	0	0	1	1	1	0	1.53	0.77	0	0.42
AMBLOPLITES CONSTELLATUS	3.96	5.56	0	0	1	1	1	0	1.6	0.94	0	0.85
AMBLOPLITES RUPESTRIS	3.20	2.20	1	0	1	1	0	0	1	0.59	0	0.61
AMEIURUS MELAS	1.69	2.69	1	0	0	1	0	1	1	0.18	0	0.51
AMEIURUS NATALIS	0.60	1.60	0	0	0	1	0	0	1	0	0	0.6
AMEIURUS NEBULOSUS	2.87	4.86	0	0	0	1	0	0	1.99	0.94	0	1
APHREDODERUS SAYANUS	1.92	1.92	0	0	0	0	0	0	1.82	0.77	0	0.53
APLODINOTUS GRUNNIENS	2.48	2.48	1	0	0	1	0	0	1.54	0.24	0	0.58
CAMPOSTOMA OLIGOLEPIS	1.01	2.01	0	0	0	0	0	1	1	0.41	0	0.6
CAMPOSTOMA PULLUM	2.71	0.71	1	1	0	0	0	0	1	0.12	0	0.59
CARPIODES CARPIO	2.49	2.49	1	0	0	1	0	0	1.44	0.47	0	0.37
CARPIODES CYPRINUS	2.83	2.83	1	0	0	1	0	0	1.65	0.41	0	0.5
CARPIODES VELIFER	4.61	4.61	0	0	1	1	0	0	1.98	0.88	0	0.88
CATOSTOMUS COMMERSIONII	3.79	3.79	1	1	1	1	1	1	1	0.18	0	0.61
CENTRARCHUS MACROPTERUS		4.86				1	0	0	1.99	0.94	0	1
COTTUS BAIRDII	3.00	5.00	1	0	0	1	1	1	1	0.47	1	0.53
COTTUS CAROLINAE	3.04	4.04	0	0	1	1	0	1	1	0.41	1	0.63
COTTUS HYPSELURUS		6.22				1	1	1	1.22	0.65	1	0.55
CYPRINELLA CAMURA	2.47	2.47	0	0	0	0	0	0	1.99	0.94	0	0.6
CYPRINELLA GALACTURA	3.98	2.17	0	0	1	0	0	0	1.81	0.82	0	0.68
CYPRINELLA LUTRENSIS	0.86	0.86	0	0	0	0	0	0	1	0.3	0	0.56
CYPRINELLA SPILOPTERA	5.82	1.98	1	0	1	0	0	0	1.92	0.71	0	0.62
CYPRINELLA VENUSTA	1.67	3.42	0	0	0	0	1	0	1.75	0.65	0	0.54
CYPRINELLA WHIPPLEI	5.13	1.89	1	0	1	0	0	0	1.62	0.77	0	0.65
DOROSOMA CEPEDIANUM	0.67	2.67	0	0	0	1	1	0	1	0.06	0	0.61
ERIMYSTAX HARRYI		5.63				0	1	1	1.87	0.82	0	0.35
ERIMYSTAX X-PUNCTATUS	1.83	5.63	0	0	0	0	1	1	1.9	0.59	0	0.71
ERIMYZON OBLONGUS	2.85	5.63	0	0	1	1	1	1	1.39	0.65	0	0.56

Scientific Name	Response	Trait	Response			Trait			Rarity	Range	Dispersal	Resilience
	Score	Score	Flow	Temp	Habitat	Flow	Temp	Habitat				
ESOX AMERICANUS VERMICULATUS		4.06				1	0	0	1.85	0.77	0	0.79
ESOX NIGER	3.94	3.94	0	0	1	1	0	0	1.86	0.77	0	0.66
ETHEOSTOMA BLENNIOIDES	3.87	2.87	1	0	1	0	0	1	1	0.24	1	0.63
ETHEOSTOMA BURRI		7.87				0	1	1	1.76	0.94	1	0.93
ETHEOSTOMA CAERULEUM	3.99	3.99	1	0	1	0	1	1	1	0.35	1	0.64
ETHEOSTOMA CRAGINI		7.82				0	1	1	1.82	0.94	1	0.65
ETHEOSTOMA EUZONUM		8.11				0	1	1	1.99	0.94	1	0.27
ETHEOSTOMA FLABELLARE	2.97	3.97	0	0	1	0	1	1	1	0.3	1	0.67
ETHEOSTOMA GRACILE		7.15				0	1	1	1.94	0.59	1	0.19
ETHEOSTOMA JULIAE		8.48				0	1	1	1.93	0.94	1	0.87
ETHEOSTOMA MICROPERCA		7.71				0	1	1	1.97	0.82	1	0.18
ETHEOSTOMA NIANGUAE		8.75				0	1	1	1.99	0.94	1	0.91
ETHEOSTOMA NIGRUM	2.94	2.94	1	0	0	0	0	1	1	0.35	1	0.59
ETHEOSTOMA PROELIARE		7.34				0	1	1	1.95	0.77	1	0
ETHEOSTOMA PUNCTULATUM	3.34	4.34	0	0	1	0	1	1	1	0.65	1	0.69
ETHEOSTOMA SPECTABILE	1.88	3.88	0	0	0	0	1	1	1	0.24	1	0.64
ETHEOSTOMA STIGMAEUM	7.82	5.84	1	0	1	0	0	1	1.98	0.71	1	0.48
ETHEOSTOMA TETRAZONUM		6.87				0	1	1	1.68	0.71	1	0.64
ETHEOSTOMA UNIPORUM		7.72				0	1	1	1.95	0.94	1	0.03
ETHEOSTOMA ZONALE	5.00	3.74	1	0	1	0	0	1	1.26	0.47	1	0.63
FUNDULUS CATENATUS	2.88	1.88	0	0	1	0	0	0	1	0.24	1	0.64
FUNDULUS DISPAR		6.71				0	1	0	1.98	0.88	1	1
FUNDULUS NOTATUS	1.88	2.88	0	0	0	0	1	0	1	0.18	1	0.7
FUNDULUS OLIVACEUS	1.93	3.93	0	0	0	0	1	1	1	0.24	1	0.69
FUNDULUS SCIADICUS		3.17				0	0	0	1.72	0.65	1	0.34
GAMBUSIA AFFINIS	0.62	0.62	0	0	0	0	0	0	1	0	0	0.62
HYBOGNATHUS ARGYRITIS		1.89				0	0	0	1.97	0.88	0	0.15
HYBOGNATHUS HANKINSONI	3.47	3.47	1	0	0	0	1	0	1.78	0.88	0	0.12
HYBOGNATHUS PLACITUS		1.91				0	0	0	1.98	0.82	0	0.28
HYBOPSIS AMBLOPS	3.65	2.56	1	0	1	0	0	1	1.09	0.65	0	0.76

Scientific Name	Response	Trait	Response			Trait			Rarity	Range	Dispersal	Resilience
	Score	Score	Flow	Temp	Habitat	Flow	Temp	Habitat				
HYPENTELIUM NIGRICANS	3.02	4.02	1	0	1	1	1	1	1	0.35	0	0.67
ICHTHYOMYZON FOSSOR		6.44				0	1	1	1.99	0.88	0	0.7
ICHTHYOMYZON GAGEI		4.66				0	0	1	1.97	0.94	0	0.83
ICTALURUS PUNCTATUS	1.63	1.63	1	0	0	1	0	0	1	0.12	0	0.51
ICTIOBUS BUBALUS	1.15	2.94	0	0	0	1	0	0	1.79	0.35	0	0.52
ICTIOBUS CYPRINELLUS	1.51	3.32	0	0	0	1	0	0	1.81	0.41	0	0.76
LAMPETRA AEPYPTERA	2.07	5.91	0	0	0	0	1	1	1.92	0.88	0	0.37
LEPISOSTEUS OCULATUS		3.66				1	0	0	1.79	0.77	0	0.5
LEPISOSTEUS OSSEUS	0.82	2.22	0	0	0	1	0	0	1.4	0.18	0	0.57
LEPISOSTEUS PLATOSTOMUS	0.96	2.46	0	0	0	1	0	0	1.5	0.3	0	0.52
LEPOMIS CYANELLUS	0.61	1.61	0	0	0	1	0	0	1	0	0	0.61
LEPOMIS GULOSUS	1.93	1.93	0	0	1	1	0	0	1.19	0.18	0	0.53
LEPOMIS HUMILIS	0.70	1.70	0	0	0	0	0	1	1	0.12	0	0.58
LEPOMIS MACROCHIRUS	0.62	1.62	0	0	0	1	0	0	1	0	0	0.62
LEPOMIS MEGALOTIS	2.68	2.68	1	0	1	1	1	0	1	0	0	0.68
LEPOMIS MICROLOPHUS	0.91	2.21	0	0	0	1	0	0	1.3	0.18	0	0.68
LEPOMIS MINIATUS		3.29				1	0	0	1.63	0.71	0	0.51
LUXILUS CARDINALIS	3.45	4.90	0	0	1	0	1	1	1.45	0.88	0	0.72
LUXILUS CHRYSOCEPHALUS	2.89	1.89	1	0	1	0	0	1	1	0.3	0	0.59
LUXILUS CORNUTUS	2.83	2.83	1	0	0	0	0	1	1.27	0.71	0	0.66
LUXILUS PILSBRYI	3.54	4.92	0	0	1	0	1	1	1.38	0.94	0	0.86
LUXILUS ZONATUS		2.05				0	0	1	1	0.47	0	0.58
LYTHRURUS UMBRATILIS	0.66	0.66	0	0	0	0	0	0	1	0.18	0	0.48
MACRHYBOPSIS STORERIANA	1.71	5.69	0	0	0	0	1	1	1.99	0.71	0	0.31
MICROPTERUS DOLOMIEU	3.99	1.99	1	1	1	1	0	0	1	0.35	0	0.64
MICROPTERUS PUNCTULATUS	0.86	2.16	0	0	0	1	0	0	1.3	0.12	0	0.71
MICROPTERUS SALMOIDES	0.62	1.62	0	0	0	1	0	0	1	0	0	0.62
MINYTREMA MELANOPS	2.56	4.15	0	0	1	1	0	1	1.59	0.3	0	0.5
MORONE CHRYSOPS	1.77	3.69	0	0	0	1	0	0	1.92	0.53	0	0.75
MOXOSTOMA ANISURUM	3.83	7.55	1	0	0	1	1	1	1.86	0.71	0	0.66

Scientific Name	Response	Trait	Response			Trait			Rarity	Range	Dispersal	Resilience
	Score	Score	Flow	Temp	Habitat	Flow	Temp	Habitat				
MOXOSTOMA CARINATUM	6.12	6.12	1	0	1	1	0	1	1.93	0.82	0	0.67
MOXOSTOMA DUQUESNEI	3.04	3.04	1	0	1	1	0	1	1	0.35	0	0.69
MOXOSTOMA ERYTHRURUM	1.96	2.96	1	0	0	1	0	1	1	0.24	0	0.72
MOXOSTOMA MACROLEPIDOTUM	2.30	5.26	1	0	0	1	1	1	1.48	0.12	0	0.65
NOCOMIS ASPER	3.80	6.94	0	0	1	1	1	1	1.57	0.94	0	0.75
NOCOMIS BIGUTTATUS	2.00	4.00	0	0	1	1	1	1	1	0.41	0	0.59
NOTEMIGONUS CRYSOLEUCAS	0.70	1.70	0	0	0	1	0	0	1	0.06	0	0.64
NOTROPIS ATHERINOIDES	2.79	2.79	1	0	0	0	1	0	1.65	0.3	0	0.65
NOTROPIS BOOPS	2.03	3.03	0	0	1	0	1	1	1	0.41	0	0.62
NOTROPIS BUCCATUS	2.34	4.10	0	0	0	0	0	1	1.76	0.82	0	0.89
NOTROPIS BUCHANANI		3.90				0	0	1	1.96	0.71	0	0.56
NOTROPIS DORSALIS	2.05	2.05	1	0	0	0	1	0	1	0.59	0	0.46
NOTROPIS GREENEI		3.20				0	0	1	1.59	0.65	0	0.58
NOTROPIS HETEROLEPIS		2.33				0	0	0	1.95	0.77	0	0.84
NOTROPIS NUBILUS	2.00	3.00	0	0	1	0	1	1	1	0.35	0	0.65
NOTROPIS OZARCANUS		5.83				0	1	1	1.93	0.82	0	0.38
NOTROPIS STRAMINEUS	2.99	0.99	1	0	1	0	0	0	1	0.41	0	0.58
NOTROPIS TELESCOPUS	2.50	3.61	0	0	1	0	1	1	1.11	0.77	0	0.54
NOTROPIS TEXANUS		1.87				0	0	0	1.86	0.82	0	0.34
NOTROPIS VOLUCELLUS	3.10	1.46	1	0	0	0	0	0	1.64	0.47	0	0.69
NOTURUS ALBATER		4.68				1	0	1	1.43	0.82	0	0.64
NOTURUS EXILIS	1.93	3.93	0	0	1	1	1	1	1	0.24	0	0.69
NOTURUS FLAVATER		6.16				1	0	1	1.98	0.88	0	0.45
NOTURUS FLAVUS	4.61	6.33	1	0	1	1	1	1	1.72	0.35	0	0.56
NOTURUS GYRINUS	1.57	3.30	0	0	0	0	0	1	1.73	0.59	0	0.55
NOTURUS MIURUS	2.20	4.11	0	0	0	0	0	1	1.91	0.82	0	0.63
NOTURUS NOCTURNUS	1.40	7.34	0	0	0	1	1	1	1.98	0.59	0	0.23
OPSOPOEODUS EMILIAE		1.34				0	0	0	1.89	0.71	0	0.01
PERCINA CAPRODES	3.79	2.79	1	0	1	0	0	1	1	0.12	1	0.67
PERCINA COPELANDI	4.43	6.40	0	0	0	0	0	1	1.97	0.94	1	0.6

Scientific Name	Response	Trait	Response			Trait			Rarity	Range	Dispersal	Resilience
	Score	Score	Flow	Temp	Habitat	Flow	Temp	Habitat				
PERCINA CYMATOTAENIA		7.88				0	1	1	1.98	0.88	1	0.19
PERCINA EVIDES	5.98	7.92	0	0	1	0	1	1	1.94	0.71	1	0.73
PERCINA MACULATA	6.17	6.17	1	0	1	0	1	1	1.59	0.53	1	0.56
PERCINA PHOXOCEPHALA	4.76	6.39	0	0	1	0	1	1	1.63	0.53	1	0.64
PERCINA SCIERA	3.90	5.89	0	0	0	0	0	1	1.99	0.77	1	0.39
PHENACOBIOUS MIRABILIS	1.84	1.84	0	0	1	0	0	1	1	0.35	0	0.49
PHOXINUS ERYTHROGASTER	1.95	1.95	0	0	1	0	1	0	1	0.35	0	0.6
PIMEPHALES NOTATUS	1.61	0.61	1	0	0	0	0	0	1	0	0	0.61
PIMEPHALES PROMELAS	1.64	0.64	1	0	0	0	0	0	1	0.18	0	0.46
PIMEPHALES VIGILAX	1.66	1.66	0	0	0	0	0	0	1.72	0.59	0	0.65
POMOXIS ANNULARIS	0.62	1.97	0	0	0	1	0	0	1.35	0.06	0	0.54
POMOXIS NIGROMACULATUS	1.17	3.04	0	0	0	1	0	0	1.87	0.24	0	0.73
PYLODICTIS OLIVARIS	0.96	2.60	0	0	0	1	0	0	1.64	0.3	0	0.48
SANDER VITREUS	4.22	6.28	1	0	0	1	1	0	1.99	0.77	0	0.7
SEMOTILUS ATROMACULATUS	0.65	2.65	0	0	0	0	1	1	1	0.06	0	0.59