Assessing the infectious reservoir of falciparum malaria: past and future

Will Stone¹, Bronner P. Gonçalves², Teun Bousema^{1, 2}, Chris Drakeley²

1. Radboud University Medical Center, Nijmegen, the Netherlands

2. London School of Hygiene & Tropical Medicine, London, UK

Corresponding author: Drakeley, C. (chris.drakeley@lshtm.ac.uk)

Keywords: *Plasmodium falciparum*, gametocytes, infectious reservoir, mosquito, transmission blocking

Abstract

Renewed interest in malaria eradication has placed greater emphasis on the development of tools to interrupt *Plasmodium* transmission, such as transmission-blocking vaccines. However, effective deployment of such tools will likely depend on improving our understanding of which individuals transmit infections to mosquitoes. To date, only a handful of studies have directly determined the infectiousness of individuals in endemic populations. Here we review these studies and their relative merits. We also highlight factors influencing transmission potential that are not normally considered; the duration of human infectiousness, frequency of sampling by mosquitoes, and variation in vector competence among different mosquito populations. We argue that more comprehensive xenodiagnostic assessments of infectivity are necessary to accurately quantify the infectious reservoir and better target interventions.

Reducing transmission of malaria: the future of elimination programs

The burden of malaria has declined in many endemic settings in Africa and elsewhere (1). Local malaria elimination is considered achievable with current control approaches in some of these areas when transmission intensity is low and re-introduction unlikely (2, 3). However, in the majority of endemic areas, operational and technical limitations are likely to hinder the complete interruption

of transmission (4, 5). New or renewed tools aimed specifically at interrupting the spread of *Plasmodium* species from human to mosquito may therefore be critical for future elimination programs (6, 7), especially if transmission efficiency increases as parasite prevalence goes down (8). Unlike traditional control strategies, which aim to reduce severe morbidity in vulnerable populations, the effectiveness of transmission reducing interventions (TRI) hinges on their coverage of individuals responsible for transmission to mosquitoes regardless of their symptomatic status (9). Despite the central importance of human infectivity for TRI (10), there have been few direct assessments of human infectivity to mosquitoes. Here we discuss previous studies that aimed to directly assess infectivity at the population level, examine factors necessary to link these controlled transmission experiments with transmission in nature, and advocate the next steps which will provide the key information necessary to better target the human infectious reservoir for malaria.

Quantifying the human infectious reservoir of malaria (1957 – 2014)

In infected humans, the life-cycle of a small portion of the total malaria parasite population culminates in differentiation and maturation into gametocytes (see Glossary). When an anopheline mosquito feeds on blood containing mature gametocytes, some may be ingested which then fuse and undergo sporogonic development making the insect infectious to humans. The human component of the infectious reservoir is the proportion of a population that are capable of infecting mosquitoes (11).

In community surveys, data from microscopy typically show that children and infants harbour *Plasmodium* gametocytes more commonly and in greater number than older age groups (12). Such observations have long galvanised belief among the scientific community that young individuals represent the main source of infection for mosquitoes (12). However, in the late 1940s and early 1950s, xenodiagnostic assessments revealed that the presence of gametocytes in blood films was not prerequisite to onward transmission (12, 13, 14, 15). The first population-based assessment of human infectivity to mosquitoes was conducted in rural Liberia, where individuals living in an

endemic region were recruited for mosquito feeding experiments regardless of their parasite status(16). Whilst this study found that young children were much more infectious to mosquitoes, it also showed that when estimates were adjusted for the demographic composition of the population, all age groups were relevant contributors to malaria transmission (Box 1). In nearly 60 years since this study a handful similar surveys have been reported (Tables 1 and 2). In general, these studies show that whilst young children are consistently more likely to be infectious *per se*(17, 18), the contribution of older age groups is by no means negligible; in previous xenodiagnostic surveys that were able to sample individuals of all ages, individuals >15 years old made up between 21.9 and 40.7% of the total infectious reservoir(16, 17, 18, 19). Several studies have assessed human-mosquito transmission using entomological parameters (20). However, these studies provide only broad population estimates of infectiousness and they are neither able to describe the distribution of the reservoir among different members of the population nor allow an examination of the role of parasite density and infection dynamics.

Ultimately, determining the infectiousness of different age groups relative to their representation in the population is just one necessary step in characterising the infectious reservoir. Contribution to transmission is influenced not only by intrinsic parasite and human determinants of infectivity, reflected in the outcome of mosquito feeding assays, but by a variety of additional factors that make the capacity to infect mosquitoes in controlled experiments distinct from the probability of onward transmission in nature (*e.g.* human exposure to mosquito biting) (Figure 1). These factors are not commonly taken into account and are not uniform across time, space, or between individuals of different age. To more accurately reflect transmission in its epidemiological context, these elements need to be considered and incorporated into future assessments.

Determinants of human transmission potential: the need for data

Duration and dynamics of infectivity

The likelihood of transmission to *Anopheles* mosquitoes is determined primarily by gametocyte density, fitness and their circulation time. The non-linear relationship between gametocyte density and mosquito infection risk has been described several times (21, 22, 23). Yet, because of the relatively high proportion of low density parasitaemias and gametocytaemias in natural infections, there is increasing focus on how important these often sub-patent infections are for the maintenance of transmission (24, 25). While the limited sensitivity of microscopy to detect gametocytes is well established (21), many claims of submicroscopic gametocyte densities contributing to transmission are based on microscopy-positive asexual parasite carriers with accompanying submicroscopic gametocyte densities. This is an important difference when considering how current diagnostics may capture the human infectious reservoir (26, 27); infections with submicroscopic gametocytes that accompany patent asexual parasites are detectable by conventional diagnostics whilst infections with no microscopically detectable parasite stages require more sensitive diagnostics (28).

Because of their cross-sectional nature, previous xenodiagnostic surveys have not formally examined the duration of infectiousness, another key factor influencing transmission potential. Data from malaria therapy studies suggest infectiousness of an individual infection can last for many months (29). However, it is difficult to extrapolate the observations from a limited age range of malaria naïve individuals to endemic populations. Broadly, in endemic settings, children (possessing limited immunity) are likely to have more acute, higher density infections that often require treatment (30). This may result in cross sectional surveys missing recently treated infections and identifying only a limited number of highly infectious children without recent treatment. Conversely, adults and older children (often semi-immune, and presenting few clinical symptoms) would have longer, chronic infections, which could result in longer infectious periods, balanced by lower infectivity at individual time points (31). These patterns of infection are likely to differ with transmission intensity. In areas of intense malaria transmission children may have repeated malaria infections with relatively high gametocyte densities and a high likelihood of malaria transmission (Figure 2). In

106 areas with lower transmission, including areas that have reduced transmission in the context of an
107 elimination program, infection duration will likely have a more significant influence on the infectious
108 reservoir. At present only one study has directly assessed population level infectiousness in an area
109 of low malaria transmission(32). Because of the utility of these data to control and elimination
110 programs, further studies assessing the infectious reservoir in areas of low transmission would be
111 advantageous.

112 In an individual infection, infectiousness per gametocyte may be higher at the beginning of an
113 infection than at the end because of changes in parasite competency (33), sex ratio (34, 35) and
114 possibly the development of sexual stage immune responses(36, 37, 38). Transmission-blocking
115 immunity is boosted by recent exposure to gametocytes (37) and may reduce infectiousness during
116 the course of infections that last several weeks or months. Superinfecting clones influence parasite
117 growth dynamics including total gametocyte biomass (39), and multiplicity of infection could in itself
118 be a determinant of transmissibility (40). This combination of factors might also indicate that
119 infectiousness of one infection may not reflect the infectiousness of subsequent infections (when
120 multiplicity of infection and transmission blocking immunity may differ), suggesting greater
121 complexity than that estimated from a single time point during the transmission season.

123 *Mosquito exposure*

124 Artificial feeding experiments in xenodiagnostic surveys have tended to use fixed numbers of
125 mosquitoes, improving the precision of estimates of infectiousness by maximising mosquito sample
126 sizes. However, in any endemic area mosquito biting rates of humans vary considerably. As an
127 example, a recent study in Uganda found biting rates varied from 270 to 7399 bites/year between
128 study sites, with each site showing distinct seasonal fluctuations (41). The transmission potential of
129 infectious individuals will be determined by the rate at which they are sampled by local mosquitoes
130 and the competence of these vectors. Where vector densities and survival rates are low the number
131 of mosquitoes that become infected and subsequently transmit their infections will be very low even

if there are infectious individuals in the human population. Although essential to accurately characterize individual-specific contributions to transmission, these entomological parameters have never been measured in conjunction with xenodiagnostic surveys.

Attractiveness to mosquitoes

In the early 1950s, Muirhead-Thomson observed that Jamaican adults attracted more *Anopheles* mosquitoes than equally exposed children (42). The conclusions of this study are supported by later reports in which wild *Anopheles* were allowed to feed on unprotected humans under observation (43, 44), or ABO blood typing allowed the identification of blood meal origin (45), though some studies show more random biting selection (46). Carnevale *et al* near Brazzaville, in the Republic of Congo, described the biting behaviours of ~6500 mosquitoes on 24 individuals of all ages (43). *Anopheles* biting increased with age, with individuals >20 years old being bitten three times more than <2 year olds. Port *et al.* were able to ascribe this age-related biting pattern to increased body weight and surface area (45). In addition to increased surface area, body temperature and chemical cues change with age and physiological maturity and may be associated with attractiveness (47, 48, 49). This might also help explain why pregnant women are thought to be more attractive to mosquitoes than non-pregnant women (50). Muirhead-Thomson also observed that infants reacted more actively in sleep to the probing of mosquitoes than adults, and that mosquitoes were less likely to feed on infants even after alighting on their skin (42). Data are less robust on the influence of *Plasmodium* infection on mosquito attractiveness. Reports from animal models and studies on a small number of humans infected with *P. falciparum* suggest that the presence of gametocytes may increase attractiveness to anopheline mosquitoes (51, 52), possibly due to the production of attractive volatiles (53). It is unclear if the small biomass of gametocytes could have specific systemic effects that would increase the attractiveness of children (harbouring higher gametocyte densities), and if this might counteract the reduced attraction associated with their smaller surface area.

158

159 *Availability to mosquitoes*

160 However attractive and infectious the individual, they must be available to mosquitoes for feeding
161 and subsequent transmission to occur. This availability is shaped primarily by human social and
162 behavioural practices, including use of mosquito nets and sleeping patterns.

163 Age-specific net use depends on the setting and extent of ownership, but in areas with good
164 coverage appears highest among young children (<5 years), and lowest among older children and
165 adolescents (5-15 years)(54, 55, 56, 57). In addition to bed-net's community-wide effect on
166 transmission, they may also reduce an individual's chance of infecting mosquitoes and divert vectors
167 to other bloodmeal sources. In the Gambia, the introduction of insecticide-treated nets (ITN)
168 protected children against malaria and diverted mosquito bites to other hosts (animals or adults)
169 (58). Net distribution has also been linked to changes in vector behaviour and species composition;
170 *Anopheles gambiae* and *Anopheles funestus* have showed signs of responsive exophagy(59), and
171 there are observations that peak biting in these species may shift to the early evening and morning
172 (60, 61). Net-related reductions in *A. gambiae*s. populations have also been linked to increasingly
173 dominant roles for *A. funestus*(62) and *Anopheles arabiensis*(63).

174 These elements together with sleeping times and time spent indoors or outdoors at night determine
175 availability to different mosquito populations. Children might be expected to sleep earlier than
176 adults, but the protection afforded by long periods of time under bed-nets depends on when local
177 vectors are most active. Data on time spent indoors, time spent under nets, and on the biting activity
178 of local vectors can be combined to determine the true protective efficacy of bed nets by age(64).
179 This is an informative composite measure, which could be used to estimate age-specific patterns of
180 mosquito exposure.

181

182 Effectively delineating the interaction of mosquito exposure and subsequent malaria transmission is
183 difficult. Individuals identified as infectious in xenodiagnostic surveys must have been exposed to

malara vectors at some stage, but heterogeneity in attractiveness and the degree of exposure should still significantly influence their relative contribution to the infectious reservoir. Surveys quantifying mosquito exposure and infectiousness in the same population could provide more realistic estimates of age-specific contributions to transmission.

Heterogeneous Mosquito Populations

Although several vector species might contribute to local transmission in endemic areas, most xenodiagnostic surveys have been performed using *A. gambiae*s.s.(Tables 1 and 2), the majority of which were colony reared.Evidence is limited on whether infectiousness to *A. gambiae* during membrane or skin feeding assays corresponds to infectivity to other vector species.Two major malaria vectors, *A. funestus* and *A. arabiensis*, are distributed throughout Sub-Saharan Africa, often co-existing with *A. gambiae*s.s.(65). The highly anthropophilic biting preferences of *A. funestus*(66),its high longevity(67)and sporozoite rate(62, 68) (higher than *A. gambiae*in some endemic areas) indicate that a non-negligible part of malaria transmission might be sustained by this vector where it is present.Conversely,*A. arabiensis* is considered more exophilic and exophagic; in Burkina Faso, *A. arabiensis* was as susceptible to infection as *A. gambiae*s.s.in feeding assays(69), while in Senegal, lower infection rates were observed in membrane feeding assays with *A. arabiensis* compared to *A. gambiae*s.s.(70).The recent description of cryptic malaria vectors (71, 72) and the conflicting results on the differential susceptibility of *A. gambiae*s.s.and *Anopheles coluzzi*(69, 70)(previously the molecular formsS and M of *A. gambiae*,respectively(73)), further illustrate the need to quantify transmission efficiency against the background of diverse vector populations.

Considerations for future studies

To more rigorously assess the human infectious reservoir using direct xenodiagnosticsurveys,severalmethodological issues require consideration. In this section, we suggest suitable inclusion criteria andmethods for measuring or predicting individual level mosquito

exposure. Other important factors that are discussed include the type of feeding assay, measures of infectiousness, and general study design.

Inclusion criteria for direct assessments of the infectious reservoir

There is abundant evidence that microscopy is insufficiently sensitive to detect low densities of asexual parasites and gametocytes (15, 19, 74, 75, 76, 77, 78). Studies including only individuals with patent gametocytes (11, 19, 79) are likely to have underestimated the proportion of populations capable of infecting mosquitoes, and biased measures of mosquito infection probability (Table 2). Since low density gametocyte carriage is prevalent in many endemic areas (25) and they may represent a major source of secondary mosquito infections (22, 80, 81), studies aiming to characterize the human infectious reservoir should not have gametocyte positivity by microscopy as an inclusion criterion.

Defining infectiousness at the population level

The simplest method for quantifying the infectious reservoir combines the prevalence of people in different age groups infecting at least one mosquito, and the proportion of those groups in the total population (16, 77). Alternatively, population level infectiousness may be presented as mosquito infection probability (K); the likelihood of a mosquito becoming infected after feeding on any member of the population, calculated as the age-adjusted mosquito infection rate. Although the former metric has value for efforts that aim to identify or target all individuals who contribute to onward malaria transmission, the latter (mosquito infection probability) makes estimates more comparable between studies, where the number of mosquitoes used in feeding assays may differ widely (18, 19), and captures variations in infectiousness intensity in different age groups that may not be apparent when reporting only infectiousness prevalence.

Assay methodology

There are two types of mosquito infection assay that have been used in previous surveys: direct skin feeding and membrane feeding assays (Table 1 & 2). Direct feeds better reflect natural infection and avoid some of the technical challenges that limit the wide-scale use of membrane feeding assays (e.g. adapting mosquito colonies to membrane feeding, preventing gametocyte activation between bleeding and feeding and a potential effect of anticoagulants)(82, 83). Although their use is currently hindered by ethical concerns that have precluded their use in repeated assessments of mosquito infection from young children (82), further direct comparisons of xenodiagnostic surveys using skin and membrane feeding experiments would be highly valuable to parameterise the extent to which the latter reflects the natural situation. Unless direct skin feeding is acceptable in all age groups and with sufficient mosquito numbers to allow precise estimates of infection, studies aiming to characterise the full infectious reservoir in endemic populations should do so using direct membrane feeding assays using locally relevant malaria vectors.

Whether feeding assays measure infection by detection of oocysts in the mosquito gut or sporozoites in the salivary glands (Table 1 & 2) should make little difference to the comparability of future studies, as even in low intensity infections (such as those occurring commonly in nature) there is a close correlation between oocyst prevalence and the prevalence of subsequent salivary gland infection (84). Salivary gland dissections are technically even more demanding than midgut dissections, will result in higher mosquito mortality between the day of feeding and the day sporozoites are first detectable and impose additional safety measures for insectaries that need to be equipped for holding mosquitoes that form a biohazard to insectary personnel. We therefore suggest that detection of *Plasmodium* oocysts is the most viable measure of mosquito infection.

Measuring mosquito exposure

As outlined earlier, the actual number of mosquito bites an individual receives is influenced by their physical characteristics and behaviour(42, 45). There is also spatial heterogeneity in exposure, with individuals living in households with higher vector densities (e.g. houses nearer to breeding sites)

receiving more bites and having greater risk of disease than those in households where vector density is lower(85). Infectious reservoir assessmentsshould ideally use a direct approach to quantify mosquito biting rates, simultaneously accounting for the availability and attractiveness of human hosts to mosquitoes: collection of mosquitoes from households and identification of blood meal sources by DNA fingerprinting would provide reliable estimates of true exposure to mosquito bites. A recent study in Peru used this approach to show that adults were more often bitten by *Aedes aegypti* mosquitoes compared to children living in the same houses (86). We suggest that in areas where transmission is dominated by indoor biting mosquitoes, the rate of secondary mosquito infections could be calculated by multiplying individual-specific mosquito biting rates (determined experimentally using the above methods) by infectiousness (the probability that a mosquito develops infection after feeding, determined with mosquito feeding assays). This measure would be informative as it accounts for all major factors influencing transmission potential. An obvious problem with this approach is that only endophilic mosquitoes are sampled and there are currently no standardized sampling strategies for outdoor biting vectors.

Where direct assessments are operationally impossible, or made unreliable by vector behaviour, we suggest that daily mosquito biting rates could be estimated for each study subject by collating mosquito (*e.g.* abundance, estimated indoor and outdoor biting rate), demographic (*e.g.* age, body surface area and attractiveness, household conditions, distance to breeding sites, number of people living in the same house/room) and behavioural (*e.g.* sleeping times, net use) data, so that a rate of secondary mosquito infections could be broadly estimated by combination with experimentally-determined infectiousness. Figure 1 is a simplified and hypothetical representation of the potential influence of some of these factors on the contribution of different age groups to the infectious reservoir.

Longitudinal studies with mosquito exposure estimation

Longitudinal studies with frequent follow-up would be valuable not only to estimate the duration of infectiousness in different age groups but also to assess whether some individuals or groups of individuals are infectious throughout a transmission season. This approach would help determine if transmissibility is associated with clusters of malaria higher infection incidence, *i.e.* where mosquito-to-human transmission is high. The more complete interpretation of such assessments could be made with concurrent mosquito biting rates.

Concluding remarks

We acknowledge that there are several malaria endemic settings in which progress toward elimination has been made in the absence of detailed local knowledge of the infectious reservoir. However, malaria remains a threat in many countries, particularly in Africa, and resurgence due to natural receptivity for transmission, drug and insecticide resistance means that a better understanding of who within a community infects mosquitoes can only benefit malaria control. The complexity of factors influencing malaria transmission means that studies aiming to fully understand how this occurs is challenging. Previous xenodiagnostic surveys provided valuable insights into malaria transmission epidemiology, but these studies need to be extended to simultaneously evaluate additional key factors. Moreover, the longitudinal nature of infectiousness during natural malaria infections needs to be considered in future studies, as this will influence the contribution of different age groups to transmission.

While human and parasite factors, including the intensity and duration of gametocytaemia, affect infectiousness at the individual level, entomological factors dictate the transmission potential of each individual and the infectious reservoir as whole. Therefore, mosquito exposure assessments need to be an integral part of xenodiagnostic surveys if we are to effectively target the reservoir of infection.

Box 1. Demography and the malaria infectious reservoir

In previous surveys, the 'bottom-heavy' age structure typical of malaria endemic regions has acted to boost the contribution of more infectious children to the total malaria infectious reservoir, while less infectious adults, representing approximately half of most populations, generally contributed to a lesser extent. Recent demographic estimations show that Sub-Saharan African populations are still disproportionately young; approximately 17% and 27% of the population are aged between 0-4 and 5-14 years, respectively (<http://esa.un.org/wpp/>). The influence of demography on patterns of transmission needs to be considered in light of the numerous other factors affecting the likelihood of infectious individuals contributing to mosquito infections. Though infants and children are over-represented in endemic areas they may be the age group least available to and, perhaps, least attractive to mosquitoes (see *Determinants of human transmission potential: the need for data*). Conversely, older children and adolescents may represent a privileged group for human-mosquito transmission; individuals in this age group possess moderate gametocyte densities, approximately double the infectivity of individuals >15 years, lack the severe symptoms that prompt treatment, and appear comparatively vulnerable to mosquito biting (Figure 1).

Women in pregnancy have been observed to attract more than twice as many anophelines as non-pregnant women over short distances [21]. However, such observations must be weighed against behavioural and demographic parameters unique to this group. Census and fertility rate data show that pregnant women comprise a small proportion of African populations. In Zambia, 8.7% of women of child bearing age were pregnant at any one time in 2013, which equates to 2-2.5% of the entire population (<http://www.dhsprogram.com/pubs/pdf/FR304/FR304.pdf>). Pregnant women are also preferentially targeted by net and drug treatment campaigns, so the high attractiveness to mosquitoes of this small proportion of the population may be balanced by interventions.

Box 2. Outstanding questions

- If and how does human population infectiousness to mosquitoes differ in areas of low malaria transmission? Are traditionally low endemic settings different from those under epidemiological transition (i.e. those settings moving from moderate to low transmission as a result of heightened control activities)?
- What is the duration of infectiousness of individuals to mosquitoes? Are certain people infectious for longer periods and, if so, what are their characteristics?
- How do fluctuations in mosquito densities and feeding rates affect human infectiousness?

Glossary

Endophilic: Arthropod behaviour describing a preference for resting indoors after blood-feeding.

Exophagic: Arthropod behaviour describing preference for blood-feeding outdoors.

Exophilic: Arthropod behaviour describing preference for resting outdoors after blood-feeding outdoors

Gametocyte: The sexual stages of the malaria parasite capable of reproduction in the mosquito.

Female (macro-) and male (micro-) gametocytes circulate in the human peripheral blood, where they may be ingested by blood-feeding *Anopheles* mosquitoes and begin sexual development.

Mosquito feeding assay: Xenodiagnostic assays developed to determine the infectiousness of *Plasmodium* gametocytes to *Anopheles* mosquitoes. Mosquito feeding assay (MFA) may refer to skin feeding assays, in which mosquitoes are allowed to feed directly on a subject's skin; to direct membrane feeding assays (DMFA), in which mosquitoes feed on venous blood from a subject maintained at body temperature in a membrane feeding device; or to standard membrane feeding assays (SMFA), in which mosquitoes feed on cultured gametocytes in a membrane-based feeder system.

Perennial transmission: Malaria transmission may be described as perennial where environmental and climatic conditions allow parasite transmission throughout the year.

Seasonal transmission: Malaria transmission may be described as seasonal where mosquito biting and thus transmission are clustered temporally, generally occurring following predictable periods of rain and subsequently decreasing in periods of drought.

Transmission blocking immunity: The development of humoral immunity to antigens present during gametocyte development. Antibodies specific to gametocyte antigens may be ingested along with mature gametocytes during blood-feeding. If these antibodies target to proteins essential for

parasite development (e.g. Pfs48/45, Pfs230) and are sufficiently abundant, antibody interaction can prevent parasite development and transmission potential is reduced or entirely blocked. There is a growing body of evidence for the development of transmission blocking immunity in individuals naturally exposed to malaria infection

Xenodiagnosis: In this context is a method to determining the infectiousness of a potentially parasitised host (i.e. malaria infected humans) by allowing vectors (i.e. mosquitoes) to feed on the subject. Infectiousness is assessed by the infection status of the vector (i.e. mosquito) after a suitable period of parasite development.

Acknowledgements

The authors would like to acknowledge the Bill & Melinda Gates Foundation for funding (grant OPP1034789).

References

1. World Health Organisation (2013) World Malaria Report.
2. Aboobakar, S. *et al.* (2012) Eliminating malaria and preventing its reintroduction: the Mauritius case study. *Malar J.* 11, O12.
3. Shamuradova, L. *et al.* (2012) Achieving malaria elimination and certification in Turkmenistan. *Malar J.* 11, O11.
4. Alonso, P.L. *et al.* (2011) A Research Agenda to Underpin Malaria Eradication. *PLoS Med* 8, e1000406.
5. Griffin, J.T. *et al.* (2010) Reducing *Plasmodium falciparum* Malaria Transmission in Africa: A Model-Based Evaluation of Intervention Strategies. *PLoS Med* 7, e1000324.
6. Targett, G. and Greenwood, B. (2008) Malaria vaccines and their potential role in the elimination of malaria. *Malar J.* 7, S10.
7. Gosling, R.D. *et al.* (2011) The role of antimalarial treatment in the elimination of malaria. *Clinical Microbiology and Infection* 17, 1617-1623.
8. Churcher, T.S. *et al.* (2015) Human-to-mosquito transmission efficiency increases as malaria is controlled. *Nat Commun* 6.
9. Vannice, K. *et al.* (2012) MALVAC 2012 scientific forum: accelerating development of second-generation malaria vaccines. *Malar J.* 11, 372.
10. Muirhead-Thomson, R.C. (1998) Where do most mosquitoes acquire their malarial (*Plasmodium falciparum*) infection? From adults or from children? *Ann. Trop. Med. Parasitol.* 92, 891-893.
11. Drakeley, C.J. *et al.* (2000) Estimates of the infectious reservoir of *Plasmodium falciparum* malaria in The Gambia and in Tanzania. *Trans R Soc Trop Med Hyg* 94, 472-476.
12. Shute, P.G. and Maryon, M. (1951) A study of gametocytes in a west african strain of *Plasmodium falciparum*. *Trans R Soc Trop Med Hyg* 44, 421-438.
13. Draper, C.C. (1953) Observations on the infectiousness of gametocytes in hyperendemic malaria. *Trans R Soc Trop Med Hyg* 47, 160-165.
14. Young MD *et al.* (1948) The infectivity of native malarias in South Carolina to *Anopheles quadrimaculatus*. *Am J Trop Med Hyg* 28, 303-311.

- 412 15. Muirhead-Thomson, R.C. (1954) Low Gametocyte Thresholds of Infection of *Anopheles* with
413 *Plasmodium falciparum*. *BMJ* 1, 68-70.
- 414 16. Muirhead-Thomson, R.C. (1957) The malarial infectivity of an African village population to
415 mosquitoes (*Anopheles gambiae*); a random xenodiagnostic survey. *Am J Trop Med Hyg* 6,
416 971-979.
- 417 17. Githeko, A.K. *et al.* (1992) The reservoir of *Plasmodium falciparum* malaria in a holoendemic
418 area of western Kenya. *Trans R Soc Trop Med Hyg* 86, 355-358.
- 419 18. Bonnet, S. *et al.* (2003) Estimation of malaria transmission from humans to mosquitoes in
420 two neighbouring villages in south Cameroon: evaluation and comparison of several indices.
421 *Trans R Soc Trop Med Hyg* 97, 53-59.
- 422 19. Graves, P.M. *et al.* (1988) Measurement of malarial infectivity of human populations to
423 mosquitoes in the Madang area, Papua New Guinea. *Parasitology* 96, 251-263.
- 424 20. Killeen, G.F. *et al.* (2006) Infectiousness of Malaria-Endemic Human Populations to Vectors.
425 *Am J Trop Med Hyg* 75, 38-45.
- 426 21. Bousema, T. and Drakeley, C. (2011) Epidemiology and Infectivity of *Plasmodium falciparum*
427 and *Plasmodium vivax* Gametocytes in Relation to Malaria Control and Elimination. *Clin.*
428 *Microbiol. Rev.* 24, 377-410.
- 429 22. Churcher, T.S. *et al.* (2013) Predicting mosquito infection from *Plasmodium falciparum*
430 gametocyte density and estimating the reservoir of infection. *eLife* 2.
- 431 23. Graves, P.M. (1980) Studies on the use of a membrane feeding technique for infecting
432 *Anopheles gambiae* with *Plasmodium falciparum*. *Trans R Soc Trop Med Hyg* 74, 738-742.
- 433 24. Lin, J.T. *et al.* (2014) The role of submicroscopic parasitemia in malaria transmission: what is
434 the evidence? *Trends Parasitol.* 30, 183-190.
- 435 25. Okell, L.C. *et al.* (2012) Factors determining the occurrence of submicroscopic malaria
436 infections and their relevance for control. *Nature Communications* 3.
- 437 26. Drakeley, C. *et al.* (2006) The epidemiology of *Plasmodium falciparum* gametocytes: weapons
438 of mass dispersion. *Trends Parasitol.* 22, 424-430.
- 439 27. Bousema, T. *et al.* (2014) Asymptomatic malaria infections: detectability, transmissibility and
440 public health relevance. *Nat Rev Micro* 12, 833-840.
- 441 28. Hofmann, N. *et al.* (2015) Ultra-Sensitive Detection of *Plasmodium falciparum* by
442 Amplification of Multi-Copy Subtelomeric Targets. *PLoS Med* 12, e1001788.
- 443 29. Jeffery, G.M. and Eyles, D.E. (1955) Infectivity to Mosquitoes of *Plasmodium falciparum*as
444 Related to Gametocyte Density and Duration of Infection. *Am J Trop Med Hyg* 4, 781-789.
- 445 30. Carneiro, I. *et al.* (2010) Age-Patterns of Malaria Vary with Severity, Transmission Intensity
446 and Seasonality in Sub-Saharan Africa: A Systematic Review and Pooled Analysis. *PLoS ONE* 5,
447 e8988.
- 448 31. Laishram, D. *et al.* (2012) The complexities of malaria disease manifestations with a focus on
449 asymptomatic malaria. *Malar J.* 11, 29.
- 450 32. Gaye, A. *et al.* (2015) Infectiousness of the Human Population to *Anopheles arabiensis* by
451 Direct Skin Feeding in an Area Hypoendemic for Malaria in Senegal. *Am J Trop Med Hyg* 92,
452 648-652.
- 453 33. Lensen, A. *et al.* (1999) *Plasmodium falciparum*:Infectivity of Cultured, Synchronized
454 Gametocytes to Mosquitoes. *Experimental Parasitology* 91, 101-103.
- 455 34. Sowunmi, A. *et al.* (2008) *Plasmodium falciparum* gametocyte sex ratios in children with
456 acute, symptomatic, uncomplicated infections treated with amodiaquine. *Malar J.* 7, 169.
- 457 35. Gbotosho, G.O. *et al.* (2011) *Plasmodium falciparum* gametocyte carriage, emergence,
458 clearance and population sex ratios in anaemic and non-anaemic malarious children. *Mem.*
459 *Inst. Oswaldo Cruz* 106, 562-569.
- 460 36. Bousema, J.T. *et al.* (2007) A longitudinal study of immune responses to *Plasmodium*
461 *falciparum* sexual stage antigens in Tanzanian adults. *Parasite Immunol.* 29, 309-317.

- 462 37. Bousema, T. *et al.* (2010) The Dynamics of Naturally Acquired Immune Responses to
463 *Plasmodium falciparum* Sexual Stage Antigens Pfs230 & Pfs48/45 in a Low Endemic Area
464 in Tanzania. *PLoS ONE* 5, e14114.
- 465 38. Ouédraogo, A.L. *et al.* (2011) Naturally Acquired Immune Responses to *Plasmodium*
466 *falciparum* Sexual Stage Antigens Pfs48/45 and Pfs230 in an Area of Seasonal Transmission.
467 *Infection and Immunity* 79, 4957-4964.
- 468 39. Bruce, M.C. and Day, K.P. (2002) Cross-species regulation of malaria parasitaemia in the
469 human host. *Curr. Opin. Microbiol.* 5, 431-437.
- 470 40. Nsango, S.E. *et al.* (2012) Genetic clonality of *Plasmodium falciparum* affects the outcome of
471 infection in *Anopheles gambiae*. *Int. J. Parasitol.* 42, 589-595.
- 472 41. Kilama, M. *et al.* (2014) Estimating the annual entomological inoculation rate for *Plasmodium*
473 *falciparum* transmitted by *Anopheles gambiae* s.l. using three sampling methods in three
474 sites in Uganda. *Malar J.* 13, 111.
- 475 42. Muirhead-Thomson, R.C. (1951) Distribution of Anopheline Mosquito Bites Among Different
476 Age Groups. *BMJ* 1, 1114-1117.
- 477 43. Carnevale, P. *et al.* (1978) The aggressiveness of *Anopheles gambiae* A in relation to the age
478 and sex of the human subjects. *Bull. World Health Organ.* 56, 147-154.
- 479 44. Thomas, T.C.E. (1951) Biting Activity of *Anopheles gambiae*. *BMJ* 2, 1402-1402.
- 480 45. Port, G.R. *et al.* (1980) The relationship of host size to feeding by mosquitoes of the
481 *Anopheles gambiae* Giles complex (Diptera: Culicidae). *Bulletin of Entomological Research* 70,
482 133-144.
- 483 46. Clyde, D.F. and Shute, G.T. (1958) Selective Feeding Habits of Anophelines Amongst Africans
484 of Different Ages. *Am J Trop Med Hyg* 7, 543-545.
- 485 47. Knols, B.G.J. *et al.* (1995) Differential attractiveness of isolated humans to mosquitoes in
486 Tanzania. *Trans R Soc Trop Med Hyg* 89, 604-606.
- 487 48. Mukabana, W. *et al.* (2002) Host-specific cues cause differential attractiveness of Kenyan
488 men to the African malaria vector *Anopheles gambiae*. *Malar J.* 1, 17.
- 489 49. Qiu, Y.T. *et al.* (2006) Interindividual variation in the attractiveness of human odours to the
490 malaria mosquito *Anopheles gambiae* s. s. *Med. Vet. Entomol.* 20, 280-287.
- 491 50. Ansell, J. *et al.* (2002) Short-range attractiveness of pregnant women to *Anopheles gambiae*
492 mosquitoes. *Trans R Soc Trop Med Hyg* 96, 113-116.
- 493 51. Batista, E. *et al.* (2014) *Anopheles darlingi* (Diptera: Culicidae) displays increased
494 attractiveness to infected individuals with *Plasmodium vivax* gametocytes. *Parasites Vectors*
495 7, 251.
- 496 52. Lacroix, R. *et al.* (2005) Malaria infection increases attractiveness of humans to mosquitoes.
497 *PLoS biology* 3, e298.
- 498 53. De Moraes, C.M. *et al.* (2014) Malaria-induced changes in host odors enhance mosquito
499 attraction. *Proceedings of the National Academy of Sciences* 111, 11079-11084.
- 500 54. Matovu, F. *et al.* (2009) How equitable is bed net ownership and utilisation in Tanzania? A
501 practical application of the principles of horizontal and vertical equity. *Malar J.* 8, 109.
- 502 55. Bernard, J. *et al.* (2009) Equity and coverage of insecticide-treated bed nets in an area of
503 intense transmission of *Plasmodium falciparum* in Tanzania. *Malar J.* 8, 65.
- 504 56. Kulkarni, M.A. *et al.* (2010) Contribution of Integrated Campaign Distribution of Long-Lasting
505 Insecticidal Nets to Coverage of Target Groups and Total Populations in Malaria-Endemic
506 Areas in Madagascar. *Am J Trop Med Hyg* 82, 420-425.
- 507 57. Baume, C.A. and Marin, M.C. (2007) Intra-household Mosquito Net Use in Ethiopia, Ghana,
508 Mali, Nigeria, Senegal, and Zambia: Are Nets Being Used? Who in the Household Uses Them?
509 *Am J Trop Med Hyg* 77, 963-971.
- 510 58. Quiñones, M.L. *et al.* (2000) Diversion of *Anopheles gambiae* from children to other hosts
511 following exposure to permethrin-treated bednets. *Med. Vet. Entomol.* 14, 369-375.

- 512 59. Reddy, M. *et al.* (2011) Outdoor host seeking behaviour of *Anopheles gambiae* mosquitoes
513 following initiation of malaria vector control on Bioko Island, Equatorial Guinea. *Malar J.* 10,
514 184.
- 515 60. Braimah, N. *et al.* (2005) Tests of bednet traps (Mbita traps) for monitoring mosquito
516 populations and time of biting in Tanzania and possible impact of prolonged insecticide
517 treated net use. *Int. J. Trop. Insect Sci.* 25, 208-213.
- 518 61. Moiroux, N. *et al.* (2012) Changes in *Anopheles funestus* Biting Behavior Following Universal
519 Coverage of Long-Lasting Insecticidal Nets in Benin. *J. Infect. Dis.* 206, 1622-1629.
- 520 62. McCann, R.S. *et al.* (2014) Reemergence of *Anopheles funestus* as a Vector of *Plasmodium*
521 *falciparum* in Western Kenya after Long-Term Implementation of Insecticide-Treated Bed
522 Nets. *Am J Trop Med Hyg* 90, 597-604.
- 523 63. Russell, T. *et al.* (2010) Impact of promoting longer-lasting insecticide treatment of bednets
524 upon malaria transmission in a rural Tanzanian setting with pre-existing high coverage of
525 untreated nets. *Malar J* 9, 187.
- 526 64. Geissbuhler, Y. *et al.* (2007) Interdependence of domestic malaria prevention measures and
527 mosquito-human interactions in urban Dar es Salaam, Tanzania. *Malar J.* 6, 126.
- 528 65. Sinka, M. *et al.* (2012) A global map of dominant malaria vectors. *Parasites Vectors* 5, 69.
- 529 66. Takken, W. and Verhulst, N.O. (2013) Host Preferences of Blood-Feeding Mosquitoes. *Annu.*
530 *Rev. Entomol.* 58, 433-453.
- 531 67. Midega, J.T. *et al.* (2007) Estimating dispersal and survival of *Anopheles gambiae* and
532 *Anopheles funestus* along the Kenyan coast by using mark-release-recapture methods. *J.*
533 *Med. Entomol.* 44, 923-929.
- 534 68. Shililu *et al.* (1998) Seasonal density, sporozoite rates and entomological inoculation rates of
535 *Anopheles gambiae* and *Anopheles funestus* in a high-altitude sugarcane growing zone in
536 western kenya. *Tropical Medicine & International Health* 3, 706-710.
- 537 69. Gneme, A. *et al.* (2013) Equivalent susceptibility of *Anopheles gambiae* M and S molecular
538 forms and *Anopheles arabiensis* to *Plasmodium falciparum* infection in Burkina Faso. *Malar J.*
539 12, 204.
- 540 70. Ndiath, M. *et al.* (2011) Comparative susceptibility to *Plasmodium falciparum* of the
541 molecular forms M and S of *Anopheles gambiae* and *Anopheles arabiensis*. *Malar J.* 10, 269.
- 542 71. Stevenson, J. *et al.* (2012) Novel vectors of malaria parasites in the western highlands of
543 Kenya. *Emerg. Infect. Dis.* 18, 1547-1549.
- 544 72. Riehle, M.M. *et al.* (2011) A Cryptic Subgroup of *Anopheles gambiae* Is Highly Susceptible to
545 Human Malaria Parasites. *Science* 331, 596-598.
- 546 73. Coetzee, M. *et al.* (2013) *Anopheles coluzzii* and *Anopheles amharicus*, new members of the
547 *Anopheles gambiae* complex. *Zootaxa* 3619, 246-274.
- 548 74. Schneider, P. *et al.* (2007) Submicroscopic *Plasmodium falciparum* gametocyte densities
549 frequently result in mosquito infection. *Am J Trop Med Hyg* 76, 470-474.
- 550 75. Muirhead-Thomson, R.C. (1954) Factors determining the true reservoir of infection of
551 *Plasmodium falciparum* and *Wuchereria bancrofti* in a West African village. *Trans R Soc Trop*
552 *Med Hyg* 48, 208-225.
- 553 76. Haji, H. *et al.* (1996) Absence of relationships between selected human factors and natural
554 infectivity of *Plasmodium falciparum* to mosquitoes in an area of high transmission.
555 *Parasitology* 113, 425-431.
- 556 77. Boudin, C. *et al.* (1993) High Human Malarial Infectivity to Laboratory-Bred *Anopheles*
557 *gambiae* in a Village in Burkina Faso. *Am J Trop Med Hyg* 48, 700-706.
- 558 78. Bonnet, S. *et al.* (2000) Comparison of artificial membrane feeding with direct skin feeding to
559 estimate infectiousness of *Plasmodium falciparum* gametocyte carriers to mosquitoes. *Trans*
560 *R Soc Trop Med Hyg* 94, 103-106.
- 561 79. Gamage-Mendis, A.C. *et al.* (1991) Infectious reservoir of *Plasmodium vivax* and *Plasmodium*
562 *falciparum* malaria in an endemic region of Sri Lanka. *Am J Trop Med Hyg* 45, 479-487.

80. Karl, S. *et al.* (2011) A Sub-Microscopic Gametocyte Reservoir Can Sustain Malaria Transmission. *PLoS ONE* 6, e20805.
81. Ouédraogo, A.L. *et al.* (2009) Substantial contribution of submicroscopical *Plasmodium falciparum* gametocyte carriage to the infectious reservoir in an area of seasonal transmission. *PLoS ONE* 4, e8410.
82. Bousema, T. *et al.* (2013) Can field-based mosquito feeding assays be used for evaluating transmission-blocking interventions? *Trends Parasitol.* 29, 53-59.
83. Solarte, Y. *et al.* (2007) Effects of Anticoagulants on *Plasmodium vivax* Oocyst Development in *Anopheles albimanus* Mosquitoes. *Am J Trop Med Hyg* 77, 242-245.
84. Stone, W.J.R. *et al.* (2013) The relevance and applicability of oocyst prevalence as a read-out for mosquito feeding assays. *Sci Rep* 3.
85. Smith, D.L. *et al.* (2004) The Risk of a Mosquito-Borne Infection in a Heterogeneous Environment. *PLoS Biol* 2, e368.
86. Liebman, K.A. *et al.* (2014) Determinants of Heterogeneous Blood Feeding Patterns by *Aedes aegypti* in Iquitos, Peru. *PLoS Negl Trop Dis* 8, e2702.
87. Ouédraogo, A.L. *et al.* (2007) Age-dependent distribution of *Plasmodium falciparum* gametocytes quantified by Pfs25 real-time QT-NASBA in a cross-sectional study in Burkina Faso. *Am J Trop Med Hyg* 76, 626-630.
88. Larson, P. *et al.* (2014) Insecticide-treated net use before and after mass distribution in a fishing community along Lake Victoria, Kenya: successes and unavoidable pitfalls. *Malar J.* 13, 466.

Table 1. Summary of studies assessing the malaria infectious reservoir where recruitment was conducted without regard for parasite status.^a

Study location	Local transmission setting	Feeding method	Assay endpoint ^b	Species	Prevalence of infectiousness (%) ^c	Proportional contribution to reservoir (%) ^d				Mosquito infection probability ^e	Refs
						1 - 4	5 - 9	10 - 19	≥20		
Papua New Guinea	Moderate perennial	Direct & Membrane	Oocysts	<i>Anopheles farauti</i>	3.8	22.7	32.9	30.3	14.2	0.013	(19)
						1 - 4	5 - 14	≥15			
Liberia	Intense perennial	Direct	Oocysts	<i>Anopheles gambiae</i>	10.6	40.5	28.4	31.1		0.023	(16)
Kenya	Intense perennial	Direct	Oocysts & CSP ELISA	<i>Anopheles gambiae</i>	10.1	23.2	36.1	40.7		0.010	(17)
Burkina Faso	Intense seasonal	Membrane	Oocysts	<i>Anopheles gambiae</i>	48.0	ND ^f	39.7	60.3		0.055	(77)
Cameroon (Bondi)	Intense seasonal	Direct	Oocysts	<i>Anopheles gambiae</i>	7.4	28.8	49.2	21.9		0.024	(18)
Cameroon (Mengang)	Intense seasonal	Direct	Oocysts	<i>Anopheles gambiae</i>	8.2	34.7	33.9	31.4		0.011	(18)
Senegal	Low, unstable	Direct	Oocysts	<i>Anopheles arabiensis</i>	8.7	ND ^f	51.8	48.2		0.002	(32)

^aOnly studies that have utilised demographic and mosquito feeding data to characterise the human infectious reservoir of *Plasmodium* in endemic populations were included. All studies were conducted in areas where *Plasmodium falciparum* and non-*falciparum* species co-circulate, so the outcome of feeding assays cannot be attributed unambiguously to *falciparum* malaria. One additional study not included in the table utilised population and xenodiagnostic data to quantify the malaria infectious reservoir in Sri Lanka (79). This study recruited individuals with incident malaria over a 17 month period, the majority of which were *Plasmodium vivax*. The study design and analysis was suitably different as to exclude the possibility of direct comparison with the other studies presented in the table.

^b Assay endpoint was either 1) oocysts counted on the mosquito midgut by microscopy; or 2) oocysts counted by microscopy or positivity of remaining live mosquitoes at the end of the experiment (day 15 post infection) in the circumsporozoite protein (CSP) ELISA (17).

^c Prevalence of infectiousness was calculated as the combined prevalence of individuals infectious to ≥1 mosquito in each age category, adjusted for proportion of the population made up by the age groups in the total population.

^d Proportional contribution to the reservoir presents the proportional contributions of three or four age groups to population-adjusted reservoir, with age groupings combined where the data in the original publication allowed.

^e Mosquito infection probability corresponds to the probability that a mosquito biting a random individual in a population will develop infection. This measure was either presented as in the published data, or was calculated by combining demographic data, age-specific prevalence of infectiousness and age-specific mosquito infection rates during infective membrane feeding experiments.

^f ND: Studies failed to perform mosquito feeds on individuals <5 years of age. Data has been adjusted to represent the infectious reservoir of individuals >5 years only.

Table 2. Summary of studies assessing the malaria infectious reservoir that recruited only gametocyte positive individuals for feeding assays.^a

Study location	Local transmission setting	Feeding method	Assay endpoint ^b	Species	Prevalence of infectiousness (%) ^c	Proportional contribution to reservoir (%) ^d				Mosquito infection probability ^e	Refs
						1 - 4	5 - 9	10 - 19	≥20		
The Gambia	Intense seasonal	Membrane	Oocysts	<i>Anophelesgambiae</i>	5.6	17.5	21.7	22.2	37.9	0.015	(11)
Tanzania	Intense perennial	Membrane	Oocysts	<i>Anopheles gambiaes.l.</i>	3.2	28.4	23.1	14.2	34.0	0.007	(11)
Papua New Guinea ^f	Moderate perennial	Direct & Membrane	Oocysts	<i>Anopheles farauti</i>	3.6 (1.4)	48.0	12.4	24.0	15.5	0.012 (0.004)	(19)

^aOnly studies that have utilised demographic and mosquito feeding data to characterise the human infectious reservoir of *Plasmodium* in endemic populations were included. All studies were conducted in areas where *Plasmodium falciparum* and non-falciparum species co-circulate, so the outcome of feeding assays cannot be attributed unambiguously to falciparum malaria.

^bAssay endpoint was oocysts counted on the mosquito midgut by microscopy.

^cPrevalence of infectiousnesswas calculated as the combined prevalence of individuals infectious to ≥1 mosquito in each age category, adjusted for proportion of the population made up by the age groups in the total population.

^dProportional contribution to the reservoir presents the proportional contributions of three or four age groups to population-adjusted reservoir, with age groupings combined where the data in the original publication allowed.

^eMosquito infection probability corresponds to the probability that a mosquito biting a random individual in a population will develop infection. This measure was either presented as in the published data, or was calculated by combining demographic data, age-specific prevalence of infectiousness and .age-specific mosquito infection rates during infective membrane feeding experiments.

^fFor this study, all species and *P. falciparum*-specific (within parenthesis) estimates are presented.

Figure 1. Age contributions to the infectious reservoir.

A. Individuals in the figure are represented by circles in three age groups; <5, 5-15, and >15 years. The abundance of individuals in each group reflects a simplified population age-structure in sub-Saharan Africa, as described in Box 1 and in the studies detailed in Table 1 and 2 (<5=15%, 5-15=30%, >15=55%) (<http://esa.un.org/wpp/>). **B.** Speckling within circles represents the presence of *Plasmodium falciparum* gametocytes detectable using quantitative-nucleic acid sequence-based amplification (QT-NASBA); age-specific prevalence presented here were taken from an area of intense seasonal transmission(87). **C.** Solid filled circles represent who might infect mosquitoes in a mosquito feeding assay. Some (or most) gametocytaemic individuals may not be infectious for a variety of reasons, including gametocyte density, maturity, recent treatment, and transmission blocking immunity. Infectiousness prevalence for the three age groups (<5=25%, 5-15=13%, >15=7%) is based on surveys described in Table 1(16, 17, 18). Pie chart 1 demonstrates the proportional contribution of each age group to the total population infectious reservoir. Demographic adjustment is not necessary as this example directly represents the age-composition of a typical population. Of the theoretical total infectious population (9.8% [5/51 individuals]), individuals in the age groups <5, 5-15 and >15 years account for 40%, 40% and 20% of the reservoir respectively. **D.** Equally exposed individuals of different ages are differently attractive to mosquito biting, partly due to differences in body surface area. To reflect this, individual circles were given a surface area proportional to the average body surface area of individuals in the three age groups (<5=0.4m², 5-15=1.2m², >15=1.6m²; Ratio of surface area 1:3:4), using data from Port *et al.* showing that body surface area correlated positively with mosquito biting rate (45). Pie chart 2 illustrates the potential effect of increased mosquito sampling due to body surface area in each age group on the infectious reservoir. Finally, to account for some of the age related differences in exposure intensity, the percentage bed net use among recorded in individuals from Western Kenya (Coverage: <5=42%, 6-17=26%, >17=57.7%; Ratio of exposure 1:1.2:0.7) was used to further adjust the surface area weighted contributions to the infectious reservoir (Pie chart 3)(88).

Figure 2. Cross-sectional surveys and infectious reservoir dynamics.

Data represent hypothetical annual transmission measures in two different endemic settings of seasonal and perennial malaria transmission. The dashed line corresponds to the timing of cross-sectional xenodiagnostic surveys. The population plots underneath the main charts reflect hypothetical population level infectiousness, as would be measured in direct assessments of the

reservoir. Individuals in the plots are represented by circles in three age groups; <5, 5-15, and >15 years. The abundance of individuals in each group reflects a simplified population age-structure in sub-Saharan Africa, as described in Box 1 and in the studies detailed in Table 1 and 2 (<5=15%, 5-15=30%, >15=55%) (<http://esa.un.org/wpp/>). Speckling within circles represents the presence of *Plasmodium falciparum* gametocytes; age-specific prevalence is hypothetical. Solid filled circles represent who might infect mosquitoes in a mosquito feeding assay; infectiousness prevalence for the three age groups is hypothetical. To reflect differences in body size and potentially attractiveness to mosquitoes, individual circles were given a surface area proportional to the average body surface area of individuals in the three age groups (<5=0.4m², 5-15=1.2m², >15=1.6m²; Ratio of surface area 1:3:4), using data from Port *et al.* showing that body surface area correlated positively with mosquito biting rate (45). The combined surface area of individuals in the plots can be interpreted broadly as the contribution each group makes to the total infectious reservoir. Combined, the figure illustrates that the total infectious reservoir and the proportional contribution of differently aged individuals are likely to vary, potentially significantly, over time in response to the dynamicity of numerous factors. Though variations in transmission potential would probably be clearest between surveys where transmission is seasonal, age-specific contributions may vary with time in areas where transmission is intense and perennial.

Figure 1

A. Age-stratified population

B. Proportion of individuals with gametocytes

C. Proportion of individuals infective to mosquitoes

Proportional contribution to total infectious reservoir by age

D. Individuals adjusted for surface area/mosquito biting frequency

Figure 2