Surprising lack of liposome-induced complement activation by artificial 1,3-diamidophospholipids in vitro

Simon Bugna, MSc^a, Marzia Buscema^a, MSc, Sofiya Matviyiv^a, MSc,
Rudolf Urbanics, M.D., PhD^b, Andreas Weinberger, PhD^c, Tamas Meszaros^b,
Janos Szebeni, M.D., PhD, DSc^b, Andreas Zumbuehl, PhD^{*c}, Till Saxer, M.D.^{*d},
Bert Müller, PhD^{*a}

Abstract

Cardio-vascular diseases are the main cause of death, emphasizing the need to improve patient treatment and survival. One therapeutic approach is a liposome-based drug carrier system specifically targeting constricted arteries. The recently discovered mechano-sensitive liposomes use hemodynamic shear-stress differences between healthy and constricted blood vessels as trigger for drug release. Liposomes are promising delivery containers but being recognized as foreign by the immune system. Complement activation as essential factor of the recognition leads to adverse effects. Here, we tested complement activation by liposomes formulated from the artificial phospholipid Pad-PC-Pad *in vitro*. Surprisingly no complement activation was detected in human sera and porcine plasma. In *in vivo* experiments with three pigs, neither anaphylactic reactions nor other significant haemodynamic changes were observed even at comparably high liposome doses. The pilot study holds promise for an absence of complement-mediated adverse effects of Pad-PC-Pad liposomes in human.

Key words: Pad-PC-Pad; CARPA; pig study; shear sensitive nano-container; physical trigger

According to the World Health Organisation, cardiovascular diseases are responsible for 30 % of worldwide deaths.¹ Many of them arise during transfer to hospital, motivating the increasing interest in pre-hospital drug delivery to critically constricted vessels.^{2, 3} One strategy involves targeting the constricted arteries by capitalizing on the endogenous, elevated shear stresses present in diseased artery segments and use the changes as a purely physical trigger for drug release.⁴

Liposomes composed of the artificial phospholipid Pad-PC-Pad can release their payload in areas of increased shear stress.⁵ In order to push this development towards clinical application, the liposomes need to be further characterized to exclude non-desired systemic effects. Here, predicting the risk of complement-activation-related-pseudo-allergy (CARPA) is fundamental, as it characterizes the hypersensitivity towards liposomal formulations.^{6, 7} This phenomenon also occurs with clinically approved lipid-based drugs and creates adverse effects in 2 to 30 % of people, occasionally leading to anaphylactic reaction or even death.^{7, 8}

Previous studies have shown that liposomal formulations including Doxil^{9, 10} and AmBisome¹¹ lead to lethal effects, if 0.5 to 5.0 mg/kg phospholipids are applied. First cell experiments with Pad-PC-Pad, however, have indicated a reasonable tolerance.⁵ Therefore, we propose *in vitro* tests with human sera and porcine plasma to characterize the CARPA reaction and animal studies to search for liposome-induced blood-pressure changes, both are fundamental to bridge the gap between bench and bedside. We hypothesize that contrary to Doxil and AmBisome the liposomes from the artificial phospholipid Pad-PC-Pad without an encapsulated active drug are reaction-free at comparable dose. Revealing such lack of complement activation, the native drug carrier could be assumed as safe from CARPA.

Methods

Pad-PC-Pad (1,3-palmitoylamido-1,3-deoxy-*sn*-glycero-2-phosphatidyl-choline) has been prepared according to the established protocol.⁵ Figure 1 displays its structure in comparison to the natural 1,2-diesterphospholipid DPPC.

The liposomes were formulated using the thin-film method.¹² The phospholipids hydrated with PBS were extruded at a temperature between 65 and 70 °C through 100 nm track-edged filters at a concentration of 20 mg/mL. The liposome stock suspension was stored without light at a temperature of 4 °C. The experiments were conducted within 48 hours after formulation. Liposome aggregation and sedimentation were observed after 12 hours storage. Therefore, before bolus injection, the storage vial was repeatedly inverted to re-suspend the sediments, and, subsequently, the suspension was passed through a 22- μ m syringe filter.

Five human sera samples, labelled H1 to H5, were tested with the Quidel MicroVue SC5b-9 ELISA kit. A volume of 20 μ L Pad-PC-Pad suspension was added to each serum at a temperature of 37.1 °C. Additionally, five pig hirudinized plasma samples, labelled P1 through P5 were tested with the Quidel MicroVue Pan-specific C3 reagent kit. A volume of 25 μ L of Pad-PC-Pad suspension was added to each plasma. After 0, 20, 40, and 60 minutes, EDTA was added to stop the reaction. Each series contained controls of PBS only and 20 or 25 μ L zymosan (1.2 mg/mL), purchased from Sigma Chemical Co. (St Louis, MO). The optical density of the samples was measured with a plate reader at a wavelength of 450 nm. The concentration of the SC5b-9 complex was calculated from a linear calibration curve.

Three female 12- to 14-week-old Yorkshire pigs with a weight of 18.4, 22.8 and 20.2 kg were injected with various bolus doses of Pad-PC-Pad vesicle suspensions and with zymosan, while monitoring the haemodynamic parameters. The protocol was approved by the local Animal Subject Review Committees, details see appendix C of supplementary information.

Results and discussion

Figure 2 summarizes the results of the Pad-PC-Pad liposome tests using human sera in the ELISA assay for SC5b-9. Within the error bars, the liposomes do not induce any complement activation, whereas the complement activator zymosan triggered an immediate complement reaction more than 20 times higher than the negative control. The variability between the five donors is given in Figure S2. The results are in line with placebo Doxil (Doxebo).¹³ The soluble terminal complement complex SC5b-9 is formed from C5 to C9 as the end result of complement activation.¹⁴ An elevation of SC5b-9 against artificial phospholipid liposomes is therefore indicative for the *in vivo* reaction. Consequently, one does not expect *in vivo* reactions in pig experiments.

Indeed, Figure 3 demonstrates that the *in vivo* reactions have been consistent with the ELISA tests. Globally, the three pigs showed a stable hemodynamic behaviour during anaesthesia and after injection of Pad-PC-Pad. Since SAP, heart rate, and ECG were constant within the error bars, Figure 3 only displays PAP. Two pigs, Figure 3A and 3C, were injected with unfiltered, aggregated Pad-PC-Pad liposomal suspensions. The third pig, Figure 3B, was injected with a filtered, non-aggregated Pad-PC-Pad suspension. These injection regimes indicate the differences between homogeneous, equally sized liposomes and liposomal aggregates. The first pig (Figure 3A1) showed no reaction after the 0.5 mg/kg injection. Even after the 5 mg/kg injection the PAP remains constant. Three minutes after an additional 5 mg/kg injection (Figure 3A3) the PAP increased by 500 Pa (4 mmHg) for two minutes. The positive reference zymosan injection led to an immediate increase by more than 2.7 kPa (20 mmHg), long lasting reaction (Figure 3A4). The second pig showed no hemodynamic reaction applying doses of 0.5 and 5 mg/kg (Figure 3B2 and B3, respectively). In the third pig (Figure 3C), the PAP was stable after the first aggregated liposome injection with a dose of 5 mg/kg, but increased by 930 Pa (7 mmHg) for a period of six minutes subsequent to the

second injection. As the third pig was reactive to a certain extent, we decided to directly proceed with histological sampling and skip the zymosan injection. This histology analysis revealed that the administration of Pad-PC-Pad liposomes did not lead to histological toxic modification in kidney, lung, heart, and liver. Table 1 summarizes the analysis of the porcine blood. The values correspond to the ones reported for healthy Yorkshire pigs.¹⁵ The lack of thromboxane changes in pigs is consistent with the *in vitro* test. C5a stimulates the production of arachidonic acid derivatives prostaglandins and thromboxane by neutrophils and macrophages. The thromboxane levels remain stable during the entire experiment. They only increased following the zymosan injection.

To support the link between the *in vitro* and *in vivo* experiments complement activation was also measured via C3 consumption in pig plasma. The data, presented in Figure 4, reveal the lack of complement activation.

Comparing the results of the *in vivo* study with data published on Doxil, and AmBisome we observed the lack of reactivity for Pad-PC-Pad liposomes. Even at one order of magnitude higher concentrations the non-aggregated Pad-PC-Pad liposomes do not trigger any reactivity *in vivo*.

In conclusion, this pilot study was performed to explore the *in vitro* complement activation by Pad-PC-Pad and the *in vivo* CARPA reactivity in a pig model, which is more sensitive for CARPA than humans. The clinical relevance of the study lies in the increased risk for non-IGE-mediated reactions in patients treated for the first time with liposomal drugs. Although premedication reduces the risk of anaphylactic reactions, the risk for such reactions remains present. The findings indicate that Pad-PC-Pad liposomes do not directly or indirectly cause anaphylactic reactions even with a dose as high as 110 mg/kg. In comparison to approved phospholipid formulations,^{7, 16} Pad-PC-Pad with a ten- or even hundred-fold higher

dose reveals much less CARPA reactivity *in vivo*. To this end, Pad-PC-Pad is regarded as a promising phospholipid for liposomal drug delivery applications in humans.

Acknowledgments

We acknowledge the experimental support of Dennis Müller as well as the help of Hans Deyhle and Georg Schulz in preparing Figure 3.

References

1. Go, A. S.; Mozaffarian, D.; Roger, V. L.; Benjamin, E. J.; Berry, J. D.; Borden, W. B., et al. Heart Disease and Stroke Statistics—2013 Update A Report From the American Heart Association, *Circulation*. 2013;**127**:e6-e245.
2. Crielgaard, B. J.; Lammers, T.; Schiffelers, R. M.; Storm, G. Drug targeting systems for inflammatory disease: One for all, all for one, *Journal of Controlled Release*. 2012;**161**:225-234.
3. Ruiz-Esparza, G. U.; Flores-Arredondo, J. H.; Segura-Ibarra, V.; Torre-Amione, G.; Ferrari, M.; Blanco, E., et al. The physiology of cardiovascular disease and innovative liposomal platforms for therapy, *International Journal of Nanomedicine*. 2013;**8**:629-640.
4. Saxer, T.; Zumbuehl, A.; Müller, B. The use of shear stress for targeted drug delivery, *Cardiovascular Research*. 2013;**99**:328-333.
5. Holme, M. N.; Fedotenko, I. A.; Abegg, D.; Althaus, J.; Babel, L.; Favarger, F., et al. Shear-stress sensitive lenticular vesicles for targeted drug delivery, *Nature Nanotechnology*. 2012;**7**:536-543.
6. Nath, P.; Basher, A.; Harada, M.; Sarkar, S.; Selim, S.; Maude, R. J., et al. Immediate hypersensitivity reaction following liposomal amphotericin-B (AmBisome) infusion, *Tropical Doctor*. 2014;**44**:241-242.
7. Szebeni, J. Complement activation-related pseudoallergy: a new class of drug-induced acute immune toxicity, *Toxicology*. 2005;**216**:106-121.
8. Chanan-Khan, A.; Szebeni, J.; Savay, S.; Liebes, L.; Rafique, N. M.; Alving, C. R., et al. Complement activation following first exposure to pegylated liposomal doxorubicin (Doxil®): possible role in hypersensitivity reactions, *Annals of Oncology*. 2003;**14**:1430-1437.
9. Lyass, O.; Uziely, B.; Ben-Yosef, R.; Tzemach, D.; Heshing, N. I.; Lotem, M., et al. Correlation of toxicity with pharmacokinetics of pegylated liposomal doxorubicin (Doxil) in metastatic breast carcinoma, *Cancer*. 2000;**89**:1037-1047.
10. Northfelt, D. W.; J., D. B.; Thommes, J. A.; Miller, B. J.; Fischl, M. A.; Friedman-Kien, A., et al. Pegylated-liposomal doxorubicin versus doxorubicin, bleomycin, and vincristine in the treatment of AIDS-related Kaposi's sarcoma: results of a randomized phase III clinical trial, *Journal of Clinical Oncology*. 1998;**16**:2445-2451.
11. Barratt, G.; Bretagne, S. Optimizing efficacy of Amphotericin B through nanomodification, *International Journal of Nanomedicine*. 2007;**2**:301-313.
12. Olson, F.; Hunt, C. A.; Szoka, F. C.; Vail, W. J.; Papahadjopoulos, D. Preparation of liposomes of defined size distribution by extrusion through polycarbonate membranes, *Biochimica et Biophysica Acta (BBA) - Biomembranes*. 1979;**557**:9-23.

13. Szebeni, J.; Bedőcs, P.; Urbanics, R.; Bünger, R.; Rosivall, L.; Tóth, M., et al. Prevention of infusion reactions to PEGylated liposomal doxorubicin via tachyphylaxis induction by placebo vesicles: A porcine model, *Journal of Controlled Release*. 2012;**160**:382-387.
14. Müller-Eberhard, H. J. *The Membrane Attack Complex*, 1984.
15. Cooper, C. A.; Moraes, L. E.; Murray, J. D.; Owens, S. D. Hematologic and biochemical reference intervals for specific pathogen free 6-week-old Hampshire-Yorkshire crossbred pigs, *Journal of Animal Science and Biotechnology*. 2014;**5**:5.
16. Szebeni, J.; Fontana, J. L.; Wassef, N. M.; Mongan, P. D.; Morse, D. S.; Dobbins, D. E., et al. Hemodynamic Changes Induced by Liposomes and Liposome-Encapsulated Hemoglobin in Pigs A Model for Pseudoallergic Cardiopulmonary Reactions to Liposomes: Role of Complement and Inhibition by Soluble CR1 and Anti-C5a Antibody, *Circulation*. 1999;**99**:2302-2309.

Figure Captions

Figure 1. Structures from the natural 1,2-diesterphospholipid DPPC, the major component of FDA-approved liposomal formulations and below of 1,3-diamidophospholipid Pad-PC-Pad that can form the lentil-shaped liposomes used in this study. The scheme explains the surgery procedure. The details are described in the supplementary material, appendix C.

Figure 2. The complement activation of Pad-PC-Pad liposomes with five human sera labelled H1 to H5 is comparable to the negative control PBS for the incubation times indicated. The bars are the standard deviations between the donors. Significance of differences among the groups was determined by Two-Way analysis of Variance (ANOVA), followed by the Newmann-Keuls test ($P < 0.05$).

Figure 3. PAP change is normalized to time zero. (A) Subsequent injections of Pad-PC-Pad liposomes at 0.5 and 5 mg/kg, followed by an injection of 0.1 mg/kg zymosan as positive control. The sample was re-suspended and filtered before the first injection. (B) Injection of directly filtered Pad-PC-Pad liposomes to avoid liposome aggregate formation. (C) Injection of non-filtered liposomes at 5 mg/kg lipid concentration.

Figure 4. Relative C3 consumption, here Pad-PC-Pad liposomes with the pig plasma labelled P1 to P5, is a standard to *in vitro* quantify complement activation in animal samples, *cf.*

Figure S3. The statistical analysis, as described for Figure 2, demonstrates that the complement activation of the Pad-PC-Pad liposomes corresponds to the negative control.

Table 1

In vivo haematology analysis of the immune reaction of the pigs towards the artificial liposomal formulations

	Baseline			Post injection follow up (6H)			Reference values ^{24,25*}
	Pig 1	Pig 2	Pig 3	Pig 1	Pig 2	Pig 3	
Glucose [mmol/L]		2.8	5.3		3.7	4.7	3.6–8.3
Creatine kinase [IU/L]	1297.0	1156.5	2107.5	1225.0	1156.5	1470.5	153–5427
GOT [IU/L]		40.5	29.0		29.0	23.5	32–84
GPT [mg/dL]		32.0	26.5		24.5	20.5	13–111
LDH [U/L]		672.5	815.0		1005.5	718.5	380–634
Bilirubin total [mmol/dL]		1.4	1.7		2.3	1.2	0.3–8.2
Na ⁺ [mmol/L]	143.0	140.5	127.5	143.0	135.5	130.0	131–150
K ⁺ [mmol/L]	4.5	4.8	5.0	5.1	5.5	5.4	3.7–7.1
Cl ⁻ [mmol/L]	100.0	101.5	91.0	99.0	98.5	96.0	93–108
Urea [mmol/L]		4.5	5.5		7.4	6.5	3.6–10.7
Creatinine [mmol/dL]		81.0	84.5		78.5	71.0	70–239
Troponin T5 [ng/dL]	14	6	5	9	5	4	< 500
CKMB [nmol/dL]	< 0.3	< 0.3	< 0.3	< 0.3	< 0.3	< 0.3	< 50
PT [s]		11.9	12.5		12.5	11.5	
APTI [s]		48.3	27.2		42.9	> 240.0	
D-dimer [µg/L]		0.02	0.0		0.83	0.18	< 500**
Hemoglobin [g/L]	94.0	97.5	108.0	94.5	105.5	103.0	88–127
Red blood cells [10 ¹² /L]	6.7	6.7	7.9	6.8	7.3	7.3	5.5–9.1
Platelet [10 ⁹ /L]	295.5	361.0	425.5	288.0	394.5	317.0	207–873
White blood cells [10 ⁹ /L]	11.1	22.5	23.2	9.4	21.2	22.2	5.44–25.1
Granulocyte [10 ⁹ /L]	6.7	16.4	16.7	5.2	18.5	17.8	1.1–13.3
Granulocyte [%]	60.2	73.0	72.0	55.2	87.3	80.0	40–75
Leucocyte [10 ⁹ /L]	4.1	5.3	5.9	3.5	2.4	3.3	3.8–15.0
Leucocyte [%]	37.1	23.6	25.3	37.5	11.5	15.1	25–45

*Pig Yorkshire reference values. **Human reference values because of lack of pig data.

GOT: glutamic oxaloacetic transaminase. GPT: glutamic-pyruvate transaminase. LDH: lactate dehydrogenase. CKMB: creatine kinase-myocardial band. PT: prothrombin time. APTI: activated partial thromboplastin time.

Text of graphical abstract

Atherosclerosis is an inflammatory disease associated with the formation of plaque-induced constrictions. Preferential vasodilator delivery can be achieved via shear-sensitive nanocontainers such as lentil-shaped liposomes from Pad-PC-Pad. Liposomal formulations, including the FDA-approved Doxil and AmBisome, are being recognized as foreign bodies and therefore cause complement activation. This communication comprises *in vitro* and *in vivo* experiments that show a surprisingly low or even absent complement-activation-related-pseudo-allergy (CARPA) for the lentil-shaped, artificial liposomes.

Figure 1

Figure 2

Figure 3

Figure 4