Ichthyotoxic *Karlodinium veneficum* (Ballantine) J Larsen in the Upper Swan River Estuary
(Western Australia): Ecological conditions leading to a fish kill

Jason E. Adolf^{a,c,1}, Tsvetan R. Bachvaroff^a, Jonathan R. Deeds^b, and Allen R. Place^a

(a)University of Maryland Center for Environmental Science, Institute of Marine and
Environmental Technology, 701 E. Pratt St., Baltimore, Maryland, USA 21202;

(b) US FDA Washington Seafood Laboratory, 8301 Muirkirk Rd., Laurel, Maryland, USA 20708

(c) Present Address: University of Hawaii Hilo, Dept. of Marine Science, 200 W. Kawili St., Hilo,
Hawaii, USA 96720

(1) Corresponding author, University of Hawaii Hilo, Dept. of Marine Science, 200 W. Kawili St.,
Hilo, Hawaii, USA 96720 phone: 808-932-7592

Email: JEA (jadolf@hawaii.edu), TRB (bachvarofft@gmail.com), JRD
(jonathan.deeds@fda.hhs.gov), ARP (place@umces.edu)

Running title: *Karlodinium* in an Australian estuary

37

38 ACKNOWLEDGEMENTS

39 The authors thank all the staff and crew of the WA DOE and SRT, particularly Ashrafi Begum,
40 Wasele (Vas) Hosja, Tarren Reitsema, Petra Ringeltaube, Malcolm Robb, and Laura Cunningham
41 for speedy delivery of phytoplankton count data. We thank Nancy Feissner who worked at
42 University of Maryland COMB as a summer technician and processed the >250 toxin samples
43 received from Australia in that time period. Parts of this work have been supported by NOAA
44 ECOHAB, CDC, and a NOAA CSCOR travel award. This is contribution #xxx from the University
45 of Maryland Center for Environmental Sciences and contribution # xxx from the ECOHAB
46 program.

47

ABSTRACT

Ichthyotoxic *Karlodinium veneficum* has become a persistent problem in the eutrophic Swan River Estuary (SRE) near Perth, Western Australia. Karlotoxin (KmTx) concentrations and *K. veneficum* were sampled from March to July 2005, spanning a bloom confirmed by microscopy and genetics (ITS sequence), and a fish kill coincident with end of the bloom. The objective of this study was to investigate *K. veneficum* cell and toxin dynamics, and water quality conditions, leading up to the bloom and fish kill in this estuarine system. Abundance of *K. veneficum* increased as diatom abundance decreased over a 3-month period (Jan – Mar) preceding the bloom. Low freshwater flow to the SRE characterized the bloom initiation period, while elevated seasonal flows altered water quality and preceded the end of the bloom and fish kill. The bloom of *K. veneficum* was localized over a bottom layer of hypoxic water in a stratified water column. Low nitrate levels, DIN:DIP (mol) near unity, and particulate C:N:P of *K. veneficum*-rich water samples were consistent with nitrogen limitation of phytoplankton. A KmTx 2 congener was present in the concentration range 0 – 1052 ng KmTx mL⁻¹, levels that were sufficient to kill larval fish in the laboratory within 4 hours. A KmTx cell quota of 2.8 pg KmTx cell⁻¹ was estimated for the bloom, which is moderately high for the species. Gill histopathology of fish from this fish kill showed signs of damage similar to those caused by KmTx in the lab. Results from this study suggest that conditions in the SRE, including elevated *K. veneficum* abundance and KmTx cell quotas, as well as hypoxia in the upper SRE, likely contribute to seasonal fish kills observed in this system.

Keywords: fish kill, *Karlodinium veneficum*, karlotoxins, Swan River Estuary

INTRODUCTION

The Swan River Estuary (SRE) is a subtropical microtidal estuary with a Mediterranean climate and has been negatively impacted by anthropogenic nutrient loading (Thompson and Hosja 1996) and hydrological alterations (Chan et al. 2002). The seasonal cycle of phytoplankton dynamics in the SRE is largely driven by nitrate supply and riverine runoff (Thompson 1998, Chan and Hamilton 2001, Horner-Rosser and Thompson 2001). Dinoflagellate blooms (*Gymnodinium* spp., *Prorocentrum*) have been observed austral summer through autumn in stratified portions of the upper estuary where surface nitrate concentrations are low and bottom-water ammonium is high (Hamilton et al. 1999, Twomey and John 2001), but no toxicity was reported associated with these blooms. Recently, the seasonal cycle of phytoplankton dynamics has been punctuated by harmful algal blooms (HAB). A major toxic bloom of *Microcystis aeruginosa* (cyanobacterium) in 2000 was associated with an anomalous, record-breaking summer rainfall event that freshened much of the SRE and dispersed the bloom population from the upper estuary into the lower basin (Hamilton 2000, Robson and Hamilton 2003). Ichthyotoxic blooms of the dinoflagellate, *Karlodinium veneficum*, had not been reported in the SRE before 2002, but have occurred periodically since that time.

The dinoflagellate *K. veneficum* (10–15 µm) is a cosmopolitan mixotrophic (Li et al. 1996, Adolf et al. 2006a, Calbet et al. 2011) protist responsible for numerous fish kills in temperate estuaries on the eastern seaboard of the United States (Deeds et al. 2002, Kempton et al. 2002, Fensin 2004, Goshorn et al. 2004, Garces et al. 2006, Hall et al. 2008) as well as in other waters around the world (Bachvaroff et al. 2008, Place et al. 2012). Previously *K. veneficum* has been called *Gyrodinium estuariale*, *Gymnodinium galatheanum*, *Gymnodinium veneficum*, and *K. micrum*, all of which are now synonymous with *K. veneficum* (Bergholtz et al. 2005). Ichthyotoxic

95 and cytotoxic bioactivity associated with *K. veneficum* was first reported by Abbot and Ballantine
 96 (1957), as *Gymnodinium veneficum* (Ballantine 1956).

97 The linear polyketide toxins produced by *K. veneficum* (Van Wagoner et al. 2008, 2010,
 98 Peng et al. 2010), karlotoxins (KmTx), have hemolytic, cytotoxic and ichthyotoxic activities (Deeds
 99 et al. 2002, Kempton et al. 2002, Mooney et al. 2009). Strains of *Karlodinium veneficum* from
 100 geographically-distinct regions produce different KmTx congeners that vary in chromatographic
 101 properties, UV absorption maxima (Deeds et al. 2004) and mass spectra (Deeds et al. 2004,
 102 Bachvaroff et al. 2008) although they are similar in bioactivity. The bioactivity of karlotoxin is
 103 membrane pore formation, dependent on the sterol composition of the target cell. The predominance
 104 of 4- α -methyl sterols (i.e. gymnodinosterol and brevestanol) in *K. veneficum* makes it immune to its
 105 own toxin, whereas organisms (i.e. fish, other protists) with a predominance of 4-*des*-methyl sterols,
 106 (i.e. cholesterol), are susceptible to KmTx bioactivity (Deeds and Place 2006, Adolf et al. 2006b).
 107 Karlotoxins have anti-grazing / allelopathic activity (Adolf et al. 2007, 2008; Stoecker et al. 2008,
 108 Brownlee et al. 2008, Waggett et al. 2008, Vaquer et al. 2005). Ichthyotoxicity of KmTx in situ
 109 appears to depend on cell lysis since intact, toxic *K. veneficum* has little effect on fish (Deeds et al.
 110 2002, Mooney et al. 2010, Dorantes-Aranda et al. 2011). In fish exposed to free KmTx, lysis of gill
 111 epithelial cells leading to collapse of the gill lamellae is observed (Deeds et al. 2006), presumably
 112 impairing gas and ion exchange across the gills.

113 Records predating 2002 for the SRE indicate observations of *Gyrodinium estuariale* (April
 114 1999 @ 32,926 cells per mL) and *Gymnodinium galatheanum* (March 2001 @ 6,000 cells per mL)
 115 (Jeff Cosgrove, pers. comm.), which were likely *K. veneficum* (Bergholtz et al. 2005) but no
 116 ichthyotoxicity was reported. From April-July 2003, numerous fish kills occurred in both the Swan
 117 and Canning River estuaries located in Perth, Western Australia. These kills coincided with a
 118 persistent bloom of *K. veneficum* starting in the Swan River in early April and spreading to the

Canning River in Mid-June. The bloom ended in early July following a large rainfall event. The bloom was preceded by unusually high rainfall in the area during early April followed by a prolonged period of calm, sunny weather. *Karlodinium veneficum* densities fluctuated during this period, often reaching 1.0×10^5 cells ml^{-1} . Numerous fish species were affected, including black bream (*Acanthopagrus butcheri*), a popular local sport fish, Perth herring (*Elops machnata*), and Swan River gobies (*Pseudogobius olorum*). Most kills contained $< 1,000$ fish, but two kills were estimated to contain in excess of 100,000 fish each (P. Musk, pers. comm.). Observations of fish gill histopathology were consistent with effects observed during exposure of zebrafish (*Danio rerio*) to purified KmTx (Deeds et al. 2006).

Toxic blooms of *K. veneficum* were first observed in the SRE in 2002 but detailed measurements of karlotoxin and *K. veneficum* abundance in the SRE have not been published before. Cell concentrations of *K. veneficum*, karlotoxin levels and SRE water quality were sampled to better understand the role of *K. veneficum* in fish mortalities that were anticipated to occur between April and June (austral autumn) based on past years' observations. It was hypothesized that increased karlotoxin concentrations, either through increased cell numbers or increases in cellular toxin cell quotas ($\text{pg toxin cell}^{-1}$) will precede or accompany major fish kills. Karlotoxin (KmTx) concentration associated with the observed 2005 SRE fish kill was closely examined and compared values with levels of KmTx observed to kill fish in laboratory trials.

MATERIALS AND METHODS

Water sampling

Swan River Estuary (Fig. 1) water quality sampling and analysis was done weekly, in the morning, according to standard procedures of the Western Australia Dept. of Environment (WA DoE) and Swan River Trust (<http://www.swanrivertrust.wa.gov.au/docs/monitoring-and-evaluation/water-quality->

[parameters.pdf](#)), including all salinity, temperature, nutrients, and oxygen measurements (Hydrolab or YSI multiparameter sonde, 0.5 m depth intervals). Nutrient samples were forwarded to Queensland Health Scientific Services (<http://www.health.qld.gov.au/qhcss/qhss/>) for analysis on the day of collection, if not the following day with overnight storage at 4 °C. Phytoplankton cell count samples were collected from the same stations as nutrient samples and counts conducted weekly by WA DoE standard procedures on surface-collected water samples. River discharge data for the Avon (Station 616011, 116.068524205 E 31.750532184 S) and Bennet Brook (Station 616084, 115.959525641 E 31.877702539 S) were obtained from the State of Western Australia Department of Water.

ITS sequence analysis

Karlodinium veneficum isolated from the SRE was grown in f/2 medium in a culture chamber at 100 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$, 12:12 L:D, 20 degrees C, salinity 15. The unialgal culture (AUS#7) was isolated using single cell isolation by micropipette, from a dense sample collected at station SUC on March 29, 2005. Cell pellets from AUS #7 of 10^5 to 10^6 cells were obtained by centrifugation (IEC Centra CL2, Thermo Fisher Scientific, Waltham, MA, USA) at 3,000 rpm (1,600 g) and frozen at 20 °C prior to DNA extraction. DNA was isolated using a CTAB detergent method followed by chloroform extraction and precipitation of the aqueous phase with isopropanol (Doyle and Doyle 1987). The DNA was diluted based on 230 nm absorbance, and 100 ng was used in a 20 μL PCR reaction in the following buffer: 3 mM MgCl₂, 500 mM Tris-HCl pH 8.3, 500 $\mu\text{g mL}^{-1}$ BSA, 2 mM dNTPs, and 4 pmoles of primer (all concentrations final). The temperature was cycled from 15 s denaturing at 94 °C to a 55 °C annealing for 15 s followed by 30 s of extension at 72 °C for 35 cycles. The primers used were based on primers suggested by Litaker et al. (2003): ITSFor CTGCGAAGCTATCGCTATT and ITSRev TGAGGGAATCCTATTTAG designed to cross from

ITS1 to the LSU rRNA including most of ITS1, 5.8S, and ITS2. PCR amplicons were purified with equal volumes of 20% w/v polyethylene glycol (mw 8000) containing 2.5 M NaCl, pelleted by centrifugation, and washed with 70% ethanol. Sequencing reactions were performed and run on an ABI 3130XL sequencer according to the manufacturer's protocol. Sequences were edited using Sequencher 4.5 software (Genecodes Corp., Ann Arbor, MI, USA) and exported to MacClade 4.05 (Sinauer Associates Inc., Sunderland, MA, USA) for manual alignment using GenBank sequences AF352365–7 for *K. veneficum* as a reference.

Large Scale Karlotoxin Isolation from a Natural Bloom

A total of 4.8 L of SRE water containing the bloom of *K. veneficum*, collected from sites SUC (March 31, 2005) and WMP, MEA, and SUC (April 4, 2005), were filtered through a GF/F (Whatman) filter, and the filtrate then passed through a 55 g tC18 cartridge (AnaLogix, Sorbtech Technologies) to collect dissolved organic material including toxin. The column was first activated by washing with 200 mL methanol followed by 200 mL HPLC-grade water. The column was stored in at 4 °C and then transported in a cooler with an ice pack until further processing in Baltimore, MD less than 2 weeks later. In Baltimore, the tC18 cartridge was washed successively with 200 mL volumes of dH₂O, 20% MeOH, and 40% MeOH (all discarded after testing for hemolytic activity). The 60% MeOH and 80% MeOH eluates, anticipated to contain partially purified KmTx, were retained, dried under vacuum at 40 °C, and analyzed by LC-MS. The concentrated toxin from the pooled 60-80% tC18 eluent was then purified using reverse phase chromatography as on a semi-preparative scale C18 column (Phenomenex Hyperprep HS C18-80S, 250 X 10 mm, 5 μ) subjected to a 4 mL min⁻¹ binary methanol/water gradient from 10% to 95% methanol over 45 minutes using an Agilent 1100 Series LC/MSD system, comprising of binary pump system, autosampler and diode array detector (DAD) with a micro high-pressure flow cell (6 mm path length, 1.7 μ L volume),

fraction collector and quadrupole mass spectrometer (G1956A SL) equipped with an electrospray ionization (ESI) interface. Toxin peaks were detected over the wavelength range 190 to 950 nm. Based on the UV/vis spectra the absorption at 225 nm was used to detect KmTx 1 while absorption at 235 nm was used to detect KmTx 2. The entire UV/Vis spectra were saved for each UV detectable peak. Fractions corresponding to KmTx^{Perth} 2-1 and KmTx^{Perth} 2-2 were collected using a fraction collector (61364A, AFC, Agilent) based on known retention times. Fractions were pooled as appropriate, dried under vacuum and resuspended in 80% methanol:20% water. From the total 4.8 L, 23 mg of KmTxPerth 2-1 and 1 mg of KmTxPerth 2-2 were obtained.

In situ Karlotoxin sampling and analysis

Routine sampling of KmTx from Swan River samples was done using a modification of the procedure described in Bachvaroff et al. (2008). Five mL of SRE water collected for cell count sampling were pushed through 13 mm 0.2 μ m pore size PTFE filters, labeled, wrapped in small manila envelopes and then stored at 4 °C, thus collecting both all toxin in the sample whether cell-associated or free. At the end of the sampling season, these syringe filters were shipped in a cooler with ice packs overnight from Perth to Baltimore. In Baltimore, the filters were eluted with 1 mL MeOH into 2 mL dH₂O to yield a crude extract of KmTx in 33% MeOH. Quantification of KmTx used HPLC-MS methods detailed in Bachvaroff et al. (2008). Ions corresponding to KmTx 2 (1367.8 amu), and a 1365 amu ion that showed strong hemolytic activity, were summed to obtain KmTx abundance for this study. The 1365 ion was in lower abundance than 1367.8, averaging 24% ($\pm$ 20.7%) of total ion counts for KmTx.

Hemolytic Assay and Fish Bioassay of KmTx

The HPLC fractions were collected for 20 s during the first 32 m of the chromatography gradient into 96 well plates. The hemolytic assay is described in detail elsewhere (Eschbach, et al., 2001). Briefly, rainbow trout (*Oncorhynchus mykiss*) erythrocytes were extracted from the caudal vein and the blood washed three times with three volumes of buffer mL^{-1} of packed erythrocytes using the incubation buffer described below minus CaCl_2 . Following the third wash the cells were resuspended in 3 mL of buffer mL^{-1} of packed cells. Resuspended cells were diluted 1:20 in buffer and 200 μL of diluted, washed erythrocytes were incubated in the presence of 10 μL of each HPLC fraction for 30 minutes in a buffer of 150 mM NaCl, 3.2 mM KCl, 1.25 mM MgSO_4 , 12.2 mM Tris Base and 3.75 mM CaCl_2 . Assays were run in 96 well, V-bottom, non-treated, polystyrene plates (Corning Inc.) sealed with Falcon 3073 pressure sensitive film (Becton Dickinson Labware). Assays were incubated on an orbital shaker (80-100 rpm) at 20 °C for 1 hour. Plates were then centrifuged at 2500 g for 5 min and the supernatant (100 μL) was transferred to a flat bottom 96 well polystyrene plate, where the absorbance of released hemoglobin was read at 540 nm using a microtiter plate spectrophotometer (Model Spectramax 340, Molecular Devices). Saponin (10 μg) (from Quillaja bark; Sigma Chemical Co.) as used as a hemolytic positive control.

To examine ichthyotoxicity of the isolated karlotoxins, sheepshead minnows (*Cyprinodon variegatus*, purchased as larvae from Chesapeake Cultures, Inc., Hayes, Virginia, and raised at salinity 12 in-house following procedures described previously for zebrafish (*Danio rerio*)) were exposed to increasing concentrations of toxin. Sheepshead minnow (60–90 d old; 2 cm long) were exposed as described (Deeds et al. 2006) except that O_2 -saturated 12% artificial seawater (Instant Ocean Brand) was used as the exposure medium. Sheepshead minnows were exposed to control (0.4% methanol), 267, 668, 1338, 2675 $\text{KmTx}^{\text{Perth}}$ 2-1 and 0, 69.3, 173, 346, and 693 $\text{KmTx}^{\text{Perth}}$ 2-2 ng/ mL (fish/beaker; $n = 2$). Fish were checked for mortality hourly, and any fish that did not die by

237 4 h were killed then, as described previously (Deeds et al. 2006). Whole gills were dissected,
 238 preserved and examined as described (Deeds et al. 2006).

239

240

241 RESULTS

242 *Molecular characterization of Karlodinium veneficum strain*

243 The 541 base ITS sequence from a cultured SRE *K. veneficum* (AUS #7, GENBANK# KP639236)
 244 had three single nucleotide differences and a six base deletion when compared to the ITS for CCMP
 245 415, isolated from the North Sea (AJ557026) (Fig. 2). The ITS2 for *K. veneficum* was able to be
 246 folded into a typical secondary structure with four base paired ‘stems’
 247 (<http://its2.bioapps.biozentrum.uni-wuerzburg.de>). The variant ITS2 sites could then be sorted into
 248 compensatory changes occurring in both bases of a base paired stem and non-compensatory changes
 249 found in single stranded loops (Gottschling and Plötner 2004). Two single nucleotide differences
 250 occurred in the single stranded loop at the end of the third base paired stem in ITS2 and were thus
 251 not compensatory changes. The six base deletion and the third nucleotide substitution shortened the
 252 fourth ITS2 stem from 9 base pairs to five and were compensatory changes.

253

254 *Physical and chemical conditions in the Swan River Estuary*

255 Discharge from the Avon River and Bennet Brook (confluence with the SRE just above
 256 SUC, Fig. 1) was negligible through DOY 84, followed by elevated flow in Bennet Brook (DOY
 257 86) and the Avon River (DOY 96) (Fig. 3). Total N in the surface SRE declined from 60 $\mu\text{mol L}^{-1}$
 258 or greater at the 35 km station (SUC), to 10 – 20 $\mu\text{mol L}^{-1}$ by the 6 km station (BLA) (Fig. 4A).
 259 Nitrate was low ($<0.2 \mu\text{mol L}^{-1}$) throughout the surface SRE except DOY >120 (May) when values
 260 increase throughout most of the estuary (Fig. 4B). Ammonium averaged 0.83 $\mu\text{mol L}^{-1}$ throughout

the surface SRE between DOY 1 - 90 (Jan to March), then showed elevated values associated with the bloom (average $2 \mu\text{mol L}^{-1}$) and thereafter (Fig 4C). Total P showed a pattern similar to that of total N, with values of $8 - 10 \mu\text{mol L}^{-1}$ in the surface waters of the upper SRE declining to $< 1 \mu\text{mol L}^{-1}$ downstream, and some of the highest values of the sampling period associated with the bloom (Fig. 4D). Dissolved phosphate was generally $1 - 2 \mu\text{mol L}^{-1}$ in the surface waters of the upper SRE and declined downstream, with elevated values of $2 - 4 \mu\text{mol L}^{-1}$ observed at the bloom (Fig. 4E). Particulate elemental analyses of high *K. veneficum* density water samples yielded average C:N (mol) of 8.0, and C:P (mol) of 97.3 (Table 1).

At the beginning of the bloom (DOY 76 - 104 and 30 - 42 km upstream), surface temperature was $25 - 26 ^\circ\text{C}$, but dropped to $20 - 21 ^\circ\text{C}$ between DOY 89 - 94 (March 30 - April 4) (Fig. 5A). A drop in salinity from ~ 27 to ~ 21 between DOY 90 - 101 (March 31 to April 11) was observed (Fig. 5B), associated with a brief stratification of the water column observed most strongly (delta salinity $6 - 9$) on DOY 101 (April 11) (Fig. 5C). A pronounced drop in salinity occurred toward the end of the study period (after DOY 140, Fig. 5B), associated with elevated flow in the Avon River (Fig. 3). Surface D.O. (Fig. 5D) was supersaturated ($9 - 11 \text{ mg L}^{-1}$) initially, while bottom water D.O. (Fig. 5E) was low ($< 1 - 6 \text{ mg L}^{-1}$) at the site of the bloom as well as at stations downstream (to 25 km upstream) of the main *K. veneficum* biomass accumulation. Low values of surface D.O. first appeared between DOY 101 - 108 (April 11 and April 18), centered at the 27 km station (MAY) but including most of the SRE from 35 km to the mouth (Fig. 5D).

Phytoplankton community structure leading up to the bloom of K. veneficum

Shifts in phytoplankton community composition over DOY 1-94 (Jan to March) were observed at three stations (SUC, KIN, and RON), showing a decline in planktonic diatoms coincident with increasing *K. veneficum* cell abundance (Fig. 6, DOY 1 - 94). The decline in

diatoms involved a shift in dominant forms within that group, from chain-forming *Thalassiosira* / *Skeletonema* to small cf. *Cyclotella* by early March (WA DOE, data not shown). Cryptophytes, a potential food source of *K. veneficum*, were present but showed no trend with time or correlations with *K. veneficum* abundance (Fig 6). No significant trends were observed for other phytoplankton groups (chlorophytes, prasinophytes, euglenophytes, chrysophytes) over the time period leading up to the bloom of *K. veneficum*.

LC-MS characterization of KmTx from the Swan River Estuary

The total ion chromatograph (TIC) for partially-purified material showed three major peaks, two associated with activity in the hemolytic assay (Fig. 7). The major ions associated with these peaks were in order of elution: 1367.8, 1365.8 and 1349 amu (Fig. 7 insets). Further purification of these peaks resulted in two fractions, 'Perth 1' or KmTx^{Perth}2-1 (Fig. 8A) dominated by the 1367.8 amu ion, and 'Perth 2' or KmTx^{Perth}2-2 containing a mixture of the 1367.8, 1349 and 1365.8 ions (Fig. 8B). Both toxin peaks were classified as KmTx2 types due to higher absorbance at 235 nm on UV absorption spectra (data not shown).

Biomass of Karlodinium veneficum and KmTx concentrations in the Swan River Estuary

Elevated phytoplankton biomass as Chl *a* (Fig. 9A) and *K. veneficum* cell concentrations (Fig. 9B; $\geq 10^3 \text{ mL}^{-1}$) were observed DOY 76 – 104 (March 17 – April 14, 2005) and between 30 - 42 km from the mouth of the SRE (Fig. 9B). Karlotoxin levels up to 1052 ng mL^{-1} were observed during this time period in the same region (Fig. 9C). Following the fish kill, low levels of *K. veneficum* ($10^2 - 10^1 \text{ cells mL}^{-1}$) and KmTx were detectable in the lower SRE with the center of the population moving approximately 20 km downstream as the time series progressed (Fig. 9B). Swan River Trust (SRT) crews observed a fish kill on the morning of April 13, 2005 (DOY 103) after

reports of sluggish and dying fish from residents (SRT Media Statement, April 13, 2005). Examination of hematoxylin and eosin stained gill arches from a moribund black seabream (*Acanthopagrus butcheri*) collected at the Swan River fish kill on April 10, 2005 showed loss of intra-lamellar spacing and secondary lamellar fusion with extensive hyperplasia of gill epithelial tissue, similar to damage caused exposure to pure KmTx2 in laboratory studies (Deeds et al. 2006) (Fig. 10).

Log toxin concentration (ng KmTx 2 mL⁻¹) correlated with log *K. veneficum* concentration (cells mL⁻¹) (Fig 11A, Pearson $r = 0.73$, $p < 0.0001$). Considering only samples where log *K. veneficum* mL⁻¹ was $\geq 10^3$, cellular toxin quota (pg KmTx cell⁻¹) averaged 4.5 ± 7.50 (median 2.8) pg KmTx cell⁻¹. Cellular KmTx (pg cell⁻¹) showed a weak positive but significant regression with total KmTx (ng mL⁻¹) ($r^2 = 0.14$, $p = 0.004$). Time series plots of KmTx cell⁻¹ (Fig 11B) for stations with cell density >100 mL⁻¹ show a high degree of variability, with a spike in cellular toxicity at SUC on DAY 108, when *K. veneficum* abundance was 2702 cells mL⁻¹.

Both purified karlotoxins exhibited dose-dependent ichthyotoxicity against juvenile sheepshead minnows (*Cyprinodon variegatus*) (Fig. 12A & B). For KmTx ‘Perth 1’, the lethal dose was between 267 – 668 ng mL⁻¹, while for KmTx ‘Perth 2’ the lethal dose was between 173 – 346 ng mL⁻¹.

DISCUSSION

The occurrence of toxic algal blooms is a relatively recent phenomenon in the SRE (Thompson 1998, Chan and Hamilton 2001, Horner-Rosser and Thompson 2001). The bloom described here is one of a series of recurrent ichthyotoxic bloom of *K. veneficum* observed in the SRE 2002 – 2006, 2010, and 2012. Gill histopathology (Fig. 10) observed on fish collected from the fish kill described here was consistent with damage from KmTx as described by Deeds et al. (2006).

Water quality properties measured in association with this bloom, including bottom layer hypoxia, suggest nutrient over-enrichment, which is considered a key factor driving the proliferation of HAB in new systems (Glibert et al. 2005, Heisler et al. 2008). The results of this study allow us to better understand the ecological context of ichthyotoxic blooms of *K. veneficum*, including co-occurrence of karlotoxin-containing cells and hypoxia (Fig. 13), both of which likely contributed to the observed fish kill.

The role of freshwater inputs in blooms of K. veneficum in the SRE

Results of this study are consistent with observations that blooms of *K. veneficum* develop under low flow conditions in the upper SRE, and that elevated flow, typically beginning in May / June, dissipates upper SRE blooms of *K. veneficum*. Abloom of *K. veneficum* documented in the Neuse River Estuary (Hall et al. 2008) similarly occurred under low flow, high temperature / salinity conditions. Linkages between river flow and phytoplankton succession have been made previously in the SRE, including Twomey and John (2001) who showed that rainfall, river flow and the positioning of the salt wedge in the estuary dictates the typical seasonal succession of phytoplankton in the SRE. Toxic blooms *Microcystis* were linked to anomalously high summer river flows (Robson and Hamilton 2003). The displacement of the *K. veneficum* population down estuary 10-20 km (Fig. 9B), to the mouth of the Canning estuary, is another potentially important role of river flow in this system that deserves further attention. Whether or not *K. veneficum* is transported back up estuary in the lower layer upon relaxation of seasonal flows (i.e. Tyler and Seliger 1978) should be investigated. Ecosystem managers can use rainfall and river flow data to help forecast bloom location along the estuary and imminent fish kills during years when a dense bloom of *K. veneficum* is present in the upper SRE. Further, understanding the linkages between ichthyotoxic blooms and river flow in the SRE will aid understanding of how changing patterns of

rainfall predicted for Western Australia (Hughes 2011, CSIRO State of the Climate Reports 2010, 2012 (<http://www.csiro.au/resources/State-of-the-Climate>)) might impact the timing and / or magnitude of blooms of *K. veneficum* in this system.

Phytoplankton community composition shift leading to the bloom of K. veneficum

Results of this study and continuing observations of annual blooms of *K. veneficum* (2010, 2012) suggest that while conditions in the SRE in late summer – autumn have historically been favorable for dinoflagellates, recently toxic *K. veneficum* has become a more prominent member of the dinoflagellate assemblage in the SRE. The bloom of *K. veneficum* and fish kill observed in March – April was preceded by a 3 month period over which planktonic diatoms were displaced by *K. veneficum* (Fig. 6) in the upper SRE. Physical / chemical conditions likely disfavored large diatoms over this time period, as freshwater (and thereby nitrate) inputs were low (Fig. 3), and the inorganic nutrient pools in surface waters suggested N- limitation (Thompson and Hosja 1996, Twomey and John 2001). The observed shift from chain-forming to small solitary diatoms (*Cyclotella*) over this time period points to low nutrients / turbulence in the surface layer. A similar shift over the period January – March, from diatoms dominated by *S. costatum* to dinoflagellates, was also reported by Thompson (1998) and Hamilton et al. (1999), but the dinoflagellates in these reports were numerically dominated by *Scrippsiella* sp. or *Gymnodinium* and *Prorocentrum* with no mention of *K. veneficum* or ichthyotoxicity. Further research is required to determine what contributes to the differential success of *K. veneficum* over other flagellates during late summer – autumn in the upper SRE, including investigation of the anti-grazing / allelopathic activity of karlotoxins (Adolf et al. 2007, 2008; Stoecker et al. 2008, Brownlee et al. 2008, Waggett et al. 2008, Vaque et al. 2005) or mixotrophic nutrition (Li et al. 1996). Although mixotrophic grazing by *K. veneficum* was not quantified in this study, feeding by *K. veneficum* upon addition of cultured

cryptophytes to SRE water collected at the bloom was observed (JEA, Pers. Obs.), indicating at least potential mixotrophic growth of SRE *K. veneficum* populations. Work by Griffin et al. (2001) suggests that anti-grazing properties of KmTx would be advantageous as dinoflagellate blooms in the upper SRE are strongly top-down regulated, and escape from grazing pressure is increasingly recognized as a strategy permitting some species to form blooms (Iriogen et al. 2005, Mitra and Flynn 2006, Sunda et al. 2006, Smayda 2008).

Physical and chemical conditions associated with the SRE bloom

By the time the bloom of *K. veneficum* was fully developed in the upper SRE, elevated Chl *a* and *K. veneficum* cell counts (DOY 76 – 90) occurred in warm surface waters that were weakly salinity-stratified (Fig. 5). Surface nutrient concentrations were low preceding and during the bloom, with molar DIN:DIP close to 1, typical for the SRE (Thompson and Hosja 1996). The bloom was initially located over a patch of hypoxic bottom water that stretched from above the 35 km station to the 25 km station (Fig. 5E), environmental conditions that are not uncommon in this portion of the SRE (Stephens and Imberger 1996, Thompson 1998, Hamilton et al. 2001) associated with strong salinity stratification (Kurup and Hamilton 2002) and elevated inorganic nutrient availability (Hamilton et al. 1999). Although not specifically addressed in this study, the ability of *K. veneficum* to undergo vertical migrations (*sensu* Hall and Paerl 2011) may give this species an advantage over other forms in the stratified upper SRE environment, although this would not explain the differential success of *K. veneficum* over other flagellates capable of vertical migrations.

Lysis of *K. veneficum* cells is an important step leading to toxin release and ichthyotoxicity (Deeds et al. 2002, Mooney et al. 2010, Dorantes-Aranda et al. 2011), and the beginning of seasonal rainfall that occurred during the bloom period may have played an important role in cell disruption leading to the fish kill investigated here. After DOY 84, flow in the Avon River and Bennet Brook

increased with an associated drop in surface salinity (5-10) and water temperatures of 4 °C, which could have contributed to osmotic lysis of the cells. Between DOY 90 and 100, bottom water oxygen levels increased compared to the prior period, but subsequently returned to hypoxic levels after DOY 101 until after DOY 108 (Fig. 5E). Low surface oxygen levels coincided with relaxation of salinity stratification and the crash of the bloom of *K. veneficum* on DOY 107 (Fig. 5E). Thus, in the narrow and shallow upper SRE, fish would have been exposed to both water column hypoxia and free KmTx simultaneously at the time of the observed fish kill event. While it is speculated here that changes in salinity and temperature led to *K. veneficum* disruption, positive identification of the factors leading to *K. veneficum* disruption in situ is an important area for future research to better understand the role of *K. veneficum* in fish kills.

Cellular toxicity of K. veneficum during the bloom period

The KmTx congener isolated from large-volume SRE samples (Fig. 8) was identical to KmTX^{Penth} 2-1 described from a clonal culture of *K. veneficum* isolated from the SRE in 2005 (strain KVSRO1) with major ions at 1367 and 1425 amu (Mooney et al. 2009). While the Australian SRE *K. veneficum* strains were considered conspecific with other isolates based on LSU sequences (Mooney 2009), the putative presence of compensatory changes in ITS2 structure that reported here (Fig. 2) and cellular toxicity (~0.1 pg KmTx cell⁻¹) (J.E. Adolf, pers. obs.) of the SRE isolate (Aus #7) relative to CCMP 415 suggests cryptic species may be present (Gottschling and Plötner 2004), although further investigation is needed. A number of *Karlodinium* species have recently been described from Australia (de Salas et al. 2005, de Salas et al 2008), but hemolytic karlotoxins were only present in *K. veneficum* (Mooney et al. 2009). However, immobilization and ingestion of copepods by *K. armiger* suggests uncharacterized toxins may be present in other species of the *Karlodinium* genus (Berge et al. 2012).

The level of cellular toxicity calculated for *K. veneficum* in this bloom (mean 4.4, median 2.8 pg KmTx cell⁻¹) is within the previously reported range of laboratory cultures (Deeds et al. 2004, Adolf et al. 2009, Bachvaroff et al. 2009) including an SRE strain of *K. veneficum* (KmTX^{Swan} 2, Mooney et al. 2009). Cellular toxin in the SRE was similar to values reported for 2005 – 2007 blooms observed in the Inner Harbor of Baltimore, MD (Adolf et al. 2008). Cellular toxicity of *K. veneficum* varies among strains grown under nutrient-replete conditions (Bachvaroff et al. 2009), and also as a result of inorganic N or P limitation (Adolf et al. 2009, Mooney et al. 2010). The particulate C, N, and P measurements measured for high density *K. veneficum* water samples from the SRE {C:N (8.0) and C:P (97.3)} suggest N-limitation when compared to nutrient replete averages from six cultured *K. veneficum* strains, where C:N averaged 4.8 and C:P averaged 136 (Adolf et al. 2009). Li et al. (2002) reported molar C:N values of 6.4 and C:P values of 164.4 for single strain (CCMP 1974) grown under 27 variable N and P conditions. The DIN:DIP values near unity surrounding the bloom also suggest N-limitation. Thus, N-limitation in surface waters at the time of the bloom likely increased the cellular toxicity of *K. veneficum* 2-5 fold (Adolf et al. 2009) compared to non-nutrient limited values, contributing to the total concentrations of KmTx observed during this event.

Some of the observations of elevated cellular toxicity levels in this study might have resulted from low *K. veneficum* cell abundance and a loss of precision in cell counts at low cell abundance. The *in situ* toxin concentrations leveled off below 10³ cells mL⁻¹, leading to cellular toxicity estimates that increased over two orders of magnitude at cell abundance below 10³ cells per mL. This most likely represents loss of precision in the ability to accurately quantify low cell abundance relative to the high precision with which KmTx can be measured by LC-MS (~1.2 ng KmTx mL⁻¹; Bachvaroff et al. 2008). Alternatively, although KmTx is labile and not likely to last more than one day after cell senescence (Deeds et al. 2002), some of the anomalously high values

for cellular toxicity may represent senescent regions of a bloom where high free toxin concentrations remained but cell abundance was diminished. For instance, the spike in cellular toxicity observed on DOY 108 at SUC (Fig. 11B) coincided with relatively low cell counts (2702 cell mL⁻¹).

*Ichthyotoxicity associated with the 2005 SRE bloom of *K. veneficum*: KmTx and hypoxia*

Determination of acute ichthyotoxicity levels of between 173 – 346 ng mL⁻¹ for purified KmTx ‘Perth2-1’, and between 267 – 668 ng mL⁻¹ for purified KmTx ‘Perth2-2’ (Fig. 12) is in agreement with the conservative environmental threshold for ichthyotoxicity of 500 ng mL⁻¹ suggested by Deeds et al. (2006). KmTx levels observed in the SRE (where >1,000 *K. veneficum* mL⁻¹ were present) were low comparably, with a median KmTx concentration of 33.4 ng mL⁻¹ (Table 2) and average cell densities of 42,000 cells mL⁻¹, and thus were unlikely to be acutely ichthyotoxic on their own. However, Deeds et al. (2006) observed sub-lethal gill damage in the laboratory at 50 ng mL⁻¹ KmTx2 (Deeds et al. 2006), and that examination of fish recovered from this SRE fish kill determined gill damage consistent with KmTx (Fig 10), which may have exacerbated effects of hypoxia on fish.

Results from the SRE show a sequence of events involving two known ichthyotoxic factors co-occurring in a eutrophic estuary – high levels of KmTx (Deeds et al. 2002, Deeds et al. 2006) and water column hypoxia (Diaz and Rosenberg 2008). Hypoxia ranks with HAB as a global symptom of ecosystem decline (Cloern 2001, Kemp et al. 2005, Diaz and Rosenberg 2008) and can co-occur spatially and temporally with HAB, including *K. veneficum* (Hall et al. 2008). Hypoxic or anoxic conditions are common in the upper SRE (Stephens and Imberger 1996, Hamilton et al. 2001), associated with water column stratification (Kurup and Hamilton 2002). Further, night time oxygen levels were likely lower than the daytime values that were measured, potentially increasing

477 their role in this fish kill. Karlotoxin affects fish by damaging cells along the exposed region of the
478 gills (Deeds et al. 2006), resulting in death by suffocation in acute cases, but impaired oxygen
479 uptake in sub-acute exposures that can potentially exacerbate sensitivity of fishes to hypoxic
480 conditions. Thus, observations of toxin levels and DO in the SRE suggest several important lines of
481 research including an examination of the effects of hypoxia on *K. veneficum* cell lysis (necessary for
482 ichthyotoxicity, Deeds et al. 2002), the dose dependency of KmTx on fish exposed to hypoxia
483 (sensu Deeds et al. 2006), and the effects of hypoxia on the lability of free KmTx that has been
484 released from cells.

485

REFERENCES

- Abbot, B. C., and Ballantine, D. 1957. The toxin from *Gymnodinium veneficum* Ballantine J. Mar. Biol. Assoc. U.K. 36, 169-189.
- Adolf, J. E., Stoecker, D.K., and Harding, L.W. Jr. 2006a. The balance of autotrophy and heterotrophy during mixotrophic growth of *Karlodinium micrum* J. Plank. Res. 28, 737-751.
- Adolf, J. E., Bachvaroff, T.R., Krupatkina, D.N., Nonogaki, H., Brown, P.J.P., Lewitus, A.J., Harvey, H.R., and Place, A.R. 2006b. Species specificity and potential roles of *Karlodinium micrum* toxin African Journal of Marine Science, Harmful Algae 2004. P. G. C. Pitcher, T.A., Verheye, H.M. 28, 415-421.
- Adolf, J. E., Bachvaroff, T.R., Krupatkina, D.N., and Place, A.R. 2007. Karlotoxin mediates grazing by *Oxyrrhis marina* on strains of *Karlodinium veneficum* Harmful Algae 6, 400-412.
- Adolf, J. E., Bachvaroff, T.R., and Place, A.R. 2008. Can cryptophyte abundance trigger toxic *Karlodinium veneficum* blooms in eutrophic estuaries? Harmful Algae 8, 119-128.
- Adolf, J. E., Bachvaroff, T.R., and Place, A.R. 2009. Environmental modulation of Karlotoxin levels in strains of the cosmopolitan dinoflagellate, *Karlodinium veneficum* (Dinophyceae). J. Phycol. 45,176-192.
- Bachvaroff, T. R., Adolf, J.E., and Place, A.R. 2009. Strain variation in *Karlodinium veneficum* (Dinophyceae): toxin profiles, pigments, and growth characteristics. J. Phycol 45, 137-153.

510

511 Bachvarroff, T. R., Adolf, J.E., Squier, A., Harvey, H.R., and Place, A.R. 2008. Characterization
 512 and quantification of karlotoxins by liquid chromatography-mass spectrometry. *Harmful Algae* 7,
 513 473-484.

514

515 Ballantine, D. 1956. Two new marine species of *Gymnodinium* isolated from the Plymouth area. *J.*
 516 *Mar. Biol. Assoc. U.K.* 35, 473-484.

517

518 Berge, T., Poulsen, L.K., Moldrup, M., Daugberg, N., and Hansen, P.J. 2012. Marine microalgae
 519 attack and feed on metazoans. *The ISME Journal* 2012, 1-11.

520

521 Bergholtz, T., N. Daugbjerg, Ø. Moestrup, and M. Ferná'ndez-Tejedor 2005. On the identity of
 522 *Karlodinium veneficum* and description of *Karlodinium armiger* Sp. Nov. (Dinophyceae), based on
 523 light and electron microscopy, nuclear-encoded LSU rDNA, and pigment composition. *J. Phycol.*
 524 42, 170-193.

525

526 Brownlee, E. F., Sellner, S.G., Sellner, K.G., Nonogaki, H., Adolf, J.E., Bachvaroff, T.R., and
 527 Place, A.R. 2008. Responses of *Crassostrea virginica* (Gmelin) and *C. ariakensis* (Fujita) to bloom-
 528 forming phytoplankton including ichthyotoxic *Karlodinium veneficum* (Ballantine). *J. Shellfish Res.*
 529 27, 581-591.

530

531 Calbet, A., Bertos, M.m Fuentes-Grunewald, C., Alacid, E., Figueroa, R., Renom, B., and Garcia, E.
 532 2011. Intraspecific variability in *Karlodinium veneficum*: growth rates, mixotrophy and lipid
 533 composition. *Harmful Algae* 10,654-667.

534

535 Chan, T. U., and Hamilton, D.P. 2001. Effect of freshwater flow on the succession and biomass of
 536 phytoplankton in a seasonal estuary. Mar. Freshwater Res. 52, 869-884.

537

538 Chan, T. U., Hamilton, D.P., Robson, B.J., Hodges, B.R., and Dallimore, C. 2002. Impacts of
 539 hydrological changes on phytoplankton succession in the Swan River, Western Australia. Estuaries
 540 25, 1406-1415.

541

542 Cloern, J. E. 2001. Our evolving conceptual model of the cultural eutrophication problem. Mar.
 543 Ecol. Progr. Ser. 210, 223-253.

544

545 Deeds, J. R., Terlizzi, D.E., Adolf, J.E., Stoecker, D.K., and Place, A.R. 2002. Toxic activity from
 546 cultures of *Karlodinium micrum* (= *Gyrodinium galatheanum*) (Dinophyceae)--a dinoflagellate
 547 associated with fish mortalities in an estuarine aquaculture facility. Harmful Algae 1, 169-189.

548

549 Deeds, J. R., Kibler, S.R., Tester, P.A., and Place, A.R. 2004. Geographic strain variation in toxin
 550 production in *Karlodinium micrum* (Dinophyceae) from Southeastern United States. Harmful Algae
 551 2002 Proceedings. K. A. Steidinger, Landsberg, J.H., Tomas, C.R., and Vargo, G.A. St. Petersburg,
 552 Florida Institute of Oceanography, and Intergovernmental Oceanographic Commission of
 553 UNESCO, 145-147.

554

555 Deeds, J. R., Reimschuessel R., and Place, A. R. 2006. Histopathological effects in fish exposed to
 556 the toxins from *Karlodinium micrum* (Dinophyceae). J. Aquat. Animal Health 18, 136-148.

557

- 558 Deeds, J. R., and Place, A.R. 2006. Sterol specific membrane interactions with the toxins from
559 *Karlodinium micrum* (Dinophyceae) - A strategy for self protection? African J. Mar. Sci. Harmful
560 Algae 2004 28, 421-427.
- 561
- 562 Diaz, R. J., and Rosenberg, R. 2008. Spreading dead zones and consequences for marine
563 ecosystems. Science 321, 926-929.
- 564
- 565 Dorantes-Aranda, J.J., Waite, T.D., Godrant, A., Rose, A.L., Tovar, C.D., Woods, G., Hallegraeff,
566 G.M. 2011. Novel application of a fish gill cell line assay to assess ichthyotoxicity of harmful
567 marine microalgae. Harmful Algae 10, 366-373
- 568
- 569 Doyle, J. J. and Doyle, J. L. 1987. A rapid DNA isolation procedure for small quantities of fresh
570 leaf tissue. Phytochem. Bull. 19, 11–15.
- 571
- 572 Eschbach, E., Scharsack, J., John, U., Medlin, L. 2001. Improved erythrocyte lysis assay in
573 microtitre plates for sensitive detection and efficient measurement of haemolytic compounds from
574 ichthyotoxic algae. J. Appl. Toxicol. 21, 513–539.
- 575
- 576 Fensin, E. E. 2004. Occurrence and ecology of the dinoflagellate *Karlodinium micrum* in estuaries
577 of North Carolina, USA. In K. A. Steidinger, Landsberg, J.H., Tomas, C.R., & Vargo, G.A. [Eds.]
578 Harmful Algae 2002 Proceedings. St. Petersburg, Florida Institute of Oceanography, and
579 Intergovernmental Oceanographic Commission of UNESCO, pp. 62-4.
- 580

- 581 Garces, E., Fernandez, M., Penna, A., Van Lenning, K., Gutierrez, A., Camp, J., and Zapata, M.
 582 2006. Characterization of NW Mediterranean *Karlodinium* spp. (Dinophyceae) strains using
 583 morphological, molecular, chemical, and physiological methodologies. J. Phycol. 42, 1096-1112.
 584
- 585 Glibert, P. M., Seitzinger, S., Heil, C.A., Burkholder, J.M., Parrow, M.W., Codispoti, L.A., and
 586 Kelly, V. 2005. The role of eutrophication in the global proliferation of harmful algal blooms.
 587 Oceanography 18, 198-209.
 588
- 589 Goshorn, D., Deeds, J., Tango, P., Poukish, C., Place, A.R., McGinty, M., Butler, W., Luckett, C.,
 590 & Magnien, R. 2004. Occurrence of *Karlodinium micrum* and its association with fish kills in
 591 Maryland estuaries. In K. A. Steidinger, Landsberg, J.H., Tomas, C.R., & Vargo, G.A. [Eds.]
 592 Harmful Algae 2002 Proceedings. St. Petersburg, Florida Institute of Oceanography, and
 593 Intergovernmental Oceanographic Commission of UNESCO, pp. 361-63.
 594
- 595 Gottschling, M. & Plötner, J. 2004. Secondary structure models of the nuclear internal transcribed
 596 spacer regions and 5.8S rRNA in Calciodinelloideae (Peridiniaceae) and other dinoflagellates.
 597 Nucleic Acids Res., 32, 307-315.
 598
- 599 Griffin, S. L., Herzfeld, M., and Hamilton, D.P. 2001. Modeling the impact of zooplankton grazing
 600 on phytoplankton biomass during a dinoflagellate bloom in the Swan River Estuary, Western
 601 Australia Ecological Engineering 16, 373-394.
 602
- 603 Hall, N. S., Litaker, R.W., Fensin, E., Adolf, J.E., Bowers, H.A., Place, A.R., and Paerl, H.W. 2008.
 604 Environmental factors contributing to the development and demise of a toxic dinoflagellate

605 (*Karlodinium veneficum*) bloom in a shallow, eutrophic, lagoonal estuary. Estuaries Coasts 31,402-
606 418.

607
608 Hall, N.S., and Paerl, H.W. 2011. Vertical migration patterns of phytoflagellates in relation to light
609 and nutrient availability in a shallow microtidal estuary. Mar Ecol Progr Ser 425,1-9.

610
611 Hamilton, D. 2000. Record summer rainfall induced the first recorded major cyanobacterial bloom
612 in the Swan River. Journal of the Environmental Engineering Society, Institute of Engineers 1, 25.

613
614 Hamilton, D. P., Thompson, P.A., and Kurup, R. 1999. Dynamics of dinoflagellate blooms in the
615 Swan River Estuary. Proceedings of the Vth International Wetlands.

616
617 Hamilton, D. P., Chan, T., Robb, M.S., Pattiaratchi, C.B., and Herzfeld, M. 2001. The hydrology of
618 the upper Swan River Estuary with focus on an artificial destratification trial. Hydrol. Processes 15,
619 2465-2480.

620
621 Heisler, J., Glibert, P.M., Burkholder, J.M., Anderson, D.M., Cochlan, W., Dennison, W.C., Dortch,
622 Q., Gobler, C.J., Heil, C.A., Humphries, E., Lewitus, A., Magnien, R., Marshall, H.G., Sellner, K.,
623 Stockwell, D.A., Stoecker, D.K., and Suddleson, M. 2008. Eutrophication and harmful algal
624 blooms: a scientific consensus. Harmful Algae 8, 3-13.

625
626 Horner-Rosser, S. M. J., and Thompson, P.A. 2001. Phytoplankton of the Swan-Canning Estuary: a
627 comparison of nitrogen uptake by different bloom assemblages. Hydrol. Processes 15, 2579-2594.

628

- 629 Hughes, L. 2011. Climate change and Australia: key vulnerable regions. *Reg Environ Change* 11
 630 (Suppl 1),S189-195.
- 631
- 632 Iriogen, X., Flynn, K.J., and Harris, R.P. 2005. Phytoplankton blooms: a 'loophole' in
 633 microzooplankton grazing impact? *J. Plank. Res.* 27, 313-321.
- 634
- 635 Kemp, W. M., Boynton, W.R., Adolf, J.E., Boesch, D.F., Boicourt, W.C., Brush, G., Cornwell, J.C.,
 636 Fisher, T.R., Glibert, P.M., Hagy, J.D., Harding, L.W., Houde, E.D., Kimmel, D.G., Miller, W.D.,
 637 Newell, R.I.E., Roman, M.R., Smith, E.M., and Stevenson, J.C. 2005. Eutrophication of
 638 Chesapeake Bay: historical trends and ecological interactions. *Mar. Ecol. Progr. Ser.* 303, 1-29.
- 639
- 640 Kempton, J. W., Lewitus, A.J., Deeds, J.R., Law, J.M., and Place, A.R. 2002. Toxicity of
 641 *Karlodinium micrum* (Dinophyceae) associated with a fish kill in a South Carolina brackish
 642 retention pond. *Harmful Algae* 1, 233-241.
- 643
- 644 Kurup, R. G., and Hamilton, D. P. 2002. Flushing of dense, hypoxic water from a cavity of the
 645 Swan River Estuary, Western Australia. *Estuaries* 25, 908-915.
- 646
- 647 Li, A., Stoecker, D.K., Coats, D.W., and Adam, E.J. 1996. Ingestion of fluorescently labelled and
 648 phycoerythrin-containing prey by mixotrophic dinoflagellates. *Aquat. Microb. Ecol.* 10, 139-147.
- 649
- 650 Li, A., Stoecker, D. K., and Coats, W. D. 2002. Use of the 'food vacuole content' method to estimate
 651 grazing by the mixotrophic dinoflagellate *Gyrodinium galtheanum* on cryptophytes. *J. Plank. Res.*
 652 23, 303-318.

653

654 Litaker, R. W., Vandersea, M. W., Kibler, S. R., Reece, K. S., Stokes, N.A., Steidinger, K. A.,
655 Millie, D. F., Bendis, B. J., Pigg, R. J. & Tester, P. A. 2003. Identification of *Pfiesteria piscicida*
656 (Dinophyceae) and *Pfiesteria*-like organisms using internal transcribed spacer-specific PCR assays.
657 J. Phycol. 39,754–61.

658

659 Mitra, A., and Flynn, K. J. 2006. Promotion of harmful algal blooms by zooplankton predator
660 activity. Biol. Lett. 2, 194-197.

661

662 Mooney, B. D., de Salas, M., Hallegraeff, G. M. and Place, A. R. 2009. Survey of karlotoxin
663 production in 15 species of gymnodinoid dinoflagellates (Kareniaceae, Dinophyta). J. Phycol 45,
664 164-175.

665

666 Mooney, B.D., Hallegraeff, G.M., and Place, A.R. 2010. Ichthyotoxicity of four species of
667 gymnodinoid dinoflagellates (Kareniaceae, Dinophyta) and purified karlotoxins to larval
668 sheepshead minnow. Harmful Algae. 9,557-562.

669

670 Peng, J., Place, A. R., Yoshida, W., Anklin, C., and Hamann, M. T. 2010. Structure and absolute
671 configuration of Karlotoxin, an ichthyotoxin with global distribution from the marine dinoflagellate
672 *Karlodinium veneficum*. J. Am. Chem. Soc. 132 , 3277–3279

673

674 Place, A. R., Bowers, H. A., Bachvaroff, T. R., Adolf, J. E., Deeds, J. E., and Sheng, J. (2012).
675 *Karlodinium veneficum* - The little dinoflagellate with a big bite. Harmful Algae. 14, 179-195.

676

- 677 Robson, B. J., and Hamilton, D.P. 2003. Summer flow event induces a cyanobacterial bloom in a
678 seasonal Western Australian estuary. *Mar. Freshwater Res.* 54, 139-151.
- 679
- 680 de Salas, M. F., Bolch, C. J. S. & Hallegraeff, G. M. 2005. *Karlodinium australe* sp. nov.
681 (Gymnodiniales, Dinophyceae), a new potentially ichthyotoxic unarmoured dinoflagellate from
682 lagoonal habitats of south-eastern Australia. *Phycologia* 44, 640– 50.
- 683
- 684 de Salas, M. F., Laza-Martinez, A. & Hallegraeff, G. M. 2008. Novel unarmored dinoflagellates
685 from the toxigenic family Kareniaceae (Gymnodiniales): five new species of *Karlodinium* and one
686 new *Takayama* from the Australian sector of the Southern Ocean. *J. Phycol.* 44,241–57.
- 687
- 688 Sheng, J., Malkiel, E., Katz, J., Adolf, J. E., and Place, A. R. 2010. A dinoflagellate exploits toxins
689 to immobilize prey prior to ingestion. *Proc. Natl. Acad. Sci. U. S. A.*, 107, 2082-2087.
- 690
- 691 Smayda, T. J. 2008. Complexity in the eutrophication-harmful algal bloom relationship, with
692 comment on the importance of grazing. *Harmful Algae* 8, 140-151.
- 693
- 694 Stephens, R., and Imberger, J. 1996. Dynamics of the Swan River Estuary: the seasonal variability.
695 *Mar. Freshwater Res.* 47, 517-529.
- 696
- 697 Stoecker, D. K., Adolf, J. E., Place, A. R., Glibert, P. M., and Merrit, D.W. 2008. Effects of the
698 dinoflagellates *Karlodinium veneficum* and *Prorocentrum minimum* on early life history stages of
699 the eastern oyster (*Crassostrea virginica*). *Mar. Biol.* 154, 81-91.
- 700

- 701 Sunda, W. G., Graneli, E., and Gobler, C. J. 2006. Positive feedback and the development and
 702 persistence of ecosystem disruptive algal blooms. *J. Phycol.* 42, 963-974.
- 703
- 704 Thompson, P. A., and Hosja, W. 1996. Nutrient limitation of phytoplankton in the upper Swan
 705 River Estuary, Western Australia. *Mar. Freshwater Res.* 47, 659-667.
- 706
- 707 Thompson, P. A. 1998. Spatial and temporal patterns of factors influencing phytoplankton in a salt
 708 wedge estuary, the Swan River, Western Australia. *Estuaries* 21, 801-817.
- 709
- 710 Twomey, L., and John, J. 2001. Effects of rainfall and salt-wedge movement on phytoplankton
 711 succession in the Swan-Canning Estuary, Western Australia. *Hydrological Processes.* 15,2655-
 712 2669.
- 713
- 714 Tyler, M.A. and Seliger, H.H. 1978. Annual subsurface transport of a red tide dinoflagellate to its
 715 bloom area: water circulation patterns and organism distributions in Chesapeake Bay. *Limnol.*
 716 *Oceanogr.* 23,227-246.
- 717
- 718 Van Wagoner, R. M., Deeds, J. R., Satake, M., Ribeiro, A. A., Place, A. R., and Wright, J. L. C.
 719 2008. Isolation and characterization of karlotoxin 1, a new amphipathic toxin from *Karlodinium*
 720 *veneficum*, *Tetrahedron Lett.* 49, 6457-6461.
- 721
- 722 Van Wagoner, R. M., Deeds, J. R., Tatters, A. O., Place, A. R., Tomas, C. R., and Wright, J. L. C.
 723 2010. Structure and relative potency of several karlotoxins from *Karlodinium veneficum*. *J. Nat.*
 724 *Prod.* 73, 1360–1365.

725
726 Wagget, R. J., Tester, P. A., and Place, A. R. 2008. Anti-grazing properties of the toxic
727 dinoflagellate *Karlodinium veneficum* during predator-prey interactions with the copepod *Acartia*
728 *tonsa*. Mar. Ecol. Progr. Ser. 366,31-42.

729
730 Vaque, D., Felipe, J., Sala, M. M., Calbet, A., Estrada, M., Alcaraz, M. 2006. Effects of the toxic
731 dinoflagellate *Karlodinium* sp. (cultured at different N/P) on micro and mesozooplankton. Sci. Mar.
732 70, 59-65.

733

734

735 Table 1. Particulate C, N, and P composition of samples taken in high-density *K. veneficum* waters.

736 DOY = day of year; km = kilometers from mouth of estuary. Site abbreviations are given on Fig. 1.

737

Date (2005)	DOY	Site	km	C:N(mol)	C:P(mol)	<i>K. veneficum</i> (cells mL ⁻¹)
3/31	90	MSB	42	7.5	78.6	12044
3/31	90	WMP	38	7.4	121.6	42420
3/31	90	MEA	36	8.2	191.6	154429
3/31	90	SUC	35	8.8	85.2	116847
3/31	90	KIN	33	8.0	140.6	51712
4/4	94	MSB	42	7.7	89.2	4621
4/4	94	CAV	39	8.2	46.1	29189
4/4	94	WMP	38	7.8	65.9	46864
4/4	94	SUC	35	8.7	66.2	104737
4/4	94	KIN	33	8.7	95.0	79586
4/4	94	BWR 10	32	7.7	97.5	21513
4/4	94	RON	31	7.5	89.9	34239
Avg.				8.0	97.3	
(±sd)				(±0.51)	(±38.80)	

738

739

740

741

Table 2. Cell-specific and water toxin statistics for records with cell abundance >10³ cells mL⁻¹ and detectable toxin levels. n = sample size; s.d. = standard deviation; Min. = minimum; Q1 = first quartile; Q3 = third quartile; Max. = maximum.

	n	Mean	s.d.	Min.	Q1	Median	Q3	Max.
pg KmTx cell ⁻¹	71	4.4	7.48	0.02	1	2.8	4.8	59.1
ng KmTx mL ⁻¹	71	113.6	185.4	0.9	5.1	33.4	162.6	1052.5

Figure captions

Fig. 1 Sampling map of the Swan River Estuary (SRE). Sampling in the SRE was conducted March - July 2005. Individual site labels and distance from the ocean are indicated on the map, as well as the location of the Bennet Brook and Avon River gages.

Fig. 2 Comparisons of an approximately 500 base ITS region between different *K. veneficum* strains. The sequence spans part of ITS1, the 5.8S rRNA and ITS2. There were no differences in the 5.8 S region. The tree contains an unresolved polytomy at the base. Above each branch are (listed in order of the corresponding strains) strain or clone designations, geographic source, and GenBank accession numbers. Where there is a single accession number with more than one strain the sequences are identical. The isolate 'Huon' refers to strain HU01 from the Huon river (Mooney et al. 2009). Filled symbols on branches are insertions, open symbols on branches are mutations.

Fig. 3. Hydrographs (2005) for the Avon River and Bennet Brook, the latter of which has a confluence with the Swan just upstream from station SUC (please see Fig. 1).

Fig. 4. Surface N and P nutrient conditions in the upper SRE. (A) Total N, (B) dissolved nitrate and nitrite, (C) dissolved ammonium, (D) total phosphorus (E) dissolved phosphate. The box represents the time / place of the bloom of *K. veneficum*.

Fig. 5. Salinity, temperature and dissolved oxygen conditions in the upper SRE. The box represents the time / place of the bloom of *K. veneficum*. Sampling stations are represented by the points on the graphs.

Fig 6. Time series plots of *K. veneficum*, planktonic diatoms, and cryptophytes for the period spanning the 2005 *K. veneficum* bloom at three SRE sites, A) SUC, B) KIN, C) RON. The double arrow along the X-axis of (C) indicates the time period of the *K. veneficum* bloom and fish kill. Regression lines and statistics are shown for *K. veneficum* and planktonic diatoms, whereas trends for cryptophytes were not significant.

Fig. 7. Partially-purified KmTx from the SRE, characterized by LC-MS. The thick line is the total ion chromatogram (left axis). The histograms are from the red blood cell hemolysis assay for each 20 s fraction of the LC run. Insets 1, 2, and 3 show the major ions in each of the peaks with hemolytic activity. Inset 1 corresponds to the toxin KmTx^{Perth}2-1, and inset 3 to the toxin KmTx^{Perth}2-2.

Fig. 8. Further purified KmTx (A. 'Perth 1'; B. 'Perth 2') characterization by LC-MS. Solid line is the OD at 235 nm, dashed line is the total ion chromatogram, insets show the major ions associated with peaks.

Fig. 9. Surface (A) Chl *a*, (B) *K. veneficum* cell numbers, and (C) KmTx concentrations in the SRE. The box represents the time / place of the *K. veneficum* bloom. Sampling stations are represented by points.

796 Fig. 10. H&E (hematoxylin and eosin) stained gill arches from a moribund black seabream
797 (*Acanthopagrus butcheri*) collected at the Swan River fishkill on April 10, 2005. Note loss of intra-
798 lamellar spacing and secondary lamellar fusion with extensive hyperplasia of gill epithelial tissue.
799 This pathology is identical to that observed when pure KmTx 2 is exposed to fish the laboratory
800 (Deeds et al. 2006).

801

802 Fig. 11. (A) Scatter plot of log KmTx mL⁻¹ versus log *K. veneficum* mL⁻¹. (B) Time series plot of
803 KmTx (pg cell⁻¹) for stations SUC, KIN and MEA.

804

805 Fig. 12. Ichthyotoxicity assay of purified KmTx from the Swan River Estuary, Perth, Australia
806 against Sheepshead minnow (*Cyprinodon variegatus*). (A) KmTx 'Perth 1' toxin; (B) KmTx 'Perth
807 2' toxin.

808

809 Fig. 13. Composite time-line of events associated with the *K. veneficum* bloom and subsequent fish
810 kill

811

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Phytoplankton abundance (log cells mL⁻¹)

Figure 7

Figure 8

Figure 9

Figure 10

Figure 11

Figure 12

Consolidated Timeline of Events - *K. veneficum* bloom (25 - 42 km region)