Global solutions to regional problems: collecting global expertise to address the problem of harmful cyanobacterial blooms. A Lake Erie case study

George S. Bullerjahn^{1*}, Robert M. McKay¹, Timothy W. Davis², David B. Baker³, Gregory L. Boyer⁴, Lesley V. D'Anglada⁵, Gregory J. Doucette⁶, Jeff C. Ho^{7,8}, Elena G. Irwin⁹, Catherine L. Kling¹⁰, Raphael M. Kudela¹¹, Rainer Kurmayer¹², Anna M. Michalak⁸, Joseph D. Ortiz¹³, Timothy G. Otten¹⁴, Hans W. Paerl¹⁵, Boqiang Qin¹⁶, Brent L. Sohngen⁹, Richard P. Stumpf¹⁷, Petra M. Visser¹⁸ and Steven W. Wilhelm¹⁹.

¹Department of Biological Sciences, Bowling Green State University, Life Sciences Building, Bowling Green, OH 43403 USA

*corresponding author: bullerj@bgsu.edu; +1-419-372-8527

²NOAA-GLERL, 4840 South State Rd., Ann Arbor, MI 48108

³National Center for Water Quality Research, Heidelberg University, Tiffin, OH 44883

⁴Department of Chemistry, SUNY College of Environmental Science and Forestry, 1 Forestry Dr. Syracuse, NY 13210

⁵U.S. Environmental Protection Agency, Office of Science and Technology, 1200 Pennsylvania Ave., N.W. Washington, DC 20460

⁶NOAA/National Ocean Service, 219 Fort Johnson Rd., Charleston, SC 29412

⁷Department of Civil & Environmental Engineering, Stanford University, 473 Via Ortega, Stanford, CA 94305, USA

⁸Department of Global Ecology, Carnegie Institution for Science, 260 Panama St., Stanford, CA 94305 USA

⁹Department of Agricultural, Environmental and Development Economics, Ohio State University, 2120 Fyffe Rd., Columbus, OH 43210

¹⁰Department of Economics, 578 Heady Hall, Ames, IA 50011-2271

¹¹Ocean Sciences, University of California Santa Cruz, 1156 High Street, Santa Cruz, CA 95064

¹²Research Institute for Limnology, University of Innsbruck, Mondseestrasse 9, Modsee, Austria 5310

¹³Department of Geology, 336 McGilvrey Hall, Kent State University, Kent, OH 44242

¹⁴Department of Microbiology, Oregon State University, 226 Nash Hall, Corvallis, OR 97331

¹⁵University of North Carolina Institute of Marine Sciences, 3431 Arendell Street, Morehead City, NC 28557

¹⁶Nanjing Institute of Geography and Limnology, 73 East Beijing Road
Nanjing, China 210008

¹⁷NOAA National Centers for Coastal Ocean Science, 1305 East-West Highway code N/SCI1,
Silver Spring, MD 20910

¹⁸Department of Aquatic Microbiology, Institute for Biodiversity and Ecosystem Dynamics,
University of Amsterdam, P.O. Box 94248, 1090 GE Amsterdam, The Netherlands

¹⁹Department of Microbiology, University of Tennessee, 1414 West Cumberland Avenue,
Knoxville, TN 37996-0845

Keywords: Cyanobacteria, CHAB, Lake Erie, microcystin, cyanotoxin, phosphorus, nitrogen,
eutrophication

Abstract

In early August 2014, the municipality of Toledo, OH (USA) issued a ‘do not drink’ advisory on their water supply directly affecting over 400,000 residential customers and hundreds of businesses (Wilson, 2014). This order was attributable to levels of microcystin, a potent liver toxin, which rose to $2.5 \mu\text{g L}^{-1}$ in finished drinking water. The Toledo crisis afforded an opportunity to bring together scientists from around the world to share ideas regarding factors that contribute to bloom formation and toxigenicity, bloom and toxin detection as well as prevention and remediation of bloom events. These discussions took place at an NSF- and NOAA-sponsored workshop at Bowling Green State University on April 13 and 14, 2015. In all, more than 100 attendees from six countries and 15 US states gathered together to share their perspectives. The purpose of this review is to present the consensus summary of these issues that emerged from discussions at the Workshop. As additional reports in this special issue provide detailed reviews on many major CHAB species, this paper focuses on the general themes common to all blooms, such as bloom detection, modeling, nutrient loading, and strategies to reduce nutrients.

1. Introduction

In early August 2014, the municipality of Toledo, OH (USA) issued a ‘do not drink’ advisory on their water supply directly affecting over 400,000 residential customers and hundreds of businesses (Wilson, 2014). This order was attributable to levels of microcystin, a potent liver toxin, which rose to $2.5 \mu\text{g L}^{-1}$ in finished drinking water and exceeded the $1 \mu\text{g L}^{-1}$ WHO advisory guideline value. This toxic drinking water was caused by a large bloom of cyanobacteria (CHAB: cyanobacterial harmful algal bloom) in the western basin of Lake Erie that was constrained by prevailing winds to the region around the city of Toledo’s water intake. CHABs are not new to Lake Erie; in fact they have recurred annually over the past two decades (Steffen et al., 2014a) and may be occurring with increasing intensity (Michalak et al., 2013; Ho and Michalak, 2015). Whereas this event gained great notoriety as national news in the US, CHABS occur annually on a global scale, promoting chronic and acute health hazards while concurrently producing serious economic effects (Roegner et al., 2014 and references therein). Wherever they are found, the suspected causes of CHABs are typically nutrients arising from various sources including fertilizer, wastewater, the atmosphere, and internal cycling from sediments (Paerl et al., 2011). Recent studies have developed models that predict CHAB intensity in Lake Erie from farm-derived nutrient loads linked to rainfall (Stumpf et al., 2012; Michalak et al., 2013; Obenour et al., 2014). The Toledo crisis afforded an opportunity to bring together scientists from around the world to share ideas regarding factors that contribute to bloom formation and toxigenicity, bloom and toxin detection as well as prevention and remediation of bloom events. These discussions took place at an NSF- and NOAA-sponsored workshop at Bowling Green State University on April 13 and 14, 2015.

The objective of the Workshop was to identify the major knowledge gaps regarding the understanding of bloom formation, detection and mitigation along with discussion of current bloom remediation efforts around the world. The workshop featured NSF- and NOAA-funded researchers and NOAA staff scientists who examined the biology of CHAB species, organism and toxin detection, nutrient management, and bloom forecasting in Lake Erie. Providing the broader international perspectives on bloom mitigation were scientists who have studied CHABs in China, Europe, Australia and Canada. In all, more than 100 attendees from six countries and 15 US states gathered together to share their perspectives. The purpose of this review is to present the consensus summary of the issues that emerged from discussions at the Workshop. Whereas the Workshop and this review focus largely on the 2014 Lake Erie CHAB as a case study, the general conclusions provided here can inform future research aims and mitigation strategies for diverse CHAB events globally. As additional reports in this special issue provide detailed reviews on many major CHAB species, this paper focuses sequentially on the general themes common to all blooms, such as nutrient loading, bloom detection methods, modeling, mitigation strategies, and economic incentives for bloom prevention.

2. Nutrient sources and watershed influences – Lake Erie case study

2.1 Phosphorus loads

Phosphorus (P) is a limiting nutrient widely responsible for controlling algal biomass in freshwater systems (Schindler, 1977). The Maumee River, which drains into the southwestern corner of Lake Erie, is the primary source of the P that is driving CHABs in the lake (Ohio EPA, 2013; US EPA, 2015a). With an area of 17,115 km², its watershed is the largest draining into the Great Lakes. In 1975 the National Center for Water Quality Research at Heidelberg University initiated detailed nutrient-loading studies on the Maumee River at the USGS stream gaging

station near Waterville, OH. Land use upstream from this station, which accounted for 95.8% of the watershed area, consisted primarily of row crop agriculture (73.3%), with smaller portions occupied by forests (6.5%), pasture/hay/grassland (6.3%) and urban areas (10.6%) (USDA Data Gateway for Ohio, 2006). Combined municipal and industrial point source inputs of total P (TP) upstream from Waterville accounted for only about 6.6% of the average annual TP export (Baker et al. 2014a). The TP load from the Maumee River was composed of 25% dissolved reactive P (DRP) and 75% total particulate P (TPP). Since DRP was considered to be 100% bioavailable while TPP in the Maumee River was potentially 25% bioavailable (Baker et al., 2014a), DRP is projected to account for ~56% of the total bioavailable P load based on the estimates at Waterville, OH.

The annual discharge and annual loads and flow-weighted mean concentrations (FWMCs) for the period-of-record (water years 1975-1978 and 1982-2014) at the Waterville station are shown in Fig. 1. The Maumee River had large annual variability in discharge (Fig. 1A) due to variations in weather conditions, including the amounts, intensities and timing of rainfall or snow melt events. Although typically less variable than discharges, the annual FWMCs of nutrients also varied from year to year, due to the interactions of variable weather conditions with seasonal changes in land cover and land management activities (Figs. 1B, D, F, H). Annual nutrient loading, as the product of annual discharge and annual FWMC, was even more variable than discharge or FWMCs (Figs. 1C, E, G, and I). These annual variations in nutrient loading contributed to annual variations in CHAB size in the lake. Comparisons between annual loads of nutrients and annual variations in CHAB size support the development of the load-response

curves that modelers have used to set revised Lake Erie target loads for both TP and DRP (US EPA, 2015; Ohio EPA, 2013; Stumpf et al., 2012).

Detection of trends in annual loading and in FWMCs of nutrients and sediments in rivers is complicated by short-term annual variability. A variety of techniques have been used to visualize and detect trends, including the use of locally weighted scatterplot smoothing (LOWESS) (Richards et al., 2008; Richards et al., 2009), linear regressions of flow adjusted, log transformed data (Richards and Baker, 2002), and running multi-year averages (Baker et al., 2014a; US EPA, 2015). The line graphs shown in Figure 1 represent running 5-year average values with each average calculated and plotted at the mid-point of its 5-year interval. Of particular interest are the changes in discharge, FWMCs and loads that occurred during the re-eutrophication of Lake Erie, beginning in the mid-1990s (Kane et al., 2014; Scavia et al., 2014). To estimate these changes, the 5-yr average value for 1994 was subtracted from that of 2012 and the difference expressed as a percentage of the 1994 value (Fig. 1). The FWMCs of DRP increased by 118% between the 5-year running averages of 1994 and 2012 (Fig. 1B). This increase in concentration, coupled with a 13% increase in discharge, resulted in a 132% increase in DRP loading. This suggests that the increase in DRP loading was primarily due to increases in FWMCs and, to a much lesser extent, to increases in discharge. In contrast with DRP, the FWMCs of TPP decreased by 11% (Fig. 1D). The TPP loads increased by 2% between these time periods due to the increasing discharge. These decreases in TPP FWMCs were attributed to farmer adoption of erosion control best management practices (BMPs) (Richards et al., 2009). Between 1994 and 2012, the TP loads (DRP + TPP) increased by 18%, due to the increase in DRP loading, while total bioavailable P loading increased by 48%, also in response to the

increase in DRP loading. Suspended sediments, which bear the TPP, settled out of the water column rather quickly within the lower Maumee River, Maumee Bay and the lake (Baker et al., 2014b). As such, even the bioavailable portion of TPP becomes less available to support algal growth. In contrast, DRP remains in solution and is accessible to algae and bacteria. Therefore, increases in DRP loading may be even more important as a cause of re-eutrophication than its role in increasing total bioavailable P loading as measured at Waterville (Baker et al., 2014b).

Causes of increased flow-weighted mean concentrations of DRP. Of the multiple factors that may contribute to the increasing DRP loads (Daloglu et al., 2012 and Smith et al., 2015), those factors that are causing the increasing FWMCs appear to be particularly important (Joose and Baker, 2011). The processes by which DRP moves into the water that flows from croplands into streams have been described as either acute or chronic (Kleinman et al., 2011a). Acute losses occur when fertilizers or manure were broadcast onto the surface of fields, and runoff-inducing rainfall or snowmelt occurred shortly thereafter. The most commonly used forms of P fertilizer (mono- and di-ammonium phosphate [MAP and DAP]) are highly soluble and DRP concentrations increase to very high levels in the runoff water (Kleinman et al., 2011b). These losses decrease rapidly with successive runoff events. Chronic losses occurred after P from fertilizers and manures was equilibrated with the soil through either precipitation or adsorption. During runoff events, P moves from surficial soil particles into runoff water as DRP. These surficial soils form a zone of interaction between soil P and the runoff water (Sharpley, 1985). The resulting concentrations in the runoff water were much lower than from acute losses, but tended to be persistent (chronic) since most soils had the capacity to bind large amounts of P relative to that removed by either crops or runoff during a given year. The DRP concentrations in

chronic losses are proportional to P soil test levels in the zone of interaction (Wang et al., 2010, Vadas et al., 2005a; Vadas et al., 2005b, Allen et al., 2006; Williams et al., 2015).

Changing farming practices in northwestern Ohio during the period of re-eutrophication can account for increased DRP losses by both acute and chronic pathways, although the relative importance of each is uncertain (OEPA, 2013). Fertilizer application by broadcasting has increased as farm size has increased, although watershed-specific changes in farm management practices are generally not available. Broadcasting is the least expensive and fastest way to apply fertilizer and it is compatible with modern variable-rate fertilizer applications and precision agriculture. As the number of rows on mechanical planters increased to accommodate planting on larger fields, fertilizer bins used for fertilizer banding were removed from planters. The resulting increased broadcasting of fertilizers increased the opportunities for acute losses of DRP.

Broadcasting of P fertilizers also contribute to increased chronic losses of DRP. Soils have a large binding capacity for P and the P distribution in soils can become very uneven. Surface application of fertilizer, coupled with P release from the breakdown of crop residues at the surface, can result in P stratification wherein surficial soils have higher soil test levels than deeper soils (Sharpley, 2003; Soldat et al., 2007). By adopting practices to reduce erosion and associated losses of TPP, farmers have largely abandoned moldboard plowing that inverted soils and reduced this stratification. They have replaced it with no-till and mulch till operations that can increase the stratification (Kleinman et al., 2015). Consequently, reductions in erosion and TPP losses are often accompanied by increased DRP losses (Kleinman et al., 2015). Soil tests in

the Sandusky River watershed, just to the east of the Maumee Watershed, showed P stratification with P levels in the upper 5 cm (zone of interaction) of the soil column 45% higher than in the 0-20 cm column normally used for agronomic soil testing to support fertilizer management (Fig. 2).

No-till and mulch till production is also accompanied by the development of macropores that provide pathways for surface runoff to move directly to tile drainage systems that lead to streams. The benefits of tile drainage to crop production in this region have led to a constantly increasing density of tile drainage (Kleinman et al., 2015). Tile drainage now constitutes the major pathway by which DRP moves from cropland to streams (Macrae et al., 2007; King et al., 2014; Smith et al., 2014).

Surficial applications of manure also contribute to both acute and chronic losses of DRP. In the Lake Erie Basin, this may be a minor contributor relative to more traditional fertilizer applications where fertilizer applications account for approximately 88% of the P applied to cropland, compared with 12% from manure (Han et al., 2012). Fertilizer application of P exceeds manure P application by a factor of five in the Maumee Watershed. (NRCS, 2009).

2.2 Nitrogen loading and CHABs

Whereas the importance of P loads contributing to CHAB biomass in freshwater has been well established, recent studies of various systems worldwide have also implicated nitrogen (N) as important in supporting both biomass, species composition and toxigenicity (Orr and Jones, 1998; Wilhelm et al., 2003; Van de Waal et al., 2009; Davis et al., 2010; Paerl et al., 2011; Harke

and Gobler, 2013; Paerl et al., 2014; Van de Wall et al., 2014; Müller and Mitrovic, 2015; Gobler et al., *this issue*). Since N in aquatic systems can rapidly cycle through multiple oxidation states, there is often high variability of N species in the water column and sediments. For example, embayments of Lake Erie and Grand Lake St. Marys (OH, USA) often have nitrate levels range from > 100 µM following rainfall events to below detection limits due to assimilation and active denitrification (Steffen et al., 2014; Davis et al., 2015). Additionally, given that microcystin is an abundant secondary metabolite (Nagata et al., 1997), the bioavailability of N likely plays a role in toxin production. Indeed, analysis of the *Microcystis* microcystin synthetase (*mcy*) gene cluster reveals that they are, at least in part, controlled by the N-responsive NtcA transcription factor (Ginn et al., 2010). Recently, western Lake Erie has been shown to exhibit N limitation of chlorophyll *a* production in late summer (Chaffin et al., 2013, Chaffin et al., 2014a, 2014b; Gobler et al., *this issue*), and that the biomass and toxigenicity of the *Planktothrix* bloom in Sandusky Bay, Lake Erie, is seasonally controlled by N inputs (Davis et al., 2015). Indeed, most of the highly aggressive and persistent CHAB blooms now occurring in Lake Erie and worldwide are dominated by non-N₂ fixing genera (*Microcystis* and *Planktothrix*). This sends researchers and managers a strong message externally supplied combined N may contribute to more intense bloom events (Paerl et al., 2014).

The long term trends in N loadings and N:P ratios will be useful in understanding toxin production in future bloom events. Stow et al., (2015) reported a steady decline in TN:TP molar ratios in the Maumee River since the late 1990s, with lowest TN:TP occurring in summertime. Assimilation by biomass and denitrification may explain the documented late summer N limitation in western Lake Erie. Given these trends and the known influences of N on

microcystin gene expression, management of N is warranted along with P (US EPA, 2015b, Steffen et al. 2014a; Gobler et al., *this issue*).

3. Bloom and Toxin Detection

Timely detection of CHABs and their associated toxins is essential from both management and research perspectives. Detection of both organisms *and* toxins is critical, due to variation in intrinsic genetic differences and external environmental factors that may influence gene expression and toxin synthesis. Real- or near real-time surveillance of toxic CHAB events, along with accurate identification and quantification of the organisms and toxins present, are required for efficient and effective mitigation efforts as well as advancing understanding of environmental factors driving population growth and toxicity (Ouellette and Wilhelm 2003). The end result is increased protection of human and animal health, and support of predictive models used to forecast CHAB development and trajectories. Advanced technologies for bloom and toxin detection are becoming commercially available for *in situ* and field-portable deployment as well as for use in the laboratory (Seltenrich, 2014a). These powerful tools are allowing the management and research communities to address critical deficiencies in their ability to rapidly respond to and investigate CHAB events, respectively. In this section, selected technologies and methods are described in the context of potential applications, along with the rationale for adopting a given approach to address a specific management and research need. Several of these are also summarized in Ho and Michalak (2015) and Srivastava et al. (2013).

The noxious (toxins) and nuisance (taste and odor compounds) properties of cyanobacteria are controlled by flexible genes/operons that vary on a strain-by-strain basis (Humbert et al., 2013;

Kurmayer et al., 2015; Simm et al., 2015). No one genus, species, or strain can be reliably considered a producer of any of these compounds without further analysis. The two principal approaches to cyanotoxin risk assessment are direct detection by ELISA, HPLC-UV or LC-MS, or indirect detection by PCR/QPCR of specific genes involved in the biosynthesis of these compounds (Table 1). Both strategies have merits and should not be viewed as mutually-exclusive. For example, many cyanotoxin classes can occur as multiple structural isoforms that are still being discovered (e.g., 6 anatoxins, 3 cylindrospermopsins, 100+ microcystins, 7 nodularins, and 21+ saxitoxins have been reported to date) (Pearson et al., 2010; Cusick and Sayler, 2013), making direct detection of all individual variants by HPLC and LC/MS difficult and expensive. Recent efforts have focused on LC-MS/MS methods that can detect a class of compounds, eliminating the need for individual standards for all of the variants (Foss and Aubel, 2015). Alternatively, ELISA can provide effective detection of some metabolites, including those of cylindrospermopsins and microcystins, as the antibodies employed recognize conserved motifs present across most variants (e.g., James et al., 2010). In other cases, however, such as the paralytic shellfish poisoning toxins (PSTs), these immunoassays react primarily with the parent compound saxitoxin and frequently exhibit low cross-reactivity with other key variants such as neosaxitoxin or the gonyautoxins (e.g., Laycock et al., 2010). Their response to the PST derivatives specific to cyanobacteria (e.g. *Lyngbya* PSTs) is currently unknown. Where they are available, functional assays (i.e., receptor or enzyme inhibition assays) provide an estimate of the integrated toxic potency of a sample regardless of the congeners present and results can correspond closely to values generated by analytical methods (e.g., Powell and Doucette, 1999; Turner et al., 2000). By comparison, PCR assays specifically targeting genes can estimate the potential for a compound to occur and may be useful in the identification of samples that require

further testing by direct analysis using the appropriate method(s) (Rinta-Kanto et al., 2005, 2009; Davis et al., 2009, 2010; Otten et al., 2015). This strategy could eliminate the need to test samples that are likely to be negative, allowing effective implementation of a more cost-effective tiered response approach. Molecular detection methods can amplify the initial target gene over one trillion-fold, making it a robust and sensitive option for early warning surveillance.

Some of the bloom and toxin detection methods identified above are now being deployed on in-water and field-portable platforms, providing greater flexibility in their application along with real- or near real-time data output. Lower cost, widely available sensors for *in situ* physicochemical analyses, including temperature, pH, dissolved oxygen, conductance, and photopigments, can provide data indicative of CHAB formation useful in guiding conventional sampling efforts. Buoy systems are routinely equipped with optical sensors for cyanobacterial pigments and their deployment in areas prone to blooms can provide an early alert for cyanobacterial blooms. A series of fifteen YSI EXO water quality sondes (YSI Inc., Yellow Springs, OH, USA) were deployed in the western basin of Lake Erie and linked by the Great Lakes Observing System (GLOS) network to provide real-time data to water treatment plant operators and the public at-large (habs.glos.us/map/). Outfitting such sensors with both *in vivo* chlorophyll *a* and phycocyanin sensors can permit the clear differentiation of eukaryotic and cyanobacterial algal blooms, as seen in an example from summer 2015 in Sandusky Bay (Fig. 3). Data generated throughout the summer allowed timely adjustment of treatment protocols as blooms migrated near Lake Erie municipal water intakes. Similar systems in the embayments of Lake Ontario allowed recreational users to track the water quality and potential CHABs in waters near their homes and marinas. Instruments such as ISCO water samplers (Teledyne Isco,

Lincoln, NE, USA) can be programmed to collect and preserve water samples autonomously at set times and depths for analyses where real-time results are not essential. Multispectral sensors for algal pigments have been installed in large ships and smaller boats suited for nearshore waters (Pavlac et al., 2012), providing spatial resolution of blooms on the order of tens of meters. This is especially important in the nearshore zone where interference from land reflectance may prevent or compromise the use of satellite imagery. Modern optical sensors such as the Optical Phytoplankton Discriminator (OPD) allow for cyanobacteria blooms to be identified at the genus level (Sullivan and Boyer, unpubl. data). Although the OPD provides very useful information about the causative organisms, determination of toxicity or potential toxicity still requires chemical or molecular testing. Advances are also occurring on this front. For example, the Environmental Sample Processor, or ESP (McLane Research Laboratories, East Falmouth MA, USA), can be deployed subsurface on a mooring and can autonomously determine cell, gene and metabolite (e.g., toxin) concentrations via molecular (sandwich hybridization and/or QPCR; Greenfield et al., 2008; Robidart et al., 2014) and antibody-based (ELISA; Doucette et al., 2009) methods. The ESP has been deployed in marine systems as part of ocean observing networks, providing insights regarding environmental factors influencing bloom dynamics (Ryan et al., 2014) and toxin production. Efforts are now underway to develop ESP applications in freshwaters for CHAB and cyanotoxin detection in Lake Erie at the Toledo, OH water intake (Davis and Doucette, unpubl. data), and to adapt the ESP for deployment on autonomous underwater vehicles for improved spatial coverage (Seltenrich, 2014a). Real-time ESP data could ultimately be assimilated into predictive models for forecasting applications that currently rely primarily on satellite-based observations of surface waters (Stumpf et al., 2012). Although wider application of this technology is currently cost-prohibitive, targeted siting of ESPs along with

selective deployment of autonomous underwater vehicles can alert managers to toxic bloom development and improve sampling efficiency for shore-based testing.

Progress has been made in the development and application of inexpensive field kits for HABs and their toxins. Hand-held, field-portable fluorometers (Algal Torch®, bbe Moldaenke GmbH, Schwentinental, Germany or AquaFluor®, Turner Designs, Sunnyvale, CA, USA) provide scientists with a quick determination if a bloom consists of cyanobacteria and can provide some indication of the level of cyanobacteria associated with the bloom. These tools can be utilized by citizen scientists and water quality managers to evaluate local water quality. Other options for water quality managers include qualitative "dip-stick" assays for cyanotoxins (e.g., Abraxis, Warminster, PA, USA) and field-portable devices for QPCR (e.g., Ubiquitome, Ltd., Dunedin, New Zealand) that could be paired with rapid DNA extraction methods such as Whatman FTA Elute cards (GE Healthcare Life Sciences, Marlborough, MA, USA) for on-site organism and/or target gene detection, producing valuable results in near real-time. While these rapid tools may not yield a definitive answer as to whether a bloom is toxic or not, they can be used to focus sampling and testing efforts and alert managers to potential problems in the early stages of bloom development.

The frequency of discrete sampling needed to detect the occurrence of CHABs should be guided by adaptive management principles, in which water sampling and water quality protocols are adjusted to changes in season. During non-bloom months such as winter and early spring in north temperate ecosystems, it is advisable to conduct occasional (e.g., bi-weekly) visual inspections of the water body and discrete samples should be collected and tested at least monthly and anytime

the water appears discolored. In the spring, as rising water temperatures reach or exceed ~12 °C and lakes begin to stratify, discrete sampling frequency should be increased. Larger bodies of water can leverage remotely sensed satellite or airborne spectroradiometer data to ascertain bloom conditions and geographic location (Section 4 below), whereas autonomous monitoring of smaller water bodies will need to rely on deployment of *in situ* sensors, such as the water quality sondes currently networked in Lake Erie. Devices that are able to detect changes in photopigment concentrations (chlorophyll *a*, phycocyanin) can yield information diagnostic of potential CHAB development. Remote sensing technologies are improving rapidly. Drones fitted with cameras or other imaging devices can already be employed to survey smaller water bodies and relatively inexpensive, low orbiting satellites (e.g., CubeSats) are continually being launched, which will improve access to remotely sensed data by a wider audience (Hand, 2015; Van der Merwe and Price, 2015). Overall, the best strategy to monitor and measure concentrations of CHAB cells and toxins will depend on the management or research application. Specific factors that will drive decision making may include the type of data needed (e.g., qualitative vs. semi-quantitative vs. quantitative), the spatio-temporal coverage required, and how quickly results are desired (e.g., real-time, hours, days) in order to inform effective management decisions or resolve bloom population and toxin dynamics. The focus for those developing future detection technologies continues to be on developing less expensive, higher throughput, and more user-friendly tools that, in some cases, have some degree of field-portability and thus allow for greater flexibility in their application.

4. Bloom prediction and modeling

4.1 Remote sensing tools

413 Remote sensing using satellites and increasingly aircraft and drones provides the capability for
414 rapid synoptic detection of blooms. Remote sensing is particularly effective for CHABs because
415 many of these blooms exhibit surface aggregation. Modeling efforts employ a mixture of
416 statistical and process models to provide hindcasts to understand the mechanisms of HAB
417 formation, nowcasts of near-term changes in conditions or forecasts of potential future CHAB
418 conditions. As remote sensing is limited to observations of the surface of the water (with the
419 signal arising from roughly the uppermost third of the euphotic zone), there are distinct
420 advantages when discrete sampling, subsurface sensors, and remote sensing data are used in
421 conjunction with or assimilated into models, enhancing our ability to quantify errors and thus
422 capitalizing on the strengths and minimizing the weaknesses of each individual component.

423

424 There are currently several CHAB-related reporting products that are regularly available for
425 Lake Erie, which to varying degrees combine results from remote sensing, *in situ* field sampling
426 and model simulations for comparison of observational and simulated fields. These approaches
427 provide products to track map bloom intensity, surface scum distribution, suspended sediment
428 and colored dissolved organic matter distribution and/or simulations of changes in the
429 distribution of particles and environmental properties with time. These systems include the
430 NOAA Lake Erie Experimental HAB Bulletin and associated Experimental HAB Tracker, which
431 compare field, remote sensing and fields from process model simulations (see
432 www.glerl.noaa.gov/res/HABs_and_Hypoxia/ and links therein). The Michigan Tech
433 Developmental Harmful and Nuisance Algal Map product also provides a comprehensive
434 interactive remote sensing atlas including cloud free daily maps of true color imagery, lake

435 surface temperature, HAB extent, chlorophyll *a*, colored dissolved organic matter, dissolved
436 organic carbon and suspended sediment (www.greatlakesremotesensing.org).

437
438 Several major challenges exist when working with remote sensing data. Because pigments are
439 detectable in the visible part of the spectrum, atmospheric biases and cloud cover pose problems
440 which must be addressed. CHAB forming environments have water that is a complex mixture of
441 living and non-living particles and dissolved substances, referred to as Case II water. Case II
442 water can absorb and scatter visible light in ways that are more complex than open ocean
443 environments (Case I water), where traditional band ratio remote sensing methods were
444 developed. Therefore, partitioning the remote sensing response found in conjunction with
445 CHABs and other water constituents is important because the optical complexity of Case II
446 waters (Ortiz et al., 2013, Ali and Ortiz, 2014; Ali et al., 2014b) can produce erroneous
447 chlorophyll *a* biomass estimates when traditional band ratio methods are applied (Witter et al.
448 2009).

449
450 Additional problems arise from differences in instrument design and the variety of algorithms in
451 use, which can make comparison of results from different locations challenging. Differences in
452 spectral resolution and band placement on different instruments can influence the quality of
453 remote sensing products. These vary from sensors such as Landsat and MODIS that have 4-10
454 bands in the visible, to the multi-band option of MERIS, to hyperspectral sensors, such as HICO.
455 MERIS and its intended replacement, the European Space Agency's Sentinel-3 Ocean and Land
456 Color Instrument (OLCI) or proposed hyperspectral instruments such as HypSIRI and PACE
457 offer the greatest capability to detect and quantify cyanobacterial blooms. Overall, instruments

with global missions have been limited by a maximum pixel resolution of 300 m, which precludes their use in many smaller lakes. Landsat (versions 1 through 8 going back over 30 years) and the Sentinel-2 Multispectral Imager (MSI; launched June 2015) offer higher spatial resolution as low as 10 m, but lack frequency of imaging with repeat times on the order of 8-14 days, introducing a challenge in establishing consistency across sensors and availability of data for timely CHAB monitoring and prediction. Whereas Landsat effectively conveyed the intensity of the record summer 2015 CHAB in Lake Erie, its efficacy was limited to a single image representing only a west-central basin swath imaged by the satellite (Fig. 4). Indeed, the low frequency of Landsat overpasses in the lower Laurentian Great Lakes in particular has been an important limitation given frequency of cloud cover over the lakes (*sensu* Vincent et al., 2004). The typical availability of acceptable, cloud-free, summer-time Landsat images around the Lake Erie Islands in the western basin is about 25% due to convective cloud formation associated with thermal differences that arise between the land and water (Ortiz et al. 2013).

The wide variety of algorithms employed by different researchers also often makes intercomparison of results difficult. Remote sensing uses optical characteristics to model biomass (generally as chlorophyll *a*), inherent optical properties (absorption and scattering due to different constituents) or bloom characterization, including the composition of different algal pigment classes. Model specification, parameter settings (without overfitting), and robust validation methods must be identified. Several classes of algorithms have been developed for MERIS, all with particular strengths and weaknesses. There is no systematic way of addressing the use, the confidence or validation of these methods for multiple applications. Sensors such as Landsat, which may provide over 30 years of data, have rarely been applied via a systematic

approach and each new study often creates another “algorithm” or tuning. In some cases attempts have been made to consistently apply multiple algorithms to one sensor (Witter et al., 2009), or one set of algorithms to multiple sensors (Wynne et al., 2013), demonstrating both the potential and the difficulties of employing a single algorithm or suite of algorithms across multiple sensors.

Building off the successful deployment of HICO on the International Space Station, the future can expect increasing use of hyperspectral sensors with higher spatial resolution, first on airborne sensors, but ultimately available from satellite platforms (Seltenrich, 2014b; Kudela et al., 2015). The development of algorithms that fully tap these sensors, thereby furthering bio-optical understanding of how different CHAB constituents can be observed spectrally, is the next challenge in the field (Ortiz et al., 2013; Ali et al., 2014a,b). Hyperspectral data can be used to simulate other sensors, so assimilation of hyperspectral aircraft with MERIS or OLCI should become easier and provides an opportunity to standardize algorithms, uncertainty estimates and validation methods.

Whereas satellites provide both the historical and regional context for CHABs, manned and unmanned aircraft hold the prospect of introducing a key enhancement to satellite data. Aircraft can target small water bodies, provide greater flexibility in deployment and can provide data during cloudy conditions by flying under cloud cover. The effectiveness of an aerial hyperspectral sensing strategy was successfully demonstrated during the summers of 2014 and 2015 by scientists at the NASA Glenn Research Center (Cleveland, OH) and their field collaborators, who conducted weekly overflights of Lake Erie aboard aircraft equipped with a

hyperspectral imager and miniature spectrometers. Flights were coordinated with ground sampling and provided high resolution mapping of the 2014 and 2015 Lake Erie CHAB event. Operations coordinated aerial assets from NASA Glenn and the ONR during 2014 and with NOAA GLERL in 2015. The flexibility offered by aircraft was also key to extending the spatial coverage of the NASA mission to include a >900 km CHAB event that developed in the Ohio River some 300 km south of Lake Erie and which spanned three US states in late August 2015. The combination of satellite, aircraft, sensor, and models will lead to a critical new era of remotely detecting CHABs.

4.2 Future model development

The objectives of modeling are to provide nowcasts and forecasts for management scenarios, and modeling to synthesize what is understood, thereby allowing scientists to capture knowledge gaps. Without validation of the model prediction with independent observations, these objectives cannot be met. Box's maxim that "all models are wrong, but some are useful" (Box and Draper, 1987) must drive the grand strategy for modeling: require clear model objectives and outcomes, robust calibration data for inputs to the models, and independent, quality data for validation of the models.

Currently, a few projects with modeling emphasis have focused on developing alternative loading scenarios for guiding nutrient management. The NOAA-funded "Ecological Forecasting: Hypoxia Assessment in Lake Erie" (EcoFore) Project incorporated field samples, remote sensing and models of varying levels of complexity to study the root causes of hypoxia in the Central Basin of Lake Erie, which largely arises from CHAB events that develop in the western basin of

Lake Erie. EcoFore developed the load response curves needed to guide revised hypoxia-based reduction targets as required by the 2012 Great Lakes Water Quality agreement. Based on these curves, Ecofore recommended phosphorous load reductions to Lake Erie of 46% to reduce HAB occurrence, decrease hypoxia and enhance fish habitat/stocks (Scavia et al. 2014). Similarly, the LimnoTech Western Lake Erie Ecosystem Model (WLEEM; LimnoTech 2010, 2013, 2014; Verhamme et al., 2015), a high-resolution process-based scenario-forecasting model, used field and remote sensing data from 2008 and 2011-2015 for calibration to establish target P loads for reducing cyanobacteria biomass. Based on WLEEM's success in assessing HAB distribution in the western basin and its successful seasonal CHAB forecast for 2015, it is currently being used to assess internal sediment P recycling in the western basin. Other modeling approaches, such as ELCOM-CAEDYM, also provide a coupled hydrodynamic and bio-geochemical modeling strategy that can be used for cyanobacteria (Leon et al., 2011). All these types of models will increasingly be relied on for assimilation of observations to guide nowcast and forecast products.

The success of future CHAB models will depend on a close coupling of models and observations, and the ability to represent processes related to anthropogenic climate forcing. Both statistical and process models can be used to develop and test hypotheses regarding the key factors driving both seasonal and interannual CHAB dynamics. Models will increasingly become assimilative—integrating new data to adjust the forecasts. This is happening now with inter-annual nutrient loads driving forecasts of bloom severity (Stumpf et al., 2012; Obenour et al., 2014) and process models that are calibrated against field and remote sensing data (Verhamme et al. 2015). Effective collaborations between modelers and observationists can also identify the

most important datasets, sampling locations and intervals, ultimately leading to more cost-effective and sustainable monitoring.

Important priorities in developing CHAB models are (1) establishing the end result the models must ultimately predict; (2) identifying the spatio-temporal constraints that impact the end result; (3) adopting best practices for the creation and validation of the model; (4) and identifying the observational methods that will support the development, input, and validation of the model.

The NSF Workshop identified two distinct end results, the modeling of bloom formation and the modeling of toxin forecasting. Including the requirements for modeling into the developing research priorities could lead to better integration between the observational and modeling strategies. For modeling bloom formation, the following actions are needed. First, the objectives of the model need to be defined, including the identification of the bloom biomass metrics, assessing how far in advance forecasts can be made, and the management and scientific needs of forecasts. Second, the models need to test the spatio-temporal occurrence of bloom-triggering events, the seasonality of bloom nutrient requirements, and the relationship between nutrients in the water column and sediments. Third, there is a need for creating substantive, rigorous and robust best practices for model development to provide guidance on a variety of factors, including the avoidance of model overfitting in accuracy assessment, scaling models between ecosystem levels, and integrating results from multiple models. Fourth, there is a need to better integrate simulated model fields with field sampling approaches to optimize sampling design (temporal and spatial acquisition) and methods (what data are best incorporated into models?). Simple sensors, such as pigment fluorescence or turbidity sensors, need to be assessed for their

suitability for CHAB biomass estimation, especially given the general lack of inter-comparability between different metrics demonstrated previously (Ho and Michalak, 2015). Water sampling and sensors need to be integrated into model validation in a way that supports the modeling effort. Key sentinel sites or long-term monitoring sites (nutrients, biomass, etc.) need to be identified and maintained for consistency in model evaluation.

Despite the many challenges presented above regarding model development, useful tools are available for short-term forecasting and generating load reduction targets. With respect to Lake Erie, current efforts integrating remote sensing, regular sampling and particle tracking models have led to the development of HAB Tracker, which provides a 5-day forecast of surface bloom intensity and migration. Future refinement of the HAB Tracker will include validated distributions resulting from migration and wind mixing events that tend to drive *Microcystis* surface scums downward in the water column where municipal water intakes are located. Other groups have coupled different physical (hydrodynamic, wind-wave, sediment transport) and biological (nutrient ingestion, algal growth) models to link nutrient loads to cyanobacteria growth, allowing for the generation of load-reduction scenarios (Verhamme et al. 2015). Visible derivative spectroscopy of hyperspectral field and remote sensing data are also providing increasing specificity for estimation of multiple bloom components, with potential for model assimilation (Ortiz et al., 2013; Ali et al., 2014a,b; Ali and Ortiz, 2014). The challenge for these coupled-models will be to move beyond generating outputs of total biomass toward modeling of algal composition and toxicity.

The first priority for modeling the formation and release of toxins themselves is to develop

models that will determine which toxins will appear and when. These models need to incorporate sufficient detail so that both long- and short-term spatio-temporal variability can be predicted. The models will need to incorporate factors such as grazing pressure and toxin gene expression with changing nutrient and environmental conditions, all of which can be expected to vary through lakes, seasons, years and CHAB species. Validation will push data collection into new areas. It will not be sufficient to simply estimate toxin concentrations from models, as the presence and expression of the genes or toxin-producing strains will be needed to address the uncertainties in the models. Biomolecular measurements from metatranscriptomics (*e.g.* Steffen et al., 2015) will need to be incorporated into the models, as it can provide not only resolution of what organisms are present, but also what each individual species is doing. Until there is a better understanding of why toxins are produced and what affects toxin production (*e.g.* grazing pressure, nutrient availability, oxidative stress [Rohrlack et al., 2001; Ginn et al., 2010; Zilliges et al., 2011]), mechanistic forecasting of toxin production will be unlikely.

Once models are developed, research should extend from predicting toxin biosynthesis within a single organism to focus on predicting spatial “patchiness” and seasonal and species variability in toxin production. Only then can long-term impacts on CHABS brought about by climate change or invasive species be widely understood.

5. Mitigation of blooms and CHAB management

5.1 Best management practices for reducing DRP loading

Best management practices (BMPs) for reducing DRP loading to Lake Erie from cropland can be divided based on their location of application. At the field scale, BMPs can be used to reduce the concentrations of DRP in runoff water or the amounts of runoff water leaving fields, while still

others work by removing DRP from streams and rivers as they transport agricultural runoff to Lake Erie. Some examples of field level BMPs are nutrient management, water management, tillage management, and crop rotations that improve soil health. The optimal combination of these field-level approaches is unique to each field and farm operation.

Nutrient management options are included in the 4-R program of nutrient stewardship: Right product, Right rate, Right time, and Right place (Roberts, 2006; Bruulsema et al., 2009). Losses of DRP associated with direct dissolution of broadcast applications can be immediately reduced by fertilizer placement using injection or banding, and by avoiding broadcast applications at times when heavy rains are forecasted or when ground is frozen or snow covered ground. Where broadcasting is used, incorporation of the fertilizer by light vertical tillage to increase fertilizer-soil contact can minimize the time the broadcast application is vulnerable to direct dissolution. Although such incorporation might increase erosion and particulate P export, the trade-off in terms of reduction of bioavailable P loading would likely yield positive water quality benefits.

Application of the right amount of fertilizer requires information on soil test levels in both the agronomic zone (0-8 inches) and the surficial zone (0-2 inches; Fig. 2). It also requires information on the relationship between soil test levels and crop yields. Since current soil test levels reflect historical past management practices, their reduction can be both challenging and time consuming. Therefore, fertilizer application recommendations have been developed based on agronomic soil test levels and divided into build-up, maintenance, draw-down, and zero application (Lambert et al., 2007). Various guidelines for erosion and particulate P control have

been developed to create awareness of the extent to which they could contribute to increased
DRP runoff to Lake Erie and the effects of that runoff on CHABs (Lambert et al., 2007).

5.2 Cover crops and buffer zones

In most instances, the biggest sources of external N and P loading are diffuse non-point sources, dominated by agricultural and stormwater runoff. The most effective ways to reduce these sources are vegetative riparian buffers, construction of wetlands for treating and removing nutrients, and building of retention ponds for impervious runoff sources (Castelle et al., 1994). These land management strategies have been underutilized in the Great Lakes basin. As an example, concurrent with changes in Lake Erie during the last several decades there have been shifts in a variety of external forcing factors. Large shifts in farm management due to technological change and rising labor costs encouraged a shift from family farms of ~100 acres that dominated the landscape until the 1980s to the large farm operations that now exceed thousands of acres. At the same time, farmers have continued to use fewer crops in crop rotations (e.g., only corn and soybeans in Ohio) and they have changed the timing of fertilizer application in response to economic incentives, including fertilizer prices and labor costs (Huang et al., 1994; Sheriff, 2005). One observation amongst these changes has been the marked decrease in the use of winter cover crops (Fig. 5). Cover crops can provide a series of ecosystem services that historically include both a decrease both erosion and nutrient runoff (Schipanski et al., 2014). Whereas some questions remain concerning how crops in winter months may redistribute specific nutrients within the soil horizon (*i.e.*, moving nutrients from the soil/roots to above ground plant components), their absence has likely lead to significant changes

in the drainage landscape. This problem is in many ways exacerbated by the increasing density of field drainage tiles across the agricultural landscape (Fig. 6).

5.3 Bloom treatment strategies: benefits and limitations

Whereas efforts to prevent blooms must ultimately be directed toward the sources of external nutrient loading, whether they are agricultural, related to urban landscape management or coincident with wastewater treatment, numerous water management strategies have been developed and tested to prevent and mitigate cyanobacterial blooms in lakes and reservoirs. A detailed discussion on mitigation strategies is provided in a separate paper in this volume (Paerl et al., 2016), but a brief summary is provided in this section.

Development of re-established and constructed wetlands offers a geo-engineering strategy to mitigate nutrient loading into lakes to help prevent the occurrence of CHABs (Falconer, 2004). Adopting this strategy to address recurrent seasonal CHABs in Hungary's Lake Balaton, the Kis-Balaton Water Protection System, a mitigation wetland constructed at the mouth of the River Zala, retains more than half of the non-point source nutrients that would otherwise enter the Lake (Tátrai et al., 2000). Combined with sewage diversion projects and geo-political changes resulting in major declines in Hungarian agriculture after 1989, the result has been marked improvement in trophic state for much of the Lake (Hatvani et al., 2014) and declines of formerly abundant CHABs (Istvánovics et al., 2007). Within the Lake Erie watershed, the Great Black Swamp dominated the landscape prior to the draining and reclamation of much of this land for agriculture in the 19th century. The few remnants of the original swamp provide ecosystem services ranging from their use as important stopovers on bird migration flyways to their role in

water quality improvement. Indeed, their efficacy in reducing nutrient loads to Lake Erie is well-documented (Mitsch and Reeder, 1991, 1992 and Rea et al., 2015). More recent wetland restoration projects situated in this region (e.g. Egan et al., 2015) whose construction has been supported by the U.S. Environmental Protection Agency's Great Lakes Restoration Initiative are expected to augment the ecosystem services of the remaining natural wetlands and should help to mitigate the external loading of both P and N into the lake.

Even when external nutrient loading is reduced to aquatic ecosystems, internal recycling of nutrients from the sediments can delay the desirable reduction of phytoplankton biomass (Søndergaard et al., 2003; Paerl et al., 2015). Multiple approaches can be used to reduce the release of P from sediment including dredging (Lürling and Faassen, 2012), hypolimnetic aeration (Beutel and Horne, 1999; Cooke et al., 2005; Singleton and Little, 2006) and chemical sequestration and capping using metal ions such as iron, lanthanum, calcium or aluminum to bind P (Spears et al., 2013; Mackay et al., 2014). In general, costs for aluminum and iron are much lower than other materials, including lanthanum-based products such as Phoslock® (Spears et al., 2013). The Fe-P bond has the disadvantage that P is released upon dissimilatory iron reduction under anaerobic conditions, whereas other ions have stronger bonding with P. The addition of P-binding metal ions is often combined with the addition of flocculants that attach to phytoplankton and aid their settling to the lake bottom. In lakes sufficiently isolated from surrounding water bodies that may otherwise supply the lake with nutrients, chemical sequestration and capping can be effective and keep the lake clear for many years (Akhurst et al., 2004; Lürling and Van Oosterhout, 2013). Reduction of internal N loads are best accomplished through selective enhancement of denitrification, by construction of artificial fringing wetlands

and marshes adjacent to impacted water bodies, and (in reservoirs) optimizing hypolimnetic water discharge to vent the system of excessive hypolimnetic N and P.

A physical technique that may mitigate cyanobacteria is artificial mixing (for a recent review see Visser et al., 2015). Artificial mixing can lead to a shift in phytoplankton composition from cyanobacterial dominance to green algae and diatoms if the imposed mixing is strong enough to keep the cyanobacteria entrained in the turbulent flow (Huisman et al., 2004). The mixing should be deep enough to limit light availability and the mixing devices should be well distributed horizontally over the lake. Mixing techniques are often most effective in relatively small water bodies, where the force of mixing devices can be exerted over a large proportion of the surface area.

Another mitigation technique that works selectively on cyanobacteria in small bodies of water is the addition of low concentrations of hydrogen peroxide. In whole lake experiments, a single treatment with hydrogen peroxide (2 mg L^{-1}) was effective at ridding the lake of cyanobacteria and cyanotoxin problems for an entire season (Matthijs et al., 2012). Other phytoplankton species (green algae, diatoms), zooplankton, macrophytes, birds and fish in the treated lake showed few, if any negative effects. Compared to other cyanocides that are currently in use (e.g. CuSO_4 , see Jancula and Marsalek, 2011), the advantage of hydrogen peroxide is that it rapidly decomposes leaving no permanent biocide traces as is seen with metal toxicants, such as copper salts which can accumulate in the sediments and food chain. Furthermore, cyanotoxins released by lysing cells degrade rapidly in the lake water following peroxide treatment (Matthijs et al., 2012). In the case of lakes too large to be treated as a whole such as Lake Erie, parts of the lake

or secondary reservoirs could be treated with hydrogen peroxide to combat cyanobacterial blooms in areas near drinking water intakes. A ‘water harrow’ on a boat ensures homogeneous addition throughout the lake in very low concentrations (15,000 times dilution from a typical 3% over-the-counter drugstore formulation). The exact dose of hydrogen peroxide needed in the reservoir depends on the phytoplankton composition present and can be accurately tested directly before an application (Weenink et al., 2015). Despite the efficacy of these methods, none of these physical-chemical approaches should be viewed as a substitute for nutrient reductions that should accompany all these protocols.

6. Economic impacts and incentives for bloom prevention and mitigation

6.1 Reducing nutrient loads – what are the costs?

Given that efforts to significantly reduce nutrient loads are often costly, it is important to understand the relative costs of both reducing those loads as well as the costs of taking no action (i.e., the economic impacts of on-going CHABs). A recent estimate, based on the empirical model in Sohngen et al. (2015), suggests that the cost would be about \$30 million per year to reduce P loading within the Lake Erie watershed by 40%. This model utilizes a tax on P inputs used by farmers, which is an economically efficient approach for pollution abatement. The 40% reduction recommended by policy makers could be achieved for about same costs with other approaches, as described below, but a reduction of this scale likely cannot be achieved for less.

What other economic instruments are available to mitigate nutrient loads and reduce the intensity of CHABs? The suite of BMPs employed in Midwestern cropping systems focuses mostly on nutrient management planning, cover crops, conservation rotations, buffer strips and riparian

759 zones. These practices currently require a \$20-\$25 million per year investment from society
760 through USDA conservation programs in the Maumee and Sandusky watersheds. Given the
761 recurrent Lake Erie summertime CHABs, they are insufficient alone as mitigation
762 strategies. Expanding the techniques already adopted in the Lake Erie region to achieve a 40%
763 reduction from today's loads will cost more than is currently expended. If these efforts could be
764 efficiently targeted to known nutrient loading hotspots, it is possible these expenditures could be
765 reduced, but current policies for implementing the USDA conservation programs limit the most
766 effective types of targeting (e.g. costs are not included in criteria for ranking implementation of
767 proposed practices by farmers). The LUMINATE modeling system, which integrates agricultural
768 decision-making in the Mississippi drainage with nutrient loading in the Gulf of Mexico (Kling
769 et al., 2014), offers an assessment model that is relevant for the Lake Erie watershed. In this
770 system, the model analyzes policy scenarios taking into consideration the costs of
771 implementation and the resulting benefits to the receiving body of water.

772
773 Some emerging BMPs such as drainage water management and bioreactors show promise in
774 mitigating CHABs although they are generally quite costly and have not been evaluated in
775 practice. For example, using assumptions of the Ohio Phosphorus Task Force (2013), the cost of
776 blind inlets was found to be ~\$93 per kg P removed. Using the assumptions of the Huron
777 County Soil and Water Conservation District, the cost is nearly \$300 per kg P removed. These
778 costs are about twice the costs of a P tax on a per kg basis. Hence, while drainage water
779 management may be effective, it is costly and must be implemented broadly across the Lake Erie
780 watershed to yield measurable results.

A regulated limit on P inputs to farms could reduce P emissions for the same overall cost as the P tax. Phosphorus limits would likely be more effective and less costly than a P tax because the regulatory system could be designed for both fertilizer and manure. In this case, regulators set a maximum input level, in pounds per acre, for an entire watershed. For instance, if the average input per acre in the Maumee River basin is currently 37 lbs per acre of P_2O_5 , then the regulation can be adjusted downward to a maximum input of 32 lbs per acre, thus achieving a 14% reduction from current levels. This total nutrient input would include manure inputs and chemical inputs. Farmers would need to decide where best to apply their allotted amount of P (*e.g.* 32 lbs in this example multiplied by their total acres). The limit could be set on a watershed-by-watershed basis and could change over time, depending on environmental conditions downstream. Reductions in P would also yield parallel reductions in N as well, so this approach would provide additional value as a general strategy for nutrient management.

While many farmers test their soils for P as a regular protocol for agricultural productivity, a regulated limit on P inputs need not make this a requirement. A farmer's individual P application limit need not be tied to the soil test level on their farm. With a P application limit in place, individual farmers will have a strong incentive to conduct soil tests, but regulators do not need to require it.

A limit on P application can also be amended to denominate P limits for specific crops, or subsidize farmers to eliminate the costs of implementing P reductions. Creating a set of crop-by-crop limits, however, is more burdensome for regulators and it reduces flexibility for farmers to substitute crops that may help achieve P reduction.

Strategies to gain public support of initiatives targeting CHAB prevention and mitigation are also relevant to an economic analysis of this issue. Recently, Egan et al. (2015) reported greater success garnering public support by linking contingent valuation surveys distributed to residents of Ohio's Lake Erie coast concerning proposed wetlands restoration funded by ongoing annual payments compared to a larger one-time payment.

7. Trends promoting future blooms

7.1 CHAB physiology and climate change

Whereas nutrient loadings from agricultural sources are the major drivers of CHAB events in Lake Erie and elsewhere, other trends in chemical and physical factors may favor the future dominance of cyanobacteria. Other changes in the Great Lakes basin may be less tightly linked to human activity and more closely related to changes in climate or biology. Regarding biological changes in the lake, in the late 1980s (and concurrent with the invasion and expansion of the Zebra mussel *Dreissena polymorpha*) there was a systematic decrease in alkalinity (in part due to dissolved inorganic carbonate species) in Lake Erie surface waters (Fig. 7). Concentrations continued to drop in all three basins until 1995, when a slow but steady recovery of alkalinity started in the central and eastern basins. The potential linkage here to cyanobacterial blooms is through the low concentration of dissolved carbon dioxide ($\text{CO}_{2\text{aq}}$), as dense cyanobacterial blooms in low- and moderately alkaline waters can deplete the dissolved CO_2 concentration to limiting levels (Verspagen et al. 2014). Highly buoyant CHAB taxa are able to directly intercept CO_2 at the air-water interface, giving them a competitive advantage under C-limited conditions (Paerl and Ustach, 1982). Moreover, all cyanobacteria and many eukaryotic algae possess efficient carbon concentrating mechanisms (CCMs) that facilitate active

uptake of inorganic carbon species against a concentration gradient of CO₂ (Badger and Price, 2003; Giordano et al., 2005). Previous work has demonstrated that under low CO₂ concentrations, CCMs in cyanobacteria are more efficient than other algae or higher plants (Badger and Price, 2003 and Badger et al., 2006) promoting cyanobacterial dominance under low CO₂ conditions (Price et al., 2008). Recent insights in the genetic diversity of C_i uptake systems of cyanobacteria indicate that there is a large variation in the effectiveness of the cyanobacterial CCMs, even among different strains within the same genus (see Visser et al., 2016 *this issue*). Some cyanobacterial strains perform well at low CO₂, whereas other strains are much better competitors under high CO₂ conditions. As these previous findings are considered in the context of climate warming, increases in CO₂ concentrations may have a more beneficial impact on species that have evolved to not have CCMs, have inferior CCMs, and/or rely primarily on CO₂ transport (Fu et al., 2007). Furthermore, to date, only species-specific responses to rising CO₂ levels have been evaluated for CHAB-forming cyanobacteria. Van de Waal et al. (2011) conducted competition experiments between toxic and non-toxic strains of cyanobacteria at high CO₂ concentrations. This study suggested that non-toxic strains have an advantage under high CO₂ conditions. Complicating this analysis was genetic analysis of several cyanobacterial strains by Sandrini et al. (2014) revealing that the two strains used in the competition experiments by Van de Waal et al. also differed in their C_i uptake systems. The strain with a high-affinity uptake system won the competition at low pCO₂ levels, while the other strain was a C_i uptake generalist (both high and low-affinity uptake C_i systems) and won the competition at high pCO₂ levels. Environmental factors affected by climate change will not occur individually but in parallel with each other. At present, little is known about the interactive effects between increased nutrients, higher temperatures (both predicted to positively influence CHAB formation and toxicity; see

references below) and rising CO₂ concentrations on the development of future CHABs (O’Neil et al., 2012).

Climate change is predicted to result in increased precipitation and more intense storm events, leading to increased nutrient runoff (Michalak et al., 2013). At the same time, changes in surface water temperature are a potential environmental driver of cyanobacterial blooms. Blooms of toxic cyanobacteria are favored by increases in temperature (Paerl and Huisman 2008; Paerl et al. 2010; Paerl et al. 2011). Temperature influences a variety of processes that shape microbial community structure, including absolute growth rates (Paerl and Huisman, 2008 and 2009), gas solubility (Hutchinson, 1957), water column nutrient chemistry (Hutchinson, 1957) and water column structure / mixing processes (Paerl et al., 2011). Several laboratory studies have found that cyanobacteria such as *Microcystis* perform better (i.e. have higher growth rates) at warmer temperatures (see Robarts and Zohary 1987). Furthermore, it was found that toxic strains of *Microcystis* grow faster than non-toxic strains at higher temperatures (Davis et al., 2009). Lake Erie has experienced a small but significant increase in surface waters temperatures (~ 0.05 °C per year over the last two decades) during recent decades (Fig. 8), consistent with global patterns of surface water temperature increases (O’Reilly et al., 2015; Sharma et al., 2015). Perhaps more important, the Great Lakes basin has experienced a period of unprecedented warmth, leading to not only higher seasonal averages, but record extreme temperature observations for both air and water during this period. While at this juncture the relationship between climate change and *Microcystis* blooms in Lake Erie remains an anecdotal observation (as there are not enough scientific data to support this), this is an area of inquiry for scientists. Indeed, changes in climate may influence CHAB distributions throughout the entire Great Lakes: Doblin et al. (2007)

876 demonstrated that *Microcystis* cells could be transported to other Great Lakes in ships' ballast
877 water, and suggested that temperature may be one of the few limiting constraints in Great Lake
878 embayments where blooms have not yet been observed.

879
880

881 **8. Workshop synthesis**

882
883 After two days of presentations and discussions in breakout groups, consensus was reached in
884 the areas listed below. These points were shared with diverse stakeholders and the general
885 public in a NOAA-sponsored open forum at the conclusion of the workshop:

886

8871) **Long term solutions for mitigating CHABs will involve integration of multiple approaches**
888 **into models – from genes to ecosystems.** To fully understand CHAB events and to predict
889 where, when, and how intense and toxic blooms will be, integration of all areas of research from
890 the cellular level to large spatial scales is necessary. There is a large and growing set of
891 approaches that is providing these new data sources, ranging from monitoring cyanobacterial
892 gene expression by transcriptomics to high resolution remote sensing. These emerging data sets
893 must be integrated with more traditional data streams to create both mechanistic and quantitative
894 models to better understand factors promoting the intensity and toxicity of CHABs.

895

8962) **Dual nutrient management is warranted in controlling CHAB events.** Whereas management
897 strategies in freshwaters to date have traditionally focused on controlling P inputs, emerging
898 research from across the globe has provided a wealth of data demonstrating the role N loadings
899 can play in controlling the growth and toxicity of CHABs. As such, N loads should be
900 controlled to mitigate the intensity and toxicity of these events.

901

9023) **Treatment strategies exist for acute bloom control in smaller lakes, but they can be limited**

903 **by scale and by regulatory barriers.** For smaller lakes, multiple methods exist to mitigate

904 bloom events. As an example, the Netherlands validated the performance of dilute hydrogen

905 peroxide as an effective algaecide specific for cyanobacteria. Similar approaches may or may not

906 be suitable in the Great Lakes basin due to individual state regulatory guidelines.

907

9084) **Land management decisions are central in preventing CHAB events.** CHABs are strongly

909 controlled by nutrient loading and within large lake ecosystems, the delivery of nutrients must be

910 restricted to control these events. With respect to the Lake Erie watershed, the use of cover

911 crops, buffer strips and construction of artificial wetlands divert nutrients away from lakes by

912 supporting terrestrial biomass instead of algal biomass. In all cases, mitigating the largest

913 nutrient sources will always be most important for lessening CHAB intensity.

914

9155) **Land use decisions can be incentivized.** If the costs of mitigating CHAB events are

916 understood, economic incentives can be established to encourage agricultural practices that

917 reduce nutrient loadings from farms. Social scientists will play critical roles toward

918 implementing practices and policies for CHAB prevention and mitigation.

919

920 As reviewed within this manuscript, these future research areas will provide information needed

921 by policymakers and stakeholders to prevent of future CHAB events in the Great Lakes and

922 elsewhere in the world. Given that Toledo Mayor D. Michael Collins has testified to the US

923 Senate that the two-day Toledo crisis cost approximately \$2.5 million in lost economic activity,

even a brief event has considerable impact to a community (Congressional Record, December 4, 2014). In this paper, suggestions are presented regarding future research on CHAB events leading to prevention and control of blooms.

Of course, outcomes leading to a decline in bloom events require implementation of new practices, whether they include economic incentives, treatment strategies, management decisions or regulatory legislation. Whatever the outcome, scientific and social science research must inform public policy decisions. Indeed, the Toledo water crisis has proven that fundamental research can lead to sound legislation aimed at protecting the water supply. Important first steps included the passage of Ohio Senate Bill 1, cosponsored by State Sens. Randy Gardner and Rob Peterson, which regulate fertilizer application and disposal of dredged P-rich lake sediments. At the Federal level, H.R. 212, *The Drinking Water Protection Act*, was drafted by Rep. Bob Latta (R, OH), cosponsored by Sens. Rob Portman (R, OH), and Sherrod Brown (D, OH) and signed into law by President Obama. H.R. 212 charges the US EPA to develop a strategic plan to assess and manage risk associated with algal toxins. The fact that new laws emerged the year following the Toledo event demonstrates that government can respond in a timely manner, provided that lawmakers have ready access to the appropriate scientific information. To serve the public effectively, future legislation must continue to rely on effective communication between researchers and elected officials.

Acknowledgements

The CHAB Workshop was supported by the National Science Foundation under grant no. CBET-1515671 (GSB). Additional support was provided by the NOAA Great Lakes Environmental Research Laboratory, the University of Michigan Water Center and by the Office

947 of the Vice President for Research & Economic Development at Bowling Green State
948 University. Additional research support was provided by Ohio Department of Education grant
949 R/HAB-2-BOR to GSB, RMM and JDO. GSB, RMM and SWW thank Mr. Jason Isakovic
950 (Senior Legislative Assistant, office of U.S. Rep. Robert Latta) for helpful discussions on the
951 legislative process. The authors also thank the many staff and students at BGSU who provided
952 assistance prior and during the Workshop.
953

References

- Akhurst, D., Jones, G.B., McConchie, D.M., 2004. The application of sediment capping agents on phosphorus speciation and mobility in a sub-tropical dunal lake. *Marine and Freshwater Research* 55, 715–725.
- Ali, K.A., Ortiz, J.D., 2014. Multivariate approach for chlorophyll-a and suspended matter retrievals in Case II waters using hyperspectral data, *Hydrological Sciences Journal*, 59, 1-14, DOI:10.1080/02626667.2014.964242.
- Ali, K.A., Witter, D.L., Ortiz, J.D., 2014a. Multivariate approach to estimate color producing agents in Case 2 waters using first-derivative spectrophotometer data, *Geocarto International*, 29, 2, 102-127, DOI:10.1080/10106049.2012.743601.
- Ali, K.A., Witter, D.L., Ortiz, J.D., 2014b. Application of empirical and semi-analytical algorithms to MERIS data for estimating chlorophyll a in case waters of Lake Erie, *Environmental Earth Sciences*, 71(9), 4209-4220, DOI 10.1007/s12665-013-2814-0.
- Allen, B.L., Mallarino, A.P., Klatt, J.G., Baker, J.L., Camara, M., 2006. Soil and surface runoff phosphorus relationships for five typical USA midwestern soils. *Journal of Environmental Quality* 35, 599-610.
- Badger, M.R., Price, G.D., 2003. CO₂ concentrating mechanisms in cyanobacteria: molecular components, their diversity and evolution. *Journal of Experimental Botany*, 54, 609-622.
- Badger, M.R., Price, G.D., Long, B.M., Woodger, F.J., 2006. The environmental plasticity and ecological genomics of the cyanobacterial CO₂ concentrating mechanism. *Journal of Experimental Botany*, 57, 249-265.

975 Baker, D.B., Confesor, R., Ewing, D.E., Johnson, L.T., Kramer, J.W., Merryfield, B.J., 2014a.
 976 Phosphorus loading to Lake Erie from the Maumee, Sandusky and Cuyahoga rivers: the
 977 importance of bioavailability. *Journal of Great Lakes Research* 40, 502-517.

978 Baker D.B., Ewing, D.E., Johnson, L.T., Kramer, J.W., Merryfield, B.J., Confesor, R., Richards,
 979 R.O., Roerdink, A.A., 2014b. Lagrangian analysis of the transport and processing of
 980 agricultural runoff in the lower Maumee River and Maumee Bay. *Journal of Great Lakes*
 981 *Research* 40, 479-495.

982 Beutel, M.W., Horne, A.J., 1999. A review of the effects of hypolimnetic oxygenation on lake
 983 and reservoir water quality. *Lake and Reservoir Management* 15: 285-297.

984 Box, G.E.P., Draper, N.R., 1987. *Empirical Model Building and Response Surfaces*, John Wiley
 985 & Sons, New York, NY.

986 Bruulsema, T., Lemunyon, J., Herz, B., 2009. Know your fertilizer rights. *Crops and Soils* 42,
 987 13-18.

988 Castelle, A.J., Johnson, A.W., Conolly, C., 1994. Wetland and stream buffer size requirements—
 989 a review. *Journal of Environmental Quality* 23, 878-882.

990 Chaffin, J. D., Bridgeman, T. B., 2014. Organic and inorganic nitrogen utilization by nitrogen-
 991 stressed cyanobacteria during bloom conditions. *Journal of Applied Phycology* 26,
 992 299–309.

993 Chaffin, J. D., Bridgeman, T. B., Bade, D. L., Mobilian, C. N., 2014. Summer phytoplankton
 994 nutrient limitation in Maumee Bay of Lake Erie during high-flow and low flow years.
 995 *Journal of Great Lakes Research* 40, 524–531.

996 Congressional Record, December 4, 2014. Testimony of Mayor D. Michael Collins to the
 997 Senate Committee on Agriculture, Nutrition and Forestry.

998 Cooke, G.D., Welch, E.B., Peterson, S.A., Nichols, S.A., 2005. Restoration and Management of
 999 Lakes and Reservoirs. 3rd edition. Editor - Cooke, G.D., Taylor and Francis, Boca Raton,
 1000 Florida. 591 p.

1001 Cusick, K.D., Sayler, G.S., 2013. An overview on the marine neurotoxin, saxitoxin: genetics,
 1002 molecular targets, methods of detection and ecological functions. *Marine Drugs* 11, 991-
 1003 1018.

1004 Dalogue, I., Cho, K.H., Scavia, D., 2012. Evaluating causes of trends in long-term dissolved
 1005 reactive phosphorus loads to Lake Erie. *Environmental Science and Technology* 4,
 1006 10660-10666.

1007 Davis, T.W., Berry, D.L., Boyer, G.L., Gobler, C.J., 2009. The effects of temperature and
 1008 nutrients on the growth and dynamics of toxic and non-toxic strains of *Microcystis* during
 1009 cyanobacteria blooms. *Harmful Algae* 8, 715-725.

1010 Davis, T.W., Harke, M.J., Marcoval, M.A., Goleski, J., Orano-Dawson, C., Berry, D.L., Gobler,
 1011 C. J., 2010. Effects of nitrogenous compounds and phosphorus on the growth of toxic and
 1012 non-toxic strains of *Microcystis* during cyanobacterial blooms. *Aquatic Microbial*
 1013 *Ecology* 61, 149–162.

1014 Davis, T.W., Bullerjahn, G.S., Tuttle, T., Rozmarynowycz, M., McKay, R.M., Watson, S.B.,
 1015 2015. Evaluation of increasing nitrogen and phosphorus concentrations on the toxicity
 1016 and chlorophyll content during cyanobacterial blooms in Sandusky Bay, Lake Erie.
 1017 *Environmental Science and Technology* 49, 7197-7207.

1018 Doblin, M.A., Coyne, K.J., Rinta-Kanto, J.M., Wilhelm, S.W., Dobbs, F.C., 2007. Dynamics and
 1019 short-term survival of toxic cyanobacteria species in ballast water from NOBOB vessels
 1020 transiting the Great Lakes-implications for HAB invasions. *Harmful Algae* 6, 519-530.

1021 Doucette, G.J., Mikulski, C.M., Jones, K.L., King, K.L., Greenfield, D.I., Marin, R., Jensen, S.,
 1022 Roman, B., Elliott, C.T., Scholin, C.A., 2009. Remote, subsurface detection of the algal
 1023 toxin domoic acid onboard the Environmental Sample Processor: assay development and
 1024 field trials. *Harmful Algae* 8, 880-888.

1025 Egan, K.J., Corrigan, J.R., Dwyer, D.F., 2015. Three reasons to use annual payments in
 1026 contingent valuation surveys: Convergent validity, discount rates, and mental accounting.
 1027 *Journal of Environmental Economics and Management* 72: 123-136.

1028 Foss, A.J., Aubel, M.T., 2015. Using the MMPB technique to confirm microcystin
 1029 concentrations in water measured by ELISA and HPLC (UV, MS, MS/MS). *Toxicon*
 1030 104, 91-101.

1031 Fu, F.X., Warner, M.E., Zhang, Y., Feng, Y., Hutchins, D.A., 2007. Effects of increased
 1032 temperature and CO₂ on photosynthesis, growth, and elemental ratios in marine
 1033 *Synechococcus* and *Prochlorococcus* (Cyanobacteria). *Journal of Phycology* 43, 485-496.

1034 Ginn, H.P., Pearson, L.A., Neilan, B.A., 2010. NtcA from *Microcystis aeruginosa* PCC 7806 is
 1035 autoregulatory and binds to the microcystin promoter. *Applied and Environmental*
 1036 *Microbiology* 76, 4362-4368.

1037 Giordano, M., Beardall, J., Raven, J.A., 2005. CO₂ concentrating mechanisms in algae:
 1038 mechanisms, environmental modulation, and evolution. *Annual Review of Plant Biology*
 1039 56, 99-131.

1040 Greenfield, D.I., Marin, R., Doucette, G.J., Mikulski, C., Jones, K., Jensen, S., Roman, B.,
 1041 Alvarado, N., Feldman, J., Scholin, C., 2008. Field applications of the second-generation
 1042 Environmental Sample Processor (ESP) for remote detection of harmful algae: 2006-
 1043 2007. *Limnology and Oceanography Methods* 6, 667-679.

1044 Han, H., Allen, B.L., Bosch, N.S., 2012. Historical pattern of phosphorus loading to Lake Erie
1045 watersheds. *Journal of Great Lakes Research* 38, 289-298.

1046 Hand, E., 2015. Startup Liftoff: How flocks of small, cheap satellites, hatched in Silicon Valley,
1047 will constantly monitor a changing Earth. *Science* 348, 172-177.

1048 Harke, M.J., Gobler, C.J., 2013. Global transcriptional responses of the toxic cyanobacterium,
1049 *Microcystis aeruginosa*, to nitrogen stress, phosphorus stress, and growth on organic
1050 matter. *PLoS One*, 8(7), e69834.

1051 Hatvani, I. G., Clement, A., Kovács, J., Kovács, I.S., Korponai, J., 2014. Assessing water-quality
1052 data: The relationship between the water quality amelioration of Lake Balaton and the
1053 construction of its mitigation wetland. *Journal of Great Lakes Research* 40, 115-125.

1054 Ho, J.C., Michalak, A.M., 2015. Challenges in tracking harmful algal blooms: a synthesis of
1055 evidence from Lake Erie. *Journal of Great Lakes Research* 41, 317-325.

1056 Huang, W.-Y., Hansen, L., Uri, N.D., 1994. The application timing of nitrogen fertilizer. *Water,*
1057 *Air, & Soil Pollution* 73, 189-211.

1058 Humbert, J.F., Barbe, V., Latifi, A., Gugger, M., Calteau, A., Coursin, T., Lajus, A., Castelli, V.,
1059 Ozta,s S., Samson, G., Longin, C., Medigue, C., Tandeau de Marsac, N., 2013. A tribute
1060 to disorder in the genome of the bloom-forming freshwater cyanobacterium *Microcystis*
1061 *aeruginosa*. *PLoS ONE* 8, e70747.

1062 Hutchinson, G.E. 1957. *A Treatise on Limnology, Vol. I, Part: 1 Geography and Physics of*
1063 *Lakes.* John Wiley & Sons, New York.

1064 Istvánovics, V., Clement, A., Somlyódy, L., Specziár, A., László, G., Padisák, J., 2007. Updating
1065 water quality targets for shallow Lake Balaton (Hungary), recovering from
1066 eutrophication. *Hydrobiologia* 581, 305-318.

1067 James, R., Gregg, A., Dindal, A., McKernan, J., 2010. Environmental technology verification
 1068 report. Abraxis microcystin test kits. Battelle/US EPA Advanced Monitoring Systems
 1069 Center Technical Report. 73 pages.

1070 Jancula, D., Marsalek, B., 2011. Critical review of actually available chemical compounds for
 1071 prevention and management of cyanobacterial blooms. *Chemosphere* 85, 1415-1422.

1072 Johnson, L.T., 2015. Cumulative Tributary Loading Report Water Year 2014 (in preparation).
 1073 National Center for Water Quality Research, Heidelberg University.
 1074 www.heidelberg.edu/ncwqr.

1075 Joose, P.J., Baker, D.B., 2011. Context for re-evaluating agricultural sources phosphorus
 1076 loadings to the Great Lakes. *Canadian Journal of Soil Science* 91, 317-327.

1077 Kane, D.D., Conroy, J.D., Richards, R.P., Baker, D.B., Culver, D.A., 2014. Re-eutrophication of
 1078 Lake Erie: Correlations between tributary nutrient loads and phytoplankton biomass.
 1079 *Journal Great Lakes Research* 40, 496-501.

1080 King, K.W., Williams, M.R., Fausey, N.R., 2014. Contributions of systematic tile drainage to
 1081 watershed-scale phosphorus transport. *Journal of Environmental Quality* 44, 486-494.

1082 Kleinman, P.J.A., Sharpley, A.N., McDowell, R.W., Flaten, D.N., Buda, A.R., Tao, L.,
 1083 Bergstrom, L., Zhu, Q., 2011a. Managing agricultural phosphorus for water quality
 1084 protection: principals for progress. *Plant Soil* 349, 169-182.

1085 Kleinman, P.J.A., Sharpley, A.N., Buda, A.R., McDowell, R.W., Allen, A.L., 2011b. Soil
 1086 controls of phosphorus in runoff: management barriers and opportunities. *Canadian*
 1087 *Journal of Soil Science* 91, 329-338.

1088 Kleinman P.J.A., Sharpley, A.N., Withers, P.J., Bergström, L., Johnson, L.T., Doody, D.G.,
1089 2015. Implementing agricultural phosphorus science and management to combat
1090 eutrophication. *Ambio* 44 (Supplement 2), S297-S310.

1091 Kling, C.L., Panagopoulos, Y., Rabotyagov, S.S., Valcu, A.M., Gassman, P.W., Campbell, T.,
1092 White, M.J. et al., 2014. LUMINATE: linking agricultural land use, local water quality
1093 and Gulf of Mexico hypoxia. *European Review of Agricultural Economics*, 41, 431-459.

1094 Kosten, S., Huszar, V.L.M., Bécares, E., Costa, L.S., van Donk, E., Hansson, L.A., Jeppesen, E.,
1095 Kruk, C., Lacerot, G., Mazzeo, N., De Meester, L., Moss, B., Lürling, M., Nöges, T.,
1096 Romo, S., Scheffer, M., 2012. Warmer climates boost cyanobacterial dominance in
1097 shallow lakes. *Global Change Biology* 18, 118-126.

1098 Kudela, R. M., Palacios, S. L., Austerberry, D. C., Accorsi, E. K., Guild, L. S., Torres-Perez, J.,
1099 2015. Application of hyperspectral remote sensing to cyanobacterial blooms in inland
1100 waters. *Remote Sensing of Environment* 167, 196-205.

1101 Kurmayer, R., Blom, J.F., Deng, L., Pernthaler, J., 2015. Integrating phylogeny, geographic
1102 niche partitioning and secondary metabolite synthesis in bloom-forming *Planktothrix*.
1103 *The ISME Journal* 9, 909-921.

1104 Laycock, M.V., Donovan, M.A., Easy, D.J., 2010. Sensitivity of lateral flow tests to mixtures of
1105 saxitoxins and applications to shellfish and phytoplankton monitoring. *Toxicon* 55, 597-
1106 605.

1107 Lambert, D.M., Lowenberg-DeBoer, J., Malzer, G., 2007. Managing phosphorus soil dynamics
1108 over space and time. *Agricultural Economics* 37, 43-53.

1109 Leon, L.F., Smith, R.E.H., Hipsey, M.R., Bocaniov, S.A., Higgins, S.N., Hecky, R.E.,
1110 Antenucci, J.P., Imberger, J., Guildford, S.J., 2011. Application of a 3D

1111 hydrodynamic-biological model for seasonal and spatial dynamics of water quality and
 1112 phytoplankton in Lake Erie. *Journal of Great Lakes Research* 37, 41-53.

1113 LimnoTech. 2010. Development, Calibration, and Application of the Lower Maumee River –
 1114 Maumee Bay Model. Technical report prepared for the U.S. Army Corps of Engineers,
 1115 Buffalo District. 113 pp. (December 30, 2010).

1116 LimnoTech. 2013. Development of an Integrated Modeling Approach for Quantifying the GLRI
 1117 Deposition Metric: Pilot Application to Toledo Harbor. Final Technical Report prepared
 1118 for U.S. Army Corps of Engineers – Buffalo District under contract to Ecology &
 1119 Environment, Inc. 125 pp.

1120 LimnoTech and Ecology and Environment. 2014. Influence of Open-Lake Placement of Dredged
 1121 Material on Western Lake Erie Basin Harmful Algal Blooms. Final Technical Report for
 1122 Contract No. W912P4-10-D-0002. Prepared for U.S. Army Corps of Engineers – Buffalo
 1123 District. 166 pp + Appendices.

1124 Lürling, M., Faassen, E.J., 2012. Controlling toxic cyanobacteria: Effects of dredging and
 1125 phosphorus-binding clay on cyanobacteria and microcystins. *Water Research* 46, 1447-
 1126 1459.

1127 Lürling, M., Van Oosterhout, F., 2013. Controlling eutrophication by combined bloom
 1128 precipitation and sediment phosphorus inactivation. *Water Research* 47, 6527-6537.

1129 Mackay, E.B., Maberly, S.C., Pan, G., Reitzel, K., Bruere, A., Corker, N., Douglas, G., et al.,
 1130 2014. Geoengineering in lakes: welcome attraction or fatal distraction? *Inland Waters* 4,
 1131 349-356.

1132 Macrae, M.L., English, M.C. Schiff, S.L., Stone, M., 2007. Intra-annual variability in the
 1133 contribution of tile drains to basin discharge and phosphorus export in a first-order
 1134 agricultural catchment. *Agricultural Water Management* 92, 171-182.

1135 Matthijs, H.C., Visser, P.M., Reeze, B., Meeuse, J., Slot, P.C., Wijn, G., et al., 2012. Selective
 1136 suppression of harmful cyanobacteria in an entire lake with hydrogen peroxide. *Water*
 1137 *Research* 46, 1460–1472.

1138 Michalak, A.M., Anderson, E.J., Beletsky, D., Boland, S., Bosch, N.S., Bridgeman, T.B., et al.,
 1139 2013. Record-setting algal bloom in Lake Erie caused by agricultural and meteorological
 1140 trends consistent with expected future conditions. *Proceedings of the National Academy*
 1141 *of Sciences* 110, 6448-6452.

1142 Mitsch, W.J., Reeder, B.C., 1991. Modelling nutrient retention of a freshwater coastal wetland:
 1143 estimating the roles of primary productivity, sedimentation, resuspension and hydrology.
 1144 *Ecological Modelling* 54, 151-187.

1145 Mitsch, W.J., Reeder, B.C., 1992. Nutrient and hydrologic budgets of a Great Lakes coastal
 1146 freshwater wetland during a drought year. *Wetlands Ecology and Management* 1, 211-
 1147 222.

1148 Müller, S., Mitrovic, M.M., 2015 Phytoplankton co-limitation by nitrogen and phosphorus in a
 1149 shallow reservoir: progressing from the phosphorus limitation paradigm. *Hydrobiologia*
 1150 744, 255–269.

1151 Nagata, S., Tsutsumi, T., Hasegawa, A., Yoshida, F., Ueno, Y., Watanabe, M. F., 1997. Enzyme
 1152 immunoassay for the direct determination of microcystins in environmental water.
 1153 *Journal of AOAC International* 80, 408–417.

1154 Natural Resources Conservation Service, 2009. Rapid Watershed Assessments. Available at

1155 www.oh.nrcs.usda.gov/programs/RWA/rapid_watershed_assessments.html.

1156 Obenour, D.R, Gronewold, A.D., Stow, C.A., and Scavia, D, 2014. Using a Bayesian
 1157 hierarchical model to improve Lake Erie cyanobacteria bloom forecasts. Water Resources
 1158 Research 50, doi:10.1002/2014WR015616.

1159 Ohio Environmental Protection Agency, 2013. Ohio Lake Erie Task Force 2. Final Report.
 1160 Available at <http://www.epa.state.oh.us>.

1161 Ohio Phosphorus Task Force II Final Report, 2013. Available at
 1162 http://lakeerie.ohio.gov/Portals/0/Reports/Task_Force_Report_October_2013.pdf

1163 O’Neil, J.M., Davis, T.W., Burford, M.A., Gobler, C.J., 2012. The rise of harmful cyanobacteria
 1164 blooms: the potential roles of eutrophication and climate change. Harmful Algae 14, 313-
 1165 334.

1166 O’Reilly, C.M., Sharma, S., Gray, D.K., Hampton, S.E., Read, J.S. Rowley, R.J., et al., 2015.
 1167 Rapid and highly variable warming of lake surface waters around the globe. Geophysical
 1168 Research Letters 42, doi:10.1002/2015GL066235.

1169 Orr, P. T., Jones, G. J., 1998. Relationship between microcystin production and cell division
 1170 rates in nitrogen-limited *Microcystis aeruginosa* cultures. Limnology and Oceanography
 1171 43, 1604–1614.

1172 Ortiz, J.D., Witter, D.L., Ali, K.A., Fela, N., Duff, M., and Mills, L., 2013. Evaluating multiple
 1173 color producing agents in Case II waters from Lake Erie, International Journal of Remote
 1174 Sensing 34, 8854-8880.

1175 Otten, T.G., Crosswell, J.R., Mackey, S., Dreher, T.W., 2015. Application of molecular tools for
 1176 microbial source tracking and public health risk assessment of a *Microcystis* bloom
 1177 traversing 300 km of the Klamath River. Harmful Algae 46, 71-81.

1178 Ouellette, A.J.A., Wilhelm, S.W., 2003. Toxic cyanobacteria: the evolving molecular toolbox.
 1179 Frontiers in Ecology and the Environment 7:359-366
 1180 Paerl, H. W., Ustach, J. 1982. Blue-green algal scums: An explanation for their occurrence
 1181 during freshwater blooms. Limnology and Oceanography 27, 212-217.
 1182 Paerl, H.W., Huisman, J., 2008. Blooms like it hot. Science 320, 57-58.
 1183 Paerl, H.W., Huisman, J., 2009. Climate change: a catalyst for global expansion of harmful
 1184 cyanobacterial blooms. Environmental Microbiology Reports 1, 27-37.
 1185 Paerl, H.W., Rossignol, K.L., Hall, S.N., Peierls, B.L., Wetz, M.S., 2010. Phytoplankton
 1186 community indicators of short- and long-term ecological change in the anthropogenically
 1187 and climatically impacted Neuse River Estuary, North Carolina, USA. Estuaries and
 1188 Coasts 33, 485-497.
 1189 Paerl, H.W., Hall, N. S., Calandrino, E.S., 2011. Controlling harmful cyanobacterial blooms in a
 1190 world experiencing anthropogenic and climactic-induced change. Science of the Total
 1191 Environment 409, 1739-1745.
 1192 Paerl, H. W., 2014. Mitigating harmful cyanobacterial blooms in a human-and climatically-
 1193 impacted world. Life 4, 988–1012.
 1194 Paerl, H.W., Gardner, W.S., McCarthy, M.J., Peierls, B.L., Wilhelm, S.W., 2014. New insights
 1195 on old paradigms – shifting thoughts on freshwater eutrophication. Science 346, 175.
 1196 Paerl, H.W., Gardner, W.S., Havens, K.E., Joyner, A.R., McCarthy, M.J., Newell, S.E., Qin, B.,
 1197 Scott, J.T., 2016. Mitigating cyanobacterial harmful algal blooms in aquatic ecosystems
 1198 impacted by climate change and anthropogenic nutrients. Harmful Algae (in press, this
 1199 issue).

1200 Pavlac, M.M., Smith, T.T., Thomas, S.P., Makarewicz, J.C., Lewis, T.W., Edwards, W.J.,
 1201 Pennuto, C.M., Basiliko, C.P., Atkinson, J.F., Boyer, G.L., 2012. Assessment of
 1202 phytoplankton distribution in the nearshore zone using continuous in situ fluorometry.
 1203 Journal of Great Lakes Research 38(suppl 4), 78-84.

1204 Pearson, L., Mihali, T., Moffitt, M., Kellmann, R., Neilan B., 2010. On the chemistry, toxicology
 1205 and genetics of the cyanobacterial toxins, microcystin, nodularin, saxitoxin and
 1206 cylindrospermopsin. Marine Drugs 8, 1650-80.

1207 Powell, C.L., Doucette, G.J., 1999. A receptor binding assay for paralytic shellfish poisoning
 1208 toxins: recent advances and applications. Natural Toxins 7, 393-400.

1209 Price, G.D., Badger, M.R., Woodger, F.J., Long, B. M., 2008. Advances in understanding the
 1210 cyanobacterial CO₂-concentrating-mechanism (CCM): functional components, Ci
 1211 transporters, diversity, genetic regulation and prospects for engineering into plants.
 1212 Journal of Experimental Botany, 59, 1441-1461.

1213 Richards, R.P, Baker, D.B., Crumrine, J.P., 2009. Improved water quality in Ohio tributaries to
 1214 Lake Erie: A consequence of conservation practices. Journal of Soil and Water
 1215 Conservation. 64, 200-211.

1216 Richards, R.P., Baker, D.B., 2002. Trends in Water Quality in LEASEQ Rivers and Streams
 1217 (Northwestern Ohio), 1975-1995. Journal of Environmental Quality 31, 90-96.

1218 Richards R.P., Baker, D.B., Crumrine, J.P., Kramer, J.W., Ewing, D.E., Merryfield, B.J., 2008.
 1219 Thirty-year trends in suspended sediments in seven Lake Erie tributaries. Journal of
 1220 Environmental Quality 37, 1894-1908.

1221 Rinta-Kanto, J.M., Ouellette, A.J.A., Boyer, G.L., Twiss, M.R., Bridgeman, T.B., Wilhelm,
 1222 S.W., 2005. Quantification of toxic *Microcystis* spp. during the 2003 and 2004 blooms in

1223 western Lake Erie using quantitative real-time PCR. Environmental Science and
 1224 Technology 39, 4198-4205.

1225 Rinta-Kanto, J.M., Konopka, E.A., DeBruyn, J.M., Bourbonniere, R.A., Boyer, G.L., Wilhelm,
 1226 S.W., 2009. Lake Erie *Microcystis*: relationship between microcystin production,
 1227 dynamics of genotypes and environmental parameters in a large lake. Harmful Algae 8,
 1228 665-673.

1229 Robarts, R.D., Zohary, T., 1987. Temperature effects on photosynthetic capacity, inspiration and
 1230 growth rates of bloom forming cyanobacteria. New Zealand Journal of Marine and
 1231 Freshwater Research 21, 391-399.

1232 Roberts, T.L., 2007. Right product, right rate, right time and right place... the foundation of best
 1233 management practices for fertilizer. In Fertilizer Best Management Practices, First
 1234 edition, International Fertilizer Industry Association, Paris, France, pp. 29-32.

1235 Robidart, J.C., Church, M.J., Ryan, J.P., Ascani, F., Wilson, S.T., Bombar, D., Marin, R.,
 1236 Richards, K.J., Karl, D.M., Scholin, C.A., Zehr, J.P., 2014. Ecogenomic sensor reveals
 1237 controls on N₂-fixing microorganisms in the North Pacific Ocean. The ISME Journal 8,
 1238 1175-1185.

1239 Roegner, A.F., Brena, B., Gonzalez-Sapienza, G., Puschner, B., 2014. Microcystins in potable
 1240 surface waters: toxic effects and removal strategies. Journal of Applied Toxicology 34,
 1241 441-457.

1242 Rohrlack, T., Dittmann, E., Börner, T., & Christoffersen, K., 2001. Effects of cell-bound
 1243 microcystins on survival and feeding of *Daphnia* spp. Applied and Environmental
 1244 Microbiology 67, 3523-3529.

1245 Ryan, J.P., McManus, M.A., Kudela, R.M., Lara Artigas, M., Bellingham, J.G., Chavez, F.P.,
 1246 Doucette, G., Foley, D., Godin, M., Harvey, J.B.J., Marin, R., Messié, M., Mikulski, C.,
 1247 Pennington, T., Py, F., Rajan, K., Shulman, I., Wang, Z., Zhang, Y., 2014. Boundary
 1248 influences on HAB phytoplankton ecology in a stratification-enhanced upwelling
 1249 shadow. *Deep-Sea Research II* 101, 63-79.

1250 Scavia, D., Allan, J.D., Arend, K.K., Bartell, S., Beletsky, D., Bosch, N.S., Brandt, S.B., Briland,
 1251 R.D., Daloğlu, I., DePinto, J.V., Dolan, D.M., Evans, M.A., Farmer, T.M., Goto, D., Han,
 1252 H., Höök, T.O., Knight, R., Ludsın, S.A., Mason, D., Michalak, A.M., Richards, R.P.,
 1253 Robert, J.J., Rucinski, D.K., Rutherford, E., Schwab, D.J., Sesterhenn, T.M., Zhang, H.,
 1254 Zhou, Y., 2014. Assessing and addressing the re-eutrophication of Lake Erie: Central
 1255 Basin Hypoxia. *J. Great Lakes Research* 40, 226-246.

1256 Schindler, D. W., 1977. Evolution of phosphorus limitation in lakes. *Science* 195, 260-262.

1257 Schipanski, M.E., Barbercheck, M., Douglas, M.R., Finney, D.M., Haider, K., Kaye, J.P.,
 1258 Kemanian, A.R. et al., 2014. A framework for evaluating ecosystem services provided by
 1259 cover crops in agroecosystems. *Agricultural Systems*, 125, 12-22.

1260 Seltenrich, N., 2014a. Keeping tabs on HABs: New tools for detecting, monitoring, and
 1261 preventing harmful algal blooms. *Environmental Health Perspectives* 122, A206-A213.

1262 Seltenrich, N., 2014. Remote-sensing applications for environmental health research.
 1263 *Environmental Health Perspectives*, 122, A268-A275.

1264 Sharma, S., Gray, D.K., Read, J.S., O'Reilly, C.M., Schneider, P., Quadrat, A., et al., 2015. A
 1265 global database of lake surface temperatures collected by *in situ* and satellite methods
 1266 from 1985–2009. *Scientific Data*, 2 doi:10.1038/sdata.2015.8.

1267 Sharpley, A.N., 1985. Depth of surface soil-runoff interaction as affected by rainfall, soil slope
 1268 and management. *Journal of the Soil Science Society of America* 49, 1010-1015.

1269 Sharpley, A.N., 2003. Soil mixing to decrease surface stratification of phosphorus in manured
 1270 soils. *Journal of Environmental Quality* 32, 1375-1384.

1271 Sheriff, G., 2005. Efficient waste? Why farmers over-apply nutrients and the implications for
 1272 policy design. *Applied Economic Perspectives and Policy* 27, 542-557.

1273 Simm, S., Keller, M., Selymes, M., Schleiff, E., 2015. The composition of the global and feature
 1274 specific cyanobacterial core-genomes. *Frontiers in Microbiology* 6, 219.

1275 Singleton, V.L., Little, J.C., 2006. Designing hypolimnetic aeration and oxygenation systems—a
 1276 review. *Environmental Science and Technology* 40, 7512–7520.

1277 Smith, D.R., King, K.W., Johnson, L., Francesconi, W., Richards, P., Baker, D., Sharpley, A.N.,
 1278 2014. Surface runoff and tile drainage transport of phosphorus in the midwestern United
 1279 States. *Journal of Environmental Quality* 44, 495-502.

1280 Smith, D.R., King, K.W., Williams, M.R., 2015. What is causing harmful algal blooms in Lake
 1281 Erie? *Journal of Soil and Water Conservation* 70, 27A-29A.

1282 Sohngen, B., King, K.W., Howard, G., Newton, J., Forster, D.L., 2015. Nutrient prices and
 1283 concentrations in Midwestern agricultural watersheds. *Ecological Economics*. 112, 141-
 1284 149.

1285 Soldat, D.J., Petrovac, A.M., 2007. Soil phosphorus levels and stratification as affected by
 1286 fertilizer and compost applications. *Applied Turfgrass Science* doi:10.1094/ATS-2007-
 1287 0815-01-RS.

1288 Søndergaard, M., Jensen, J.P., Jeppesen, E., 2003. Role of sediment and internal loading of
 1289 phosphorus in shallow lakes. *Hydrobiologia* 506, 135-145.

1290 Spears, B.M., Dudley, B., Reitzel, K., Rydin, E., 2013. Geo-engineering in lakes - a call for
1291 consensus. *Environmental Science & Technology* 47, 3953-3954.

1292 Srivastava, A., Singh, S., Ahn, C.-Y., Oh, H.-M., Asthana, R.K., 2013. Monitoring approaches
1293 for a toxic cyanobacterial bloom. *Environ. Sci. Technol.* 47, 8999–9013.

1294 Steffen, M.M., Belisle, B.S., Watson, S.B., Boyer, G.L., Wilhelm, S.W., 2014a. Review: Status,
1295 causes and consequences of cyanobacterial blooms in Lake Erie. *Journal of Great Lake*
1296 *Research* 40, 215-225.

1297 Steffen, M.M., Zhu, Z., McKay, R.M.L., Wilhelm, S.W., Bullerjahn, G.S., 2014b. Taxonomic
1298 assessment of a toxic cyanobacteria shift during the 2010 bloom in hypereutrophic Grand
1299 Lake St. Marys (Ohio, USA). *Harmful Algae* 33, 12-18.

1300 Steffen, M.M., Belisle, B.S., Watson, S.B., Bourbonniere, R. A., Boyer, G.L., Wilhelm,
1301 S.W., 2015. Metatranscriptomic evidence for co-occurring top-down and bottom-up
1302 controls on toxic cyanobacterial communities. *Applied and Environmental Microbiology*
1303 81, 3268 - 3275.

1304 Stow, C.A., Cha, Y., Johnson, L.T., Confesor, R., and Richards, R.P. 2015. Long-term and
1305 seasonal trend decomposition of Maumee River nutrient inputs into western Lake Erie.
1306 *Environmental Science and Technology* 49, 3392-3400.

1307 Stumpf, R.P., Wynne, T.T., Baker, D.B., Fahnenstiel, G.L., 2012. Interannual variability of
1308 cyanobacterial blooms in Lake Erie. *PLoS ONE* 7, e42444.

1309 Tátrai, I., Mátyás, K., Korponai, J., Paulovits, G., Pomogyi, P., 2000. The role of the Kis-Balaton
1310 Water Protection System in the control of water quality of Lake Balaton. *Ecological*
1311 *Engineering* 16, 73-78.

1312 Tiessen, K.H.D., Elliot, J.A., Yarotski, J., Lobb, D.A., Flaten, D.N., Glozier, N.E., 2010.
 1313 Conventional and conservation tillage: influence on seasonal runoff, sediment, and
 1314 nutrient losses in the Canadian prairies. *Journal of Environmental Quality* 39, 964-980.
 1315 Turner, J.T., Doucette, G.J., Powell, C.L., Kulis, D.M., Keafer, B.A., Anderson, D.M., 2000.
 1316 Accumulation of red tide toxins in larger size fractions of zooplankton assemblages from
 1317 Massachusetts Bay, USA. *Marine Ecology Progress Series* 203, 95-107.
 1318 United States Department of Agriculture, National Agricultural Statistics Service, 2006. USDA
 1319 Data Gateway for Ohio. Available at:
 1320 www.nass.usda.gov/Research_and_Science/Cropland/metadata/metadata_oh06.htm
 1321 United States Environmental Protection Agency, 2015a. Recommended phosphorus loading
 1322 targets for Lake Erie. Annex 4 Objectives and Targets Task Team Final Report to the
 1323 Nutrients Annex Subcommittee, Great Lake Water Quality Agreement of 2012.
 1324 Available as PDF at www2.epa.gov.
 1325 United States Environmental Protection Agency, 2015b. Preventing eutrophication: scientific
 1326 support for dual nutrient criteria. EPA Office of Water, Document 820-S-15-001.
 1327 Washington DC.
 1328 Vadas, P.A., Kleinmann, P.J.A., Sharpley, A.N., Turner, B.L., 2005a. Relating soil phosphorus
 1329 to dissolved phosphorus in runoff: a single extraction coefficient for water quality
 1330 modeling. *Journal of Environmental Quality* 34, 572- 580.
 1331 Vadas, P.A., Mallarino, A.P., McFarland, A., 2005b. The importance of sampling depth when
 1332 testing soils for their potential to supply phosphorus to surface runoff. SERA17. Position
 1333 Paper. <http://sera17.org/publications/>

1334 Vadas, P.A., Good, L.W., Moore, P.A. Jr., Widman, A., 2009. Estimating phosphorus loss in
 1335 runoff from manure and fertilizer for a phosphorus loss quantification tool. *Journal of*
 1336 *Environmental Quality* 38, 1645-1653.

1337 Van de Waal, D. B., Verspagen, J. M. H., Lurling, M., van Donk, E., Visser, P. M., et al., 2009.
 1338 The ecological stoichiometry of toxins produced by harmful cyanobacteria: an
 1339 experimental test of the carbon-nutrient balance hypothesis. *Ecology Letters* 12,
 1340 1326–1335.

1341 Van de Waal, D.B., Verspagen, J.M., Finke, J.F., Vournazou, V., Immers, A.K., Kardinaal,
 1342 W.E.A. et al., 2011. Reversal in competitive dominance of a toxic versus non-toxic
 1343 cyanobacterium in response to rising CO₂. *The ISME Journal*, 5, 1438-1450.

1344 Van de Waal, D. B., Smith, V. H., Declerck, S. A. J., Stam, E. C. M., Elser, J. J., 2014.
 1345 Stoichiometric regulation of phytoplankton toxins. *Ecology Letters* 17, 736–742.

1346 Van der Merwe, D., Price, K.P., 2015. Harmful algal bloom characterization at ultra-high spatial
 1347 and temporal resolution using small unmanned aircraft systems. *Toxins* 7, 1065-1078.

1348 Verhamme, E., DePinto, J.V., Redder, T., Schlea, D., 2015. Western Lake Erie Ecosystem
 1349 Model - connecting phosphorus loads to HABs biomass. International Association for
 1350 Great Lakes Research Annual Conference, Burlington, VT.
 1351 http://www.limno.com/pdfs/2015_IAGLR_Verhamme.pdf

1352 Verspagen, J. M.H., Van de Waal, D.B., Finke, J.F., Visser, P.M., Van Donk, E., Huisman, J.,
 1353 2014b. Rising CO₂ levels will intensify phytoplankton blooms in eutrophic and
 1354 hypertrophic lakes. *PloS One*, 9(8), e104325.

1355 Vincent, R.K., Qin, X., McKay, R.M.L., Miner, J., Czajkowski, K., Savino, J., Bridgeman, T.,
 1356 2004. Phycocyanin detection from LANDSAT TM data for mapping cyanobacterial
 1357 blooms in Lake Erie. *Remote Sensing of the Environment* 89, 381-392.

1358 Visser, P.M., Ibelings, B.W., Bormans, M., Huisman, J., 2015. Artificial mixing to control
 1359 cyanobacterial blooms: a review. *Aquatic Ecology* (Special Issue 'Cyanobacterial
 1360 blooms. Ecology, prevention, mitigation and control.' Eds. Visser, P.M., Ibelings, B.W.,
 1361 Fastner, J. & Bormans, M.) DOI: 10.1007/s10452-015-9537-0

1362 Visser, P.M., Verspagen, J.M.H, Sandrini, G., Stal, L.G., Matthijs, H.C.P., Davis, T.W., Paerl,
 1363 H.W. & Huisman, J. Effects of rising CO₂ and global warming on harmful
 1364 cyanobacteria. *Harmful Algae*, this issue.

1365 Vitosh, M.L., Johnson, J.L., Mengel, D.B., 1995. Tri-State Fertilizer Recommendations for
 1366 Corn, Soybeans, Wheat and Alfalfa. Extension Bulletin E-2567 (Michigan State
 1367 University, The Ohio State University, Purdue University).

1368 Wang, Y.T., Zhang, T.Q., Hu, Q.C., Tan, C.S., O'Halloran, I.P., Drury, C.F., Reid, D.K., Ma,
 1369 B.L., Ball-Coelho, B., Lauson, J.D., Reynolds, W.D., Welacky, T., 2010. Estimating
 1370 dissolved reactive phosphorus concentration in surface runoff water from major Ontario
 1371 soils. *Journal of Environmental Quality* 39, 1771-1781.

1372 Weenink, E.F.J., Luimstra, V.M., Schuurmans, J.M., Van Herk, M.J., Visser, P.M., Matthijs,
 1373 H.C.P., 2015. Combatting cyanobacteria with hydrogen peroxide: a laboratory study on
 1374 the consequences for phytoplankton community and diversity. *Frontiers in Microbiology*
 1375 6, 714. <http://doi.org/10.3389/fmicb.2015.00714>

1376 Wilhelm, S.W., DeBruyn, J.M., Gillor, O., Twiss, M.R., Livingston, K., Bourbonniere, R.A.,
 1377 Pickell, L.D., Trick, C.G., Dean, A.L., R. Michael L. McKay, 2003. Effect of

1378 phosphorus amendments on present day plankton communities in pelagic Lake Erie.
 1379 Aquatic Microbial Ecology 32, 275-285.
 1380 Williams, M.R., King, K.W., Dayton, E., LaBarge, G.A., 2015. Sensitivity analysis of the Ohio
 1381 phosphorus risk index. Transactions of the American Society of Agricultural and
 1382 Biological Engineers 58, 93-102.
 1383 Wilson, E.K., 2014. Danger from microcystins in Toledo water unclear. Chemical & Engineering
 1384 News 92, 9.
 1385 Witter, D., Ortiz, J.D, Palm, S. Heath, R., Budd, J., 2009. Assessing the application of SeaWiFS
 1386 ocean color algorithms to Lake Erie. Journal of Great Lakes Research 35, 361-370.
 1387 Wynne, T. T., Stumpf, R.P, Briggs, T.O., 2013. Comparing MODIS and MERIS spectral shapes
 1388 for cyanobacterial bloom detection. International Journal of Remote Sensing 34, 6668-
 1389 6678.
 1390 Zilliges, Y., Kehr, J.-C., Meissner, S., Ishida, K., Mikkat, S., Hagemann, M., Kaplan, A., Börner,
 1391 T., Dittmann, E., 2011. The cyanobacterial hepatotoxin microcystin binds to proteins and
 1392 increases the fitness of *Microcystis* under oxidative stress conditions. PLoS ONE 6,
 1393 e17615.
 1394
 1395

Figure legends

Figure 1. The relationships between annual discharge (A) and annual flow-weighted mean concentrations and loads of dissolved reactive phosphorus (DRP [B and C]), total particulate phosphorus (TPP [D and E]), Nitrate-N [F and G] and total Kjeldahl nitrogen (TKN [H and I]) for the Maumee River at Waterville, OH. The percent change between 5-year running average values for 1994 and 2012 are shown for each parameter.

Figure 2. A comparison between Mehlich 3 phosphorus soil test levels in the agronomic zone (0-20 cm. cores) and the environmental zone (zone of interactions with runoff water – 0-5 cm. cores) for 1610 fields in the Sandusky River Watershed.

Figure 3. Summer 2015 surface water *in situ* chlorophyll *a* (black) and phycocyanin (gray) profiles expressed as relative fluorescence units (RFU) measured with an EXO2 sonde (YSI Inc.) deployed from a buoy at the city of Sandusky (OH) principal water intake in coastal waters of Lake Erie. Use of a multiparameter sonde resolved an algal bloom in mid-July (arrow) that was not dominated by cyanobacteria, thus guiding water treatment options by city utilities personnel.

Figure 4. Image of CHAB in west-central Lake Erie captured on July 28, 2015 by the Operational Land Imager (OLI) on Landsat 8. The CHAB enveloped the Lake Erie islands and continued into Canadian waters east of Point Pelee. The persistent CHAB in Sandusky Bay is also visible. Landsat imagery courtesy of NASA Goddard Space Flight Center and U.S. Geological Survey.

Figure 5. Annual acreage of winter wheat planted in the Maumee River watershed for counties in the states of A) Ohio B) Indiana and C) Michigan. Data from the United States Department of Agriculture National Agriculture Statistical Service (www.nass.usda.gov, accessed on Feb 4, 2015).

Figure 6. Tile drainage deployment in the USA. Key indicates percent of acreage drained by tile (data from 2012 US Census of Agriculture).

Figure 7. Mean summer alkalinity estimates for the three basins of Lake Erie since 1983. Data demonstrate a drop in CaCO_3 equivalents since the early 1980's that is coincident with the invasion of *D. polymorpha* into the system. While some level of recovery has been observed in the Central and Eastern basins, alkalinity in the western basin remains low. Data from the Great Lakes National Program Office (courtesy. R. Barbiero).

Figure 8. Mean summer temperature in Lake Erie's western basin since 1983. Surface water temperatures for the western basin of Lake Erie have increased $\sim 0.05\text{ }^{\circ}\text{C yr}^{-1}$ ($R^2 = 0.129$, $p = 0.066$).

1438

1439 **Table 1.** Overview of detection methods for assessing cyanobacterial abundance and toxicity.

1440 ‘Prep’ is time required for preparation of a sample(s) for analysis. ‘Analysis’ is time required for

1441 analysis and initial processing of data into a usable form. ‘na’ = not applicable.

1442

Detection method	Location	Target	Prep (h)	Analysis (h)	Samples/Run	Instrument cost (USD)
ELISA	Lab/Field	Toxin class	1	2	40	\$3,000 - \$25,000*
Receptor Assay	Lab	Toxin class	1	4	10 - 20	\$3,000 - \$50,000* [@]
Enzyme Inhibition Assay	Lab	Toxin class	1	2	40	\$3,000 - \$25,000*
HPLC-UV	Lab	Toxin congener	1	0.5	1	\$35,000 - \$70,000
LC-MS/MS	Lab	Toxin congener	1	0.5	1	\$250,000 - \$500,000
PCR/QPCR	Lab/Field	Toxin gene	1	2	40	\$5,000 - \$40,000
Light Microscope	Lab/Field	Cells	0.25	1	1	\$200 - \$20,000
FlowCAM	Lab/Field	Cells	0.5	8	1	\$65,000 - \$95,000
Imaging Flow Cytobot	Field ^{&}	Cells	na	na	Continuous	\$135,000 - \$160,000
Autonomous Underwater Vehicle	Field	Phys-Chem-Pigments	na	na	Continuous	\$50,000 - \$100,000
Environmental Sample Processor	Field ^{&}	Genes & Toxins	0.5 - 2.0 [§]	1	1	\$180,000 - \$300,000 [#]

* same instrument – 96 well plate reader spectrophotometer

[@] for radioreceptor assays a standard or microplate scintillation counter is required

[#] cost range reflects core instrument to fully deployable assembly

[&] pier-based or in-water deployment

[§] prep time depends on sample volume

1443

Figure 1

Figure 2

Figure 3

Chl *a* (RFU)

10
8
6
4
2
0

Jul

Aug

Sep

Oct

Date

14
12
10
8
6
4
2
0

Phycocyanin (RFU)

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8