Many disjoint edges in topological graphs

Andres J. Ruiz-Vargas^{1,2}

Mathgeom-DCG
Ecole Polytechnique Fédérale de Lausanne
Lausanne, Switzerland

Abstract

A *monotone cylindrical* graph is a topological graph drawn on an open cylinder with an infinite vertical axis satisfying the condition that every vertical line intersects every edge at most once. It is called *simple* if any pair of its edges have at most one point in common: an endpoint or a point at which they properly cross. We say that two edges are *disjoint* if they do not intersect. We show that every simple complete monotone cylindrical graph on n vertices contains $\Omega(n^{1-\epsilon})$ pairwise disjoint edges for any $\epsilon > 0$. As a consequence, we show that every simple complete topological graph (drawn in the plane) with n vertices contains $\Omega(n^{\frac{1}{2}-\epsilon})$ pairwise disjoint edges for any $\epsilon > 0$. This improves the previous lower bound of $\Omega(n^{\frac{1}{3}})$ by Suk which was reproved by Fulek and Ruiz-Vargas. We remark that our proof implies a polynomial time algorithm for finding this set of pairwise disjoint edges.

Keywords: Topological graphs, disjoint edges, cylindrical graphs, complete graph, graph drawings.

¹ Research partially supported by Swiss National Science Foundation grant 200020-144531 and Swiss National Science Foundation grant 200021-137574 and by Hungarian Science Foundation EuroGIGA Grant OTKA NN 102029.

² Email: andres.ruizvargas@epfl.ch

1 Introduction

A *topological graph* is a graph drawn on the plane so that its vertices are represented by points and its edges are represented by Jordan arcs connecting the respective endpoints. Moreover, in topological graphs we do not allow overlapping edges or edges passing through a vertex. A *topological graph* is *simple* if every pair of its edges meet at most once, either in a common vertex or at a proper crossing. We use the words “vertex” and “edge” in both contexts, when referring to the elements of an abstract graph and also when referring to their corresponding drawings. A graph is *complete* if there is an edge between every pair of vertices. We say that two edges are *disjoint* if they do not intersect. Throughout this note n denotes the number of vertices in a graph.

By applying a theorem of Erdős and Hajnal [5], every complete n -vertex simple topological graph contains $e^{\Omega(\sqrt{\log n})}$ edges that are either pairwise disjoint or pairwise crossing. However, it was thought [15] that this bound is far from optimal. Fox and Pach [8] showed that there exists a constant $\delta > 0$ such that every complete n -vertex simple topological graph contains $\Omega(n^\delta)$ pairwise crossing edges. In 2003, Pach, Solymosi, and Tóth [15] showed that every complete n -vertex simple topological graph has at least $\Omega(\log^{1/8} n)$ pairwise disjoint edges. This lower bound was later improved by Pach and Tóth [16] to $\Omega(\log n / \log \log n)$. Fox and Sudakov [10] improved this to $\Omega(\log^{1+\epsilon} n)$, where ϵ is a very small constant. Furthermore, the previous two bounds hold for dense simple topological graphs. Pach and Tóth conjectured (see Problem 5, Chapter 9.5 in [3]) that there exists a constant $\delta > 0$ such that every complete n -vertex simple topological graph has at least $\Omega(n^\delta)$ pairwise disjoint edges. Using the existence of a perfect matching with a low stabbing number for set systems with polynomially bounded dual shattered function [4], Suk [17] settled this conjecture by showing that there are always at least $\Omega(n^{1/3})$ pairwise disjoint edges. This was later reproved by Fulek and Ruiz-Vargas [12] using completely different techniques. We are now able to improve the lower bound to $\Omega(n^{1/2-\epsilon})$ for any $\epsilon > 0$.

Theorem 1.1 *A complete simple topological graph on n vertices, which is drawn on the plane, contains $\Omega(n^{\frac{1}{2}-\epsilon})$ pairwise disjoint edges.*

To the best of our knowledge, no sub-linear upper bound is known for this problem.

Algorithmic aspects

Theorem 1.1 gives a lower bound on the size of the largest independent set in the intersection graph of the edges in a complete simple topological graph. Besides the fact that computing the maximum number of pairwise disjoint elements in an arrangement of geometric objects is an old problem in computational geometry, this line of research is also motivated by applications, e.g., in frequency assignment [6], computational cartography [1], or VLSI design [13]. Determining the size of a largest independent set is NP-hard already for intersection graphs of sets of segments in the plane lying in two directions [14], disks [7] and rectangles [2]. For most known cases, efficient algorithms searching for a large independent set in intersection graphs of geometric objects can only approximate the maximum. It is for this reason that a lot of research has been developed for finding such approximations (see [9] for more references).

Our proof of Theorem 1.1 yields an efficient algorithm for finding $\Omega(n^{\frac{1}{2}-\epsilon})$ pairwise disjoint edges in a complete simple topological graph. Because the number of pairwise disjoint edges is at most $\lfloor n/2 \rfloor$ then our algorithm is a $O(n^{\frac{1}{2}+\epsilon})$ -approximation algorithm for the problem of finding the maximal set of pairwise disjoint edges in a complete simple topological graph, which beats the factor of $n^{1-\delta}$ for any small $\delta > 0$ in the inapproximability result for the independence number in general graphs due to Zuckerman [18].

2 Cylindrical drawings of graphs

Let $\mathcal{C}$ be the surface of an infinite open cylinder. Formally, $\mathcal{C} = S^1 \times \mathbb{R}$. We may assume that S^1 is the interval $[0, 1]$ after gluing the point 0 to the point 1. Then, we can characterise each point $p \in \mathcal{C}$ by its coordinates: we use p_x to denote its x -coordinate and p_y to denote its y -coordinate with $0 \leq p_x < 1$ and $p_y \in \mathbb{R}$. A *cylindrical graph* is a graph drawn on $\mathcal{C}$ so that its vertices are represented by points and its edges are represented by Jordan arcs connecting the respective endpoints (this is similar to topological graphs with the only difference that the latter are drawn in the plane). As in topological graphs, cylindrical graphs do not allow overlapping edges or edges passing through a vertex. A *cylindrical graph* is *simple* if every pair of its edges meet at most once either in a common vertex or at a proper crossing.

The straight lines of $\mathcal{C}$ with fixed x -coordinate will be called the vertical lines of $\mathcal{C}$. We let $l_{x=a}$ denote the vertical line with x -coordinate equal to a .

Fig. 1. Three different plane representations of cylindrical graphs.

We say that a curve $\gamma \in \mathcal{C}$ is *x-monotone* if every vertical line intersects γ in at most one point. We say that a cylindrical graph is *monotone* if each of its edges is an *x-monotone* curve and furthermore no pair of vertices have the same *x*-coordinate.

It is easy to work with monotone cylindrical graphs. Let G be a monotone cylindrical graph, then G be easily represented on the plane: we simply cut $\mathcal{C}$ along the line $l_{x=0}$ and embed the resulting surface $[0, 1) \times \mathbb{R}$ in the plane in the natural way. We call this the *plane representation* of $\mathcal{C}$. For a graph G drawn on $\mathcal{C}$ we say that the *plane representation* of G is the drawing of G given by the plane representation of $\mathcal{C}$, note that some edges of G might be cut into two connected components while some edges will remain intact, but with both cases, the edges will stay *x-monotone* curves consisting of either one or two connected pieces, see Figure 1.

It is easy to see that a complete topological graph drawn on the plane with its edges being *x-monotone* curves always contains $\lfloor \frac{n}{2} \rfloor$ pairwise disjoint edges. In [12] it was conjectured that a similar statement is true for complete monotone simple cylindrical graphs, that is that every complete monotone simple cylindrical graph contains at least cn pairwise disjoint edges, for some fixed constant c . We proved this conjecture up to a factor of n^ϵ for any $\epsilon > 0$.

Theorem 2.1 *A complete monotone simple cylindrical graph on n vertices contains $\Omega(n^{1-\epsilon})$ pairwise disjoint edges for any $\epsilon > 0$.*

Theorem 2.1 is our main contribution. Theorem 1.1 follows by combining Theorem 2.1 with the following theorem of R. Fulek.

Theorem 2.2 [11] *Let $c(n')$ denote the maximum number c such that every complete monotone simple cylindrical graph on n' vertices contains a disjoint matching of size at least c . For every complete simple topological graph G with n vertices there exists $\Delta = \Delta(G)$, $0 < \Delta < n$, such that G contains a set of at least $\max\{\frac{n}{\Delta}, c(\Delta)\}$ pairwise disjoint edges.*

Proof of Theorem 1.1

Let $\epsilon > 0$ and G be a simple topological graph with n vertices. By Theorem 2.1 it follows that every complete monotone simple cylindrical graph with n' vertices has a set of $cn'^{1-\epsilon}$ pairwise disjoint edges for some constant $c(\epsilon)$. By Theorem 2.2 there exists a $\Delta = \Delta(G)$ such that G has a set of at least $\max\{\frac{n}{\Delta}, c\Delta^{1-\epsilon}\}$ pairwise disjoint edges. If $\Delta \leq n^{1/2}$ then $n^{1/2-\epsilon} \leq n^{1/2} \leq n/\Delta \leq \max\{\frac{n}{\Delta}, c\Delta^{1-\epsilon}\}$. If, on the other hand, $\Delta > n^{1/2}$ then

$$cn^{1/2-\epsilon} \leq c(n^{1/2})^{1-\epsilon} \leq c\Delta^{1-\epsilon} \leq \max\left\{\frac{n}{\Delta}, c\Delta^{1-\epsilon}\right\}.$$

Hence in both cases, we obtain a set with $\Omega(n^{\frac{1}{2}-\epsilon})$ pairwise disjoint edges. $\square$

The proof of Theorem 2.1, not presented here due to lack of space, is inductive and implicitly provides a polynomial time algorithm. Furthermore, the application of Theorem 2.2 can also be done in polynomial time [11]. Hence we have an implicit polynomial time algorithm for finding a set of $\Omega(n^{1/2-\epsilon})$ pairwise disjoint edges in every simple complete topological graph.

References

- [1] Agarwal, P. K., M. van Kreveld and S. Suri, *Label placement by maximum independent set in rectangles*, Comput. Geom. Theory Appl. **11** (1998), pp. 209–218.
- [2] Asano, T. and H. Imai, *Finding the connected components and a maximum clique of an intersection graph of rectangles in the plane*, J. Algorithms **4** (1983), pp. 310–323.
- [3] Brass, P., W. Moser and J. Pach, “Research Problems in Discrete Geometry,” Springer, New York, 2005.
- [4] Chazelle, B. and E. Welzl, *Quasi-optimal range searching in spaces of finite vc-dimension*, Discrete Comput. Geom. **4** (1989), pp. 467–489.
- [5] Erdős, P. and A. Hajnal, *Ramsey-type theorems*, Discrete Applied Mathematics **25** (1989), pp. 37 – 52.
- [6] Even, G., Z. Lotker, D. Ron and S. Smorodinsky, *Conflict-free colorings of simple geometric regions with applications to frequency assignment in cellular networks*, SIAM J. Comput. **33** (2004), pp. 94–136.

- [7] Fowler, R. J., M. S. Paterson and S. L. Tanimoto, *Optimal packing and covering in the plane are NP-complete*, Information Processing Letters **12** (1981), pp. 133–137.
- [8] Fox, J. and J. Pach, *Coloring K_k -free intersection graphs of geometric objects in the plane*, in: *Proceedings of the 2008 Symposium on Computational Geometry*, ACM, 2008, pp. 346–354.
- [9] Fox, J. and J. Pach, *Computing the independence number of intersection graphs*, in: *Proceedings of the Twenty-Second Annual ACM-SIAM Symposium on Discrete Algorithms*, SIAM, 2011, pp. 1161–1165.
- [10] Fox, J. and B. Sudakov, *Density theorems for bipartite graphs and related ramsey-type results*, Combinatorica **29** (2009), pp. 153–196.
- [11] Fulek, R., *Estimating the number of disjoint edges in simple topological graphs via cylindrical drawings*, SIAM Journal on Discrete Mathematics **28** (2014), pp. 116–121.
- [12] Fulek, R. and A. J. Ruiz-Vargas, *Topological graphs: empty triangles and disjoint matchings*, in: *Proceedings of the twenty-ninth annual symposium on Computational geometry*, ACM, 2013, pp. 259–266.
- [13] Hochbaum, D. S. and W. Maass, *Approximation schemes for covering and packing problems in image processing and VLSI*, J. ACM **32** (1985), pp. 130–136.
- [14] Kratochvíl, J. and J. Nešetřil, *Independent set and clique problems in intersection-defined classes of graphs*, Commentationes Mathematicae Universitatis Carolinae **31** (1990), pp. 85–93.
- [15] Pach, J., J. Solymosi and G. Tóth, *Unavoidable configurations in complete topological graphs*, Discrete & Computational Geometry **30** (2003), pp. 311–320.
- [16] Pach, J. and G. Tóth, *Disjoint edges in topological graphs*, in: *Combinatorial geometry and graph theory*, Springer, 2005 pp. 133–140.
- [17] Suk, A., *Disjoint edges in complete topological graphs*, Discrete and Computational Geometry **49** (2013), pp. 280–286.
- [18] Zuckerman, D., *Linear degree extractors and the inapproximability of max clique and chromatic number*, Theory of Computing **3** (2007), pp. 103–128.