**To Treat or Not to Treat: Diminishing Effectiveness of Emamectin Benzoate Tree
Injections in Ash Trees Heavily Infested by Emerald Ash Borer**

**Charles E. Flower^{1*}, Marie Brikha¹, Jennifer E. Dalton¹, Kathleen S. Knight², Miquel A.
Gonzalez-Meler¹,**

¹ Department of Biological Sciences, University of Illinois at Chicago, SES 3262 (M/C 066), 845
West Taylor Street, Chicago, IL 60607, USA. ² USDA Forest Service, Northern Research
Station; 359 Main Rd; Delaware, OH, 43015, USA

Submitted to *Urban Forestry and Urban Greening*

*corresponding author. Tel.: +1 312 996 5929 fax: +1 312 413 243

E-mail addresses: cflowe3@uic.edu (C.E. Flower), mbrikh2@uic.edu (M. Brikha),
jenniferedalton@gmail.com (J.E. Dalton), mmeler@uic.edu (M.A. Gonzalez-Meler)

Abstract

Emerald ash borer (EAB), a non-native invasive tree-boring beetle, is the primary agent behind the widespread mortality of ash trees (*Fraxinus spp.*) in both natural forests and urban areas of North America. While a variety of insecticide options have been adopted for protection against EAB attacks, little has been reported on the success of insecticide treatments on EAB-infested trees. This is particularly important because EAB is difficult to detect in early stages of its infestation and protective treatment may be undertaken on trees already in decline. Here, we investigated the effectiveness over a four-year period of biennial emamectin benzoate injections in trees exhibiting different initial EAB infestation levels (estimated using visual ash canopy condition metrics) in an ash forest in central Ohio. Results indicate that emamectin benzoate treated trees exhibited less canopy decline relative to non-treated control trees over the course of the experiment. In fact, all untreated trees died over the course of the four-year experiment, while on average treated trees did not exhibit a significant decline. Furthermore, initially healthy and moderately EAB impacted trees treated with insecticide either maintained or improved their canopy condition, while initially heavily EAB impacted trees stabilized, declined slightly, or died. Our results suggest that by using trunk injections of emamectin benzoate, private and public landowners may effectively be able to preserve lightly or moderately EAB infested trees or delay the replacement of ash trees at varying levels of EAB infestation.

Keywords: *Fraxinus*, insecticide, canopy decline, invasive species, urban forestry

Introduction:

Disturbances caused by non-native and native pests and pathogens can cause extensive environmental and economic damage to trees in both urban and natural environments, with the costs of non-native species alone estimated at ~\$138 billion USD annually in the United States (Pimentel et al. 2005). In recent years, the inadvertent introduction of emerald ash borer (*Agrilus planipennis* Fairmaire; EAB) has shed light on the importance of tree protection from such disturbances. Relative to other phytophagous insects, phloem and wood-boring insects like the EAB cause the greatest degree of economic damage in North America (Aukema et al. 2011). In fact, the direct and indirect costs solely associated with EAB may be upwards of \$1.7 billion USD per year (Aukema et al. 2011). Serving as both an economic and aesthetic commodity, the potential loss of ash trees in urban and natural environments represents an increasingly pressing concern for public and private landowners. Trees in urban and natural areas provide a variety of ecosystem services including the removal of carbon dioxide and other airborne pollutants, nutrient retention, water filtration, stormwater management, and microclimate regulation (Costanza et al. 1997; Bolund and Hunhammar 1999; Flower and Gonzalez-Meler 2015). Additionally, homeowners place emotional, aesthetic and other subjective values on trees and neighborhood tree cover has been linked to increased home values (Harris et al. 2013; Sander et al. 2010). As a consequence, EAB-induced ash decline has increased awareness and environmental stewardship in human populations surrounded by landscapes experiencing extensive ash mortality (Hunter 2011). Maintaining the health of susceptible tree species vulnerable to species-specific pests and pathogens using insecticides can promote tree diversity in urban areas and may contribute to increased forest resilience (Peterson et al. 1998). Effective insecticides provide an alternative tool to homeowners and local communities to manage the health of their tree populations (McKenney and Pedlar 2012).

Emerald ash borer is a phloem feeding beetle native to Asia, which was accidentally introduced into North America in the 1990s and has since spread to two Canadian provinces and 25 states across the United States resulting in the widespread decline in ash trees (Siegert et al. 2007; Wang et al. 2010; Flower et al. 2013a). The serpentine galleries created in the cambium of ash trees by EAB larvae sever conductive tissue between roots and shoots resulting in tree mortality in as little as two years, although typically mortality occurs in 2-5 years (Flower et al.

2013b). At the tree level, the presence of EAB can be identified by characteristic D-shaped exit holes, woodpecker holes, epicormic growth, and canopy decline (Cappaert et al. 2005; Smith 2006; Flower et al. 2013b; 2014). In EAB-infested regions, tree-level canopy health condition (or tree decline) is well correlated with tree-level EAB densities (Flower et al. 2013b), however ash trees with low EAB densities seldom exhibit symptoms, making them difficult to distinguish from non-infested trees and treat with insecticides before infestation takes place.

In the conterminous United States, urban counties (metropolitan areas) contain over 74.4 billion trees which cover ~33 percent of these areas (Dwyer et al. 2000). Many of these urban and residential trees are ash trees (Kovacs et al. 2010), which were intentionally planted due to their tolerance of high pH, salinity, drought and compacted soils (MacFarlane and Meyer 2005). Prior to the EAB outbreak in Chicago, for example, ash trees ranked as the second most common species across the Cook and DuPage County study area (19.4 % of trees) (Nowak 1994). The widespread distribution of ash trees and their high abundance in urban areas highlights the magnitude of potential economic, environmental, and emotional costs associated with EAB induced ash mortality. In light of the EAB outbreak, landowners and managers in urban and residential areas have three options regarding tree management: 1) repeatedly treat infested tree with insecticide, 2) remove and perhaps replace the infested tree with a non-ash species, and 3) allow the tree to succumb to EAB and remain in place. While a variety of effective insecticide treatments have been deployed to maintain tree health during an EAB outbreak (Table 1), it is commonly thought that because of treatment costs and mixed success, trees with declining canopies should not be treated (Herms et al. 2014; McKenney and Pedlar 2012). However, as long as plant vasculature remains intact insecticide treatment has the potential to slow the senescence of trees in later stages of decline and may reduce or delay capital expenditures associated with removal and replacement. Furthermore, due to the risks to people and property in urban and residential areas, leaving standing dead trees across the landscape is not plausible. If effective, insecticide treatment of infested trees can provide alternatives to managers and citizens by delaying or slowing ash mortality in areas of interest.

The objectives of this research were to investigate the response of EAB infested ash trees in different stages of EAB infestation to biennial emamectin benzoate insecticide injections over four years from 2010-2013. Information regarding the survival of emamectin benzoate injected

ash trees across different canopy condition classes will allow for informed cost-effective decisions based on the value placed at the individual tree or stand level by the local communities.

Methods:

Site description

This study was initiated in May 2010 in an ash plantation with a silty loam soil characterization located at the United States Department of Agriculture (USDA) Forest Service Northern Research Station near Delaware, Ohio (40° 23' N, 83° 02' W). The plantation was established on an agricultural field and subsequently left unmanaged. At the start of the experiment canopy trees were on average ~35 years old (determined dendrochronologically). The forest is at an elevation of 290 m above sea level and its average precipitation from 2006-2010 was 92.3 cm per year. Upon initial inspection of the forest in 2010, the presence of D-shaped exit holes on ash trees (> 80% of ash trees exhibiting exit holes) and the positive trapping of EAB with purple panel traps hung in the canopy (28 EAB captured from May-August 2010) indicated that EAB was present. The forest canopy is dominated by *Fraxinus americana* L. (white ash) and *F. pennsylvanica* Marshall (green ash), which together represent ~ 80% of tree basal area, as well as *Ulmus americana* L. (American elm), *Tilia americana* L. (basswood), and *Morus rubra* L. (mulberry). The understory is dominated by *Lonicera mackii* (amur honeysuckle), *Multiflora rosea* (multiflora rose), *Alliaria petiolata* (garlic mustard), *Poa* spp. and saplings of the canopy species.

The experimental plots were located > 10 m from the edge of the forest to reduce edge effects. The forest was divided into 12 m x 12 m plots and insecticide (n = 4 plots, containing 32 ash trees) and control (n = 3, containing 21 ash trees) treatments were randomly assigned to plots. All ash trees > 7 cm diameter at breast height (DBH) were measured. Pre-treatment DBH (control = 16.43 ± 1.37 and insecticide = 17.21 ± 1.28 ; mean cm $\pm$ SE) and ash canopy condition (AC; control = 2.23 ± 0.22 and insecticide = 1.94 ± 0.18) were not different between control and insecticide treated plots (ttest; $P >> 0.05$). Ash trees > 7 cm DBH residing in the insecticide treatment plots were treated in the spring (May 2010 and 2012) with trunk injections of emamectin benzoate (TREE-ägeTM, Arborjet Inc, Woburn, MA, USA) in quantities recommended by the manufacturer (0.2 g active ingredient for every 2.5 cm DBH). Injection

plugs (Arborplug # 4 for trees < 10 cm DBH and # 5 for trees > 10 cm DBH) were installed evenly around the base every 5 cm at a height of no more than 30 cm from the soil. Areas exhibiting decay were avoided. Injections were conducted using the Arborjet Quik-Jet system (Arborjet Inc.). Ash trees within the control plots were not exposed to any experimental manipulation, but were exposed to moderate natural EAB densities.

Emamectin benzoate

Emamectin benzoate is a systemic insecticide from the avermectin family of macrocyclic lactones which acts as a chlorine channel agonist (Lasota and Dybas 1991). This insecticide was selected for the study as it has been shown to be highly effective at tree level protection from bark beetles, including mountain pine beetle, spruce beetle and western pine beetle (Herms et al. 2014; Grosman et al. 2009; Smitley et al. 2010; Fettig et al. 2013). Emamectin benzoate is thought to provide two levels of protection against EAB. First, larvae will be exposed if compounds diffuse from the xylem to the phloem or if larvae penetrate the xylem during feeding. Second, as emamectin benzoate is transported in the xylem to the canopy adult EAB feeding on leaf tissue will be exposed. A study by McCullough et al. (2011) found emamectin benzoate trunk injections reduced EAB larval densities by almost 100% in treated trees compared to non-treated trees and resulted in near 100% mortality of adults in bioassay feeding trials. Additionally, a study by Herms (2011) indicated that while emamectin benzoate injections provided effective control for 2 years (< 10% decline) at several application rates, efficacy past 2 years was less successful. Despite these findings, it remains unknown whether biennial emamectin benzoate application can provide control to infested trees over a wide range of tree diameters and canopy condition classes for extended time periods (> 2 years), particularly those in later stages of decline > 50% (AC4).

Ash canopy condition monitoring

Annually (2010-2013), following leaf expansion, each ash tree from the study was evaluated for EAB infestation using the visual assessment of canopy health called ash canopy condition (AC) developed by the USDA (Knight et al. 2014; adopted in part from Ball and Simmons 1980; Figure 1). Ash canopy condition exhibits a significant relationship with tree-level EAB densities (larvae/adult) and EAB gallery cover (determined via quantification of EAB

larval gallery cover in 2-100cm² grids on the north and south side of 1m long debarked logs) representing a visual proxy for EAB infestation levels in infested regions not exposed to ash yellows or other tree stress (Flower et al. 2011; 2013b). Briefly, AC is graded on a 1 (healthy) to 5 (dead) scale: 1) healthy/full canopy – a healthy ash canopy is full and exhibits no defoliation; 2) thinning canopy – slight reduction in leaf area, all top branches exposed to sunlight have leaves; 3) dieback – canopy is thinning and some top branches exposed to sunlight are defoliated, lower branches which exhibit natural thinning are not considered; 4) > 50 % dieback – the canopy has less than 50 % of the leaves of a class 1 tree and/or over half of the top branches are defoliated; and 5) dead canopy – no leaves remain in the canopy portion of the tree, regardless of epicormic sprouts (Knight et al. 2014; Figure 1). This canopy condition rating system was utilized in contrast to the 10 % canopy decline system used by Smitley et al. 2008, because it was developed for easy visual canopy classification by untrained individuals (such as homeowners and land managers) and thus minimizes observer bias (in excess of 20 % in the middle range in the Smitley scheme). Furthermore, the proposed scheme is established and in widespread use by the USDA Forest Service and has been linked to EAB densities and backed by neutral tree physiological characteristics such as water use, sapflux, and foliar carbon isotope ratios (Flower et al. 2011; 2013b; Knight et al. 2014). Therefore our scheme and the results of the study are directly applicable to practitioners, planners and the scientific community.

Statistical analyses

To determine whether pre-treatment mean tree diameter (DBH) or initial ash canopy condition (AC) differed between ash trees in the control and insecticide treatment groups, we used two-tailed t-tests ($P < 0.05$) with homogeneity of variance between treatments. No significant differences were observed between treatments for DBH or initial AC ($P = 0.579$ and $P = 0.205$ respectively) permitting further analysis between treatments. To test for differences in the change in ash tree canopy condition (ΔAC ; $AC_{2010} - AC_{2013}$) between treatments, we used an analysis of covariance (ANCOVA; $\alpha = 0.05$) with DBH treated as a covariate. An a priori test of the mean ΔAC of trees in the insecticide treatment group and zero was conducted to determine if there was significant decline in the mean canopy condition ($\alpha = 0.05$). We also conducted a repeated measures analysis of variance (RM ANOVA) test in which treatments (control v. insecticide) were treated as the between subjects effect and annual ash canopy conditions as the

repeated measure. While the ΔAC approach only considers shifts in tree health between the beginning and end of the experiment, the RM ANOVA examines the changes in tree condition each year throughout the treatment period. Because significant differences were found in the canopy decline between insecticide treated trees and control trees, we conducted a follow-up analysis excluding the control treatment to test our hypotheses regarding the differential decline in ash trees with different initial canopy conditions. Here we conducted an ANCOVA with ΔAC as the dependent variable but the with initial ash canopy condition as a categorical effect and with DBH as a covariate. Post-hoc Tukey's HSD pairwise comparisons ($P < 0.05$) were used to test differences between canopy decline ash trees with different initial canopy conditions. All statistics were analyzed using SYSTAT v. 12 statistical software (Systat 2007).

Results:

Effectiveness of Enamectin Benzoate Injections

The EAB-induced ash canopy decline (ΔAC) was significantly different between the control (untreated) and insecticide treatments (ANCOVA, $F_{1,50} = 93.001$; $P < 0.001$), with canopies in the control treatment declining ($\Delta AC = -2.38$) more than the insecticide treatment ($\Delta AC = -0.41$) (Figure 2). Canopies of all ash trees in the control group declined to AC 5 (dead) while only ~16 % (5 trees) of trees in the insecticide treatment group declined to AC 5. The marginal but non-significant decline in mean ΔAC of insecticide treated trees over the course of the experiment indicates general success of the treatment; this decline was driven most strongly by trees in AC 3 and AC 4 (Figure 3 a & b). There was no relationship between initial DBH and tree decline across treatments (ANCOVA, $F_{1,50} = 1.213$; $P = 0.276$). Similarly, the RM ANOVA also revealed the significant treatment effect (RM ANOVA $F_{1,51} = 32.762$; $P < 0.001$) and time effect (RM ANOVA, $F_{3,49} = 46.686$; $P < 0.001$, Wilks's Lambda $P < 0.001$) (Figure 2). The significant time effect indicates that there is a change in AC over time, an increase in AC this case, when averaged over treatment levels. There was a significant treatment*time interaction whereby control trees declined over time relative to the insecticide treated trees (RM ANOVA, $F_{3,153} = 72.324$; $P < 0.001$).

Importance of Initial Canopy Conditions on Treatment Effectiveness

In insecticide treated trees, ash canopy decline varied significantly among initial canopy condition classes (initial AC) (ANOVA, $F_{3,27} = 5.514$; $P = 0.004$; Figure 3). Trees with lightly impacted canopies (AC 1 and 2) maintained canopy health relative to the other canopy condition classes (Figure 3). While the average canopy condition in trees initially ranked AC 1 declined marginally, the shift was not significantly different from zero, and because these trees could not possibly improve in canopy condition this change was not surprising. Insecticide treated trees initially AC 2 showed the greatest improvement ($\Delta AC = 0.36$) relative to AC 3 and AC 4 which declined significantly over the course of the experiment despite treatment (Figure 3).

Discussion:

The results of our research indicate that biennial emamectin benzoate treatment of ash trees can successfully maintain or improve tree health in lightly and moderately EAB infested trees (AC 1 and 2 respectively) over short to moderate periods. Overall, the mean canopy health of emamectin benzoate treated trees did not significantly change over the four-year course of the experiment. These findings are consistent with other studies which highlight the effectiveness of emamectin benzoate tree injections on EAB adults and larvae which exhibited nearly 99% mortality feeding on treated tissue (Herms et al. 2014; Smitley et al. 2010; McCullough et al. 2011). No relationship between treated ash tree size and canopy senescence was observed, indicating that trees ranging from 7-27.6 cm DBH (as studied herein) exhibited similar beneficial responses to the emamectin benzoate treatment. This result also suggests that systemic emamectin benzoate injections may be equally effective on larger trees than utilized in the study. Higher application rates of emamectin benzoate (0.4-0.6 g active ingredient for every 2.5 cm DBH) have been approved and are recommended for larger trees, although results indicate that treatment at these higher levels may not confer added protection (Smitley et al. 2010; Herms 2011).

While previous research has indicated that emamectin benzoate can maintain ash tree health at low pretreatment EAB densities (McCullough et al. 2011), its effectiveness on trees in later stages of decline (AC4 or >50 % canopy decline) was previously unknown. Our results confirm that emamectin benzoate injections can be used to stabilize canopy health in lightly impacted trees (AC 1) and highlight that it can improve trees exhibiting light to moderate EAB infestations (AC 2). Additionally, our results indicate the mixed success of spring emamectin

benzoate injections on trees in later stages of decline (AC 3 and 4), with nearly 63 % of treated trees in later stages of decline dying over the course of the experiment (Figure 3). Our data reveals success in treating trees exhibiting canopy thinning and lends support to the theory that insecticide treatment of trees with > 50 % canopy decline may not be as successful as it is for trees with healthy canopies (Herms et al. 2014); yet 37 % of the heavily impacted trees treated were able to recover or stabilize so it may be worth the investment to treat valuable landscape trees. Furthermore, the lower effectiveness of emamectin benzoate on trees with high EAB densities and in later stages of canopy decline is likely be attributed to their inability to distribute the compound from the injection site to the rest of the canopy because of reduced sapflux rates caused by EAB damage to tree vascular tissues (Flower et al. 2011).

Although the results of this study highlight the effectiveness of emamectin benzoate treatments at controlling EAB induced tree decline after a tree is infested, we reiterate current recommendations and suggest that trees should be treated before symptoms are evident for maximum efficacy (Herms et al. 2014). Spring treatment of vulnerable ash trees with emamectin benzoate and other insecticides (Table 1) has been demonstrated to be most effective at controlling EAB as the insecticide is distributed throughout the cambium and transmitted to the leaf tissue prior to adult emergence (Herms et al. 2014; McCullough et al. 2011). In EAB impacted areas, insecticide treatments of ash trees in residential areas will maintain tree health and help foster wellbeing of citizens and of the urban ecosystem (Hunter 2011). To maintain ash across the landscape, homeowners and municipalities should recognize and plan for sustained biennial emamectin benzoate injections throughout the duration of the EAB outbreak to prevent future infestation.

Although treatment of ash trees may help maintain biogeochemical and hydrological cycling in EAB impacted regions (Fissore et al. 2012), insecticide use should be minimized as there can be negative consequences for non-target insects. Non-target insects may be impacted through three primary sources of exposure: 1) direct insecticide contact, although this is virtually eliminated with applications via tree injection systems, 2) residual uptake from another surface and 3) food chain uptake (Croft 1990). Applications of insecticides via soil drenching and broadcast spraying likely have the highest non-target impacts relative to injections. Soil drenches of imidacloprid have been shown to negatively impact beneficial soil-dwelling arthropods

(Kreutzweiser et al. 2008 a,b). Although of minor concern because of the current recommended use patterns, its plausible that long-term application of emamectin benzoate for ash tree protection from EAB could result in the evolution of resistant EAB or other non-target insects, as has been observed in reduced sensitivity of salmon lice to emamectin benzoate (Lees et al. 2008; Espedal et al. 2013).

Acknowledgements:

We thank K. Costilow for assisting with experimental maintenance and M. Mason for assistance with pesticide application. The authors also thank D.A. Herms for supplying emamectin benzoate for the 2012 treatments and for his insights into insecticide applications. We also thank the anonymous reviewers for their insightful comments. This research was funded in part by the National Science Foundation Grant DGE-0549245, “Landscape Ecological and Anthropogenic Processes” and the University of Illinois at Chicago (Hadley Award and Provost Fellowship). This research would not have been possible without access to and permission to manipulate an experimental ash forest located at the USDA FS Northern Research Station Delaware, OH laboratory.

References:

- Aukema, J.E., Leung, B., Kovacs, K., Chivers, C., Britton, K.O., Englin, J., Frankel, S.J., Haight, R.G., Holmes, T.P., Liebhold, A.M., McCullough, D.G., Von Holle, B. 2011. Economic impacts of non-native forest insects in the continental United States. PLoS ONE 6, e24587. doi:10.1371/journal.pone.0024587
- Ball, J., Simmons, G. 1980. The relationship between bronze birch borer and birch dieback. J. Arboric. 6(12), 309-314.
- Bolund, P., Hunhammar, S. 1999. Ecosystem services in urban areas. Ecol. Econ. 29, 293-301.
- Cappaert, D., McCullough, D.G., Poland, T.M., Siegert, N.W. 2005. Emerald ash borer in North America: A research and regulatory challenge. Am. Entomol. 51, 152-165.
- Costanza, R., d'Arge, R., de Groot, R., Farber, S., Grasso, M., Hannon, B., Limburg, K., Naeem, S., O'Neill, R., Paruelo, J., Raskin, R.G., Sutton, P., van den Belt, M. 1997. The value of the world's ecosystem services and natural capital. Nature 387, 253-260.
- Croft, B.A. 1990 Arthropod biological control agents and pesticides. Wiley and Sons: New York. 703 p.
- Dwyer, J.F., Nowak, D.J., Noble, M.H., Sisinni, S.M. 2000. Connecting people with ecosystems in the 21st century: an assessment of our nation's urban forests. Gen. Tech. Rep. PNW-GTR-490. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 483 p.
- Espedal, P.G., Glover, K.A., Horsberg, T.E., Nilsen, F. 2013. Enamectin benzoate resistance and fitness in laboratory reared salmon lice (*Lepeophtheirus salmonis*). Aquaculture 416-417, 111-118.
- Fettig, C.J., Grosman, D.M., Munson, A.S. 2013. Advances in insecticide tools and tactics for protecting conifers from bark beetle attack in the western United States. In: S. Trdan (Ed.) Insecticides - Development of Safer and More Effective Technologies. InTech, Rijeka, Croatia. P. 472-492.
- Fissore, C., McFadden, J.P., Nelson, K.C., Peters, E.B., Hobbie, S.E., King, J.Y., Baker, L.A., Jakobsdottir, I. 2012. Potential impacts of emerald ash borer invasion on biogeochemical and water cycling in residential landscapes across a metropolitan region. Urban Ecosyst. 15, 1015-1030.
- Flower, C.E., Gonzalez-Meler, M.A. 2015. Responses of temperate forest productivity to insect and pathogen disturbances. Ann. Rev. Plant Biol. 66: 547-569.
- Flower, C.E., Lynch, D.J., Knight, K.S., Gonzalez-Meler, M.A. 2011. EAB induced tree mortality impacts ecosystem respiration and tree water use in an experimental forest. In: Parra G, Lance D, Mastro V, Reardon R, Benedict C, comps. 2011 emerald ash borer national research and technology development meeting; 2011 October 12-13; Wooster,

331 OH. FHTET-2011-06. Morgantown, WV; U.S. Department of Agriculture, Forest
 332 Service, State and Private Forestry, Forest Health Protection, Forest Health Technology
 333 Enterprise Team: 115-116.

334 Flower, C.E., Knight, K.S., Gonzalez-Meler, M.A. 2013a. Impacts of the emerald ash borer
 335 (*Agrilus planipennis* Fairmaire) induced ash (*Fraxinus* spp.) mortality on forest carbon
 336 cycling and successional dynamics in the eastern United States. *Biol. Invasions* 15, 931-
 337 944.

338 Flower, C.E., Knight, K.S., Rebbeck, J., Gonzalez-Meler, M.A. 2013b. The relationship between
 339 emerald ash borer (*Agrilus planipennis* Fairmaire) and ash (*Fraxinus* spp.) tree decline:
 340 using visual ash condition assessments and leaf isotope measurements to assess pest
 341 damage. *For. Ecol. Manag.* 303, 143-147.

342 Flower, C.E., Long, L.C., Knight, K.S., Rebbeck, J., Brown, J.S., Gonzalez-Meler, M.A.,
 343 Whelan, C.J. 2014. Native bark-foraging birds preferentially forage in infected ash
 344 (*Fraxinus* spp.) and prove effective predators of the invasive emerald ash borer (*Agrilus*
 345 *planipennis* Fairmaire). *For. Ecol. Manag.* 313, 300-306.

346 Grosman, D.M., Clarke, S.R., Upton, W.W. 2009. Efficacy of two systemic insecticides injected
 347 into loblolly pine for protection against southern pine bark beetles (Coleoptera:
 348 Curculionidae). *J. Econ. Entomol.* 102, 1062-1069.

349
 350 Harris, E.M., Martin, D.G., Polsky, C., Denhardt, L., Nehring, A. 2013. Beyond “Lawn People”:
 351 The role of emotions in suburban yard management practices. *Prof. Geogr.* 65 (2), 345-
 352 361.

353
 354 Herms, D.A. 2011. Multiyear evaluation of efficacy of soil and trunk applied systemic
 355 insecticides for ash canopy conservation. *In: Parra G, Lance D, Mastro V, Reardon R,*
 356 *Benedict C, comps. 2011 emerald ash borer national research and technology*
 357 *development meeting; 2011 October 12-13; Wooster, OH. FHTET-2011-06.*
 358 *Morgantown, WV; U.S. Department of Agriculture, Forest Service, State and Private*
 359 *Forestry, Forest Health Protection, Forest Health Technology Enterprise Team: 26-27.*

360 Herms, D.A., McCullough, D.G., Smitley, D.R., Clifford, C.S., Cranshaw, W. 2014. Insecticide
 361 options for protecting ash trees from emerald ash borer. *North Central IPM Center*
 362 *Bulletin. 2nd Edition. 16 pp.*

363 Hunter, M.R. 2011. Impact of ecological disturbance on awareness of urban nature and sense of
 364 environmental stewardship in residential neighborhoods. *Landsc. Urban Plan.* 101 (2),
 365 131-138.

366 Knight, K.S., Flash, B.P., Kappler, R.H., Throckmorton, J.A., Grafton, B., Flower, C.E. 2014.
 367 Monitoring ash (*Fraxinus* spp.) decline and emerald ash borer (*Agrilus planipennis*)
 368 symptoms in infested areas. *Gen. Tech. Rep. NRS-139. Newtown Square, PA: U.S.*
 369 *Department of Agriculture, Forest Service, Northern Research Station. 18p.*

370 Kovacs, K.F., Haight, R.G., McCullough, D.G., Mercader, R.J., Siegert, N.W., Liebhold, A.M.

371 Cost of potential emerald ash borer damage in U.S. communities, 2009-2019. *Ecol. Econ.*
372 69, 569-578.

373 Kreutzweiser, D.P., Good, K.P., Chartrand, D.T., Scarr, T.A., Holmes, S.B., Thompson, D.G.
374 2008a. Effects on litter-dwelling earthworms and microbial decomposition of soil-applied
375 imidacloprid for control of wood-boring insects. *Pest Manag. Sci.* 64, 112-118.

376 Kreutzweiser, D.P., Good, K.P., Chartrand, D.T., Scarr, T.A., Thompson, D.G. 2008b. Toxicity
377 of the systemic insecticide, imidacloprid, to forest stream insects and microbial
378 communities. *B. Environ. Contam. Tox.* 80, 211-214.

379 Lees, F., Baillie, M., Gettinby, G., Revie, C.W. 2008. Factors associated with changing efficacy
380 of emamectin benzoate against infestations of *Lepeophtheirus salmonis* on Scottish
381 salmon farms. *J. Fish Dis.* 31, 947-951.

382 MacFarlane, D.W., Meyer, S.P. 2005. Characteristics and distribution of potential ash tree hosts
383 for emerald ash borer. *For. Ecol. Manag.* 213, 15-24.

384 McCullough, D.G., Smitley, D.R., Poland, T.M. 2003. Evaluation of insecticides to control
385 emerald ash borer adults and larvae. Available online at [www.emeraldashborer.info/](http://www.emeraldashborer.info/files/bulletin.pdf)
386 files/bulletin.pdf; last accessed April 2014.

387
388 McCullough, D.G., Poland, T.M., Anulewicz, A.C., Lewis, P., Cappaert, D. 2011. Evaluation of
389 *Agrilus planipennis* (Coleoptera: Buprestidae) control provided by emamectin benzoate
390 and two neonicotinoid insecticides, one and two seasons after treatment. *J. Econ.*
391 *Entomol.* 104, 1599-1612.

392
393 McKenney, D.W., Pedlar, J.H. 2012. To treat or remove: an economic model to assist in
394 deciding the fate of ash trees threatened by emerald ash borer. *Arboric. Urban For.* 38 (4),
395 121-129.

396
397 Nowak, D.J. 1994. Urban forest structure: the state of Chicago's urban forest, pp. 3-19. In:
398 McPherson, E.G., Nowak, D.J. & Rowntree, R. A. (Eds.). *Chicago's urban forest*
399 *ecosystem: results of the Chicago Urban Forest Climate Project*. Gen. Tech. Rep. NE-
400 186. Radnor, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest
401 Experiment Station: 201 p.

402
403 Peterson, G., Allen, C.R., Holling, C.S. 1998. Ecological resilience, biodiversity, and scale.
404 *Ecosystems* 1, 6-18.

405 Pimentel, D., Zuniga, R., Morrison, D. 2005. Update on the environmental and economic costs
406 associated with alien-invasive species in the United States. *Ecol. Econ.* 52, 273-288.

407 Sander, H.A., Polasky, S., Haight, R.G. 2010. The value of urban tree cover: A hedonic property
408 price model in Ramsey and Dakota Counties, Minnesota, USA. *Ecol. Econ.* 69, 1545-
409 1656.

410 Siegert, N.W., McCullough, D.G., Liebhold, A.M., Telewski, F.W. 2007. Resurrected from the
411 ashes: as historical reconstruction of emerald ash borer dynamics through

- dendrochronological analysis. In: Mastro, V., Reardon, R. (eds) Proceedings of the Emerald Ash Borer Research and Technology Development Meeting; 2006 October 29–November 2; Cincinnati, OH. FHTET 2007-04. Morgantown, WV: US Forest Service, Forest Health Technology Enterprise Team, pp 18–19.
- Smith, A. 2006. Effects of community structure on forest susceptibility and response to the emerald ash borer invasion of the Huron River watershed in southeast Michigan. M.S. Thesis. The Ohio State University. Columbus, OH. 122p.
- Smith, A., Herms, D.A., Long, R.P., Gandhi, K.J.K. 2015. Community composition and structure had no effect on forest susceptibility to invasion by the emerald ash borer (Coleoptera: Buprestidae). *Can. Entomol.* 147: 318-328.
- Smitley, D., Davis, T., Rebek, E.. 2008. Progression of ash canopy thinning and dieback outward from the initial infestation of emerald ash borer (Coleoptera: Buprestidae) in Southeast Michigan. *J. Econ. Entomol.* 101: 1643 – 1650.
- Smitley, D.R., Doccia, J.J., Cox, D.L. 2010. Multiple-year protection of ash trees from emerald ash borer with a single trunk injection of emamectin benzoate, and single-year protection with an imidacloprid basal drench. *Arboric. Urban For.* 36 (5), 206-211.
- SYSTAT. 2007. SYSTAT 12.0 for windows. SPSS, Chicago.
- Wang, X.Y., Yang, Z.Q., Gould, J.R., Zhang, Y.N., Liu, G.J., Liu, E.S. 2010. The biology and ecology of the emerald ash borer, *Agrilus planipennis*, in China. *J. Insect Sci.* 10, 128. doi: 10.1673/031.010.12801.

Table 1: Insecticide products, application methods, timing and effectiveness for treating *Fraxinus* sp. trees for emerald ash borer control.

Active Ingredient	Insecticide Formulation	Application Method	Application Timing	Effectiveness
Acephate	Orthene	Trunk Spray	Spring-Early Summer	Low to Moderate ^d
Bidrin®	Inject-A-Cide B®	Trunk Injection	Spring-Early Summer	Low to Moderate ^a
Carbaryl	Sevin®	Trunk Spray		
Dinotefuran	Safari TM	Trunk Spray	Spring-Early Summer	Moderate ^c
Emamectin Benzoate	TREE-äge TM	Trunk Injection	Spring-Early Summer	High ^{b,c,e}
Imidacloprid	Bayer Advanced Tree & Shrub Insect Control [†]	Soil Drench	Fall-Spring	Mixed ^d
Imidacloprid	IMA-jet®	Trunk Injection	Spring-Early Summer	High ^d
Imidacloprid	Imicide®	Trunk Injection	Spring-Early Summer	Moderate ^c
Imidacloprid	Merit®	Soil Injection/Drench	Spring-Early Summer	Mixed ^{d,e}
Imidacloprid	Xytect TM	Soil Injection/Drench	Spring-Early Summer	Mixed ^d
Imidacloprid	Macho 2F	Trunk Spray	Spring-Early Summer	Moderate ^c
Pyrethroid (Bifenthrin)	Onyx	Trunk/Foliar Spray	Biannual Application (spring and summer)	Moderate ^d
Pyrethroid (Cyfluthrin)	Tempo SC Ultra	Whole Tree spray	Spring-Early Summer	Moderate ^d
Pyrethroid (Cyfluthrin)	Pointer	Trunk Injections	Spring-Summer	Moderate ^d

[†]Appropriate for homeowner application, ^aMcCullough, et al. (2004), ^b Flower et al. (herein), ^c McCullough et al. (2011), ^dHermes et al. (2009; 2014), ^e Smitley et al. (2010).

Figure 1: Visual classification scheme denoting ash canopy condition class (AC) 1-5 (Knight et al. 2014; Smith et al. 2015).

Figure 2: Ash canopy condition class (AC) in the control and insecticide treated ash trees from 2010-2013. Values represent means $\pm$ SE.

Figure 3: (A) The relationship between initial ash canopy condition and the timeseries progression of each canopy condition class of insecticide treated trees over the course of the experiment and (B) the relationship between initial ash tree canopy condition (initial AC) and ash canopy decline over the course of the experiment (ΔAC ; $AC_{2010} - AC_{2013}$) (ANCOVA; $P < 0.001$). Error bars denote ± 1 SE. In panel B, letters represent significance levels determined by Post Hoc Tukey's HSD pairwise comparisons ($P < 0.05$).