Identification of putative circadian clock genes in the American horseshoe crab, *L. polyphemus*

Kevin N. Chesmore^a, Winsor H. Watson III^b, Christopher C. Chabot^c

For submission to Comp Biochem Physiol D

^aDepartment of Genetics,
Dartmouth College
Hanover, NH 03755 USA

^bDepartment of Biological Sciences
University of New Hampshire
Durham, NH 03824 USA

^cDepartment of Biological Sciences, MSC#64
Plymouth State University
Plymouth, NH 03264 USA

Running Head: Putative circadian proteins in horseshoe crabs

ms. has 20 pages, 12 figures, 3 tables, 4 suppl. figures & 1 suppl. table

Corresponding author contact:

Kevin N. Chesmore
508 479-7914 24

Kevin.N.Chesmore.GR@dartmouth.edu

Abstract:

While the American horseshoe crab, *Limulus polyphemus*, has robust circadian and circatidal rhythms, virtually nothing is known about the molecular basis of these rhythms in this species or any other chelicerate. In this study, next generation sequencing was used to assemble transcriptomic reads and then putative homologs of known core and accessory circadian genes were identified in these databases. Homologous transcripts were discovered for one circadian clock input gene, five core genes, 22 accessory genes, and two possible output pathways. Alignments and functional domain analyses showed generally high conservation between the putative *L. polyphemus* clock genes and homologs from *Drosophila melanogaster* and *Daphnia pulex*. The presence of both *cry1* and *cry2* in the *L. polyphemus* transcriptome would classify its system as an “ancestral”, type 2 clock system. In addition, a novel duplication of CYCLE, and a novel triplication of PERIOD were found. Investigations are currently underway to determine if any of these “circadian” genes also participate in the molecular processes that drive the *Limulus* circatidal clock.

Key words: circadian, biological rhythm, clock, transcriptome, horseshoe crab, chelicerate, circatidal, de novo, Illumina, clock genes.

1. Introduction.

Endogenously driven biological rhythms have been observed in virtually all organisms in which a concerted effort has been made to look for them (Dunlap, 1999). These rhythmic changes in physiology and behavior are driven by internal biological clocks and help organisms to both anticipate, and synchronize to, predictable rhythmic changes in their environment. These clocks are composed of three essential elements: 1) the input proteins that allow the clock to synchronize to environmental rhythms, such as light:dark cycles; 2) the clock itself, made up of the proteins that allow the clock to keep time; and 3) the output proteins that help to mediate changes in physiology and behavior (Allada and Chung, 2010; Dunlap, 1999). The molecular basis of these three components of the clock have been elucidated in a few model systems such as the fruit fly, *Drosophila melanogaster* and the mouse, *Mus musculus* (Zhang and Kay, 2010) and these findings have provided the framework for studies in other organisms. Although the specific molecular mechanisms of the circadian clock vary greatly between distantly related animal models (i.e. *M. musculus* and *D. melanogaster*), the general architecture appears to be well preserved across nearly all living organisms (Dunlap, 1999). In plants, fungi, and animals, circadian clocks appear to be composed of interlocking transcription-translation cycles that feedback to control their own transcription.

These circadian clocks involve two sets of proteins: the core proteins (which are part of a negative feedback loop) and the accessory proteins (which modulate the core proteins and are necessary for maintaining the ~24hr periodicity of the core clock). In *D. melanogaster* the core clock is composed of four proteins: PERIOD (PER), TIMELESS (TIM), CLOCK (CLK), and CYCLE (CYC) (Allada and Chung, 2010). When the *per* and *tim* genes are transcribed and translated, they form a heterodimer that acts at the *clk* promoter to up-regulate CLK expression (Allada and Chung, 2010; Chang and Reppert, 2003). In other insect species Non-photoreceptive(np) CRYPTOCHROME 2 (npCRY2) acts in addition to, or in place of, TIM (Rubin et al., 2006; Yuan et al., 2007). As PER and TIM build up over time, they dimerize and become phosphorylated by accessory genes such as CASEIN KINASE I ϵ (CKI ϵ). This phosphorylation allows this heterodimer to enter the nucleus and to remove CLK-CYC from the E-box, turning off *per* and *tim* (Allada and Chung, 2010). The proteins CLK and CYC (which appear to be constitutively expressed) are, in turn, capable of repressing the activity of *per* and *tim*, completing the negative feedback loop.

This core negative feedback loop is also modulated by a host of accessory proteins, such as VRILLE (VRI), CLOCKWORK ORANGE (CWO), and PAR DOMAIN PROTEIN 1 ϵ (PDP1 ϵ), which form a secondary feedback loop with CLK (Allada and Chung, 2010). Additionally, the proteins SUPERNUMERARY LIMBS (SLIMB) and ARYL HYDROCARBON RECEPTOR NUCLEAR TRANSPORTER (ARNT) also help to modulate the transcription of the core clock components. The protein kinases CASEIN KINASE II α (CKII α), and CASEIN KINASE II β (CKII β) serve to phosphorylate PER, while TIM is phosphorylated by CKII α , CKII β , CASEIN KINASE I α (CKI α), SHAGGY (SGG), and JETLAG (JET). Along with these kinases, the protein phosphatases PROTEIN PHOSPHATASE 1 α (PP1 α), PROTEIN PHOSPHATASE 1 β (PP1 β), PROTEIN PHOSPHATASE 2 α -MTS (PP2 α -MTS), PROTEIN PHOSPHATASE 2 α -WBT (PP2 α -WBT), PROTEIN PHOSPHATASE 2 α -WS (PP2 α -TWS) are also crucial for proper clock function (Allada and Chung, 2010).

The input portion of the circadian clock system is provided by the photoreactive(p) protein CRYPTOCHROME 1 (pCRY1). In *D. melanogaster*, pCRY1 undergoes a conformational change in the presence of blue light (such as in sunlight) and ubiquitinates TIM, tagging it for degradation via the proteasome (Emery et al., 1998; Tauber et al., 2004; Yuan et al., 2007). This degradation acts to “reset” the clock, because when TIM is degraded, it can no longer form a heterodimer with PER and the transcriptional regulation of *clk* is terminated.

The neuropeptides Pigment Dispersing Hormone (PDH) and NEUROPEPTIDE F (NPF) (Lee et al., 2006; Strauss et al., 2011; Taghert and Shafer, 2006) are crucial for the output of the clock in *D. melanogaster*. Ablation of PDH cells, or mutations of the *pdh* gene, disrupts the expression of circadian rhythms in this species *D. melanogaster* (Renn et al., 1999), and these effects are exacerbated by the ablation of NPF containing cells (Hermann et al., 2012). While these peptides are thought to induce numerous physiological and behavioral changes (Lee et al., 2006; Strauss et al., 2011), the mechanism of the control of either NPF or PDH release by the core circadian clock is not known (Depetris-Chauvin et al., 2011).

While much is known about the molecular mechanisms of the circadian clock in *D. melanogaster* and a few other insect species (Dunlap, 1999; Tomioka and Matsumoto, 2010), much less is known about the circadian clocks of non-insect arthropods, with the exception of *Daphnia pulex* (Tilden et al., 2011) and *Calanus finmarchicus* (Christie et al., 2013). One entire sub-phylum of Arthropoda that has been ignored is the chelicerata, which encompasses a range of organisms including spiders, ticks, mites, scorpions, and horseshoe crabs (Giribet and Edgecombe, 2012). The American horseshoe crab, *Limulus polyphemus*, has long been known to exhibit robust circadian rhythms of lateral eye sensitivity to light and has served as a model species for studies in photophysiology and its circadian control (Barlow, 1983; Barlow et al., 1980). Bob Barlow, his colleagues, and other investigators have shown that more than twenty independent changes occur in the eye of *L. polyphemus* to achieve an approximately 100,000 fold change in eye sensitivity between night and day (Barlow et al., 1980; Battelle, 2013). Yet, despite decades of study of this system, the molecular mechanisms that drive this clock in *L. polyphemus* are completely unknown.

Horseshoe crabs also have an additional timing system that serves to synchronize its locomotor activity to the ~ 12.4 hour tidal cycles (Chabot and Watson, 2010; Watson et al., 2008). While circatidal rhythms such as this have been documented in several species, the molecular basis of these clocks is unknown (de la Iglesia and Hsu, 2010; Takekata et al., 2014; Tessmar-Raible et al., 2011). Furthermore, there is some controversy surrounding the nature of the timing system that drives these rhythms. The two primary competing theories that have been put forth to explain the underlying mechanisms that give rise to these 12.4 h rhythms are: 1) The circatidal oscillator theory, which states that two bouts of activity per day are controlled by one, ~ 12.4 h, circatidal clock (which can also be modulated by a circadian clock (Naylor, 1996)) and; 2) the circalunidian theory, which states that each bout of activity is controlled by a separate, ~24.8 hr (the time between successive moonrises), circalunidian oscillator (Palmer and Williams, 1986). This circalunidian model has been well supported by behavioral evidence in *L. polyphemus* (Chabot et al., 2016; Chabot and Watson, 2010) and our working hypothesis is that each bout of activity is controlled by a modified circadian (Palmer and Williams, 1986) clock that has a slightly longer period than 24 hours. This would be similar to the situation in *D. melanogaster*, where dawn and dusk bouts of activity are controlled by separate, coordinated, circadian clocks (Stoleru et al., 2004). The goal of this study was to determine if horseshoe crabs possess some of the circadian genes found in other model species. The identification of these genes in *L. polyphemus* would not only provide the scientific community with a resource to begin studying circadian clock in chelicerates, but may also allow us to begin to understand the molecular basis of circatidal/circalunidian clocks.

2. Methods

2.1. Animals and RNA extraction. *L. polyphemus* individuals were collected from Great Bay in Durham NH. Animals were treated in accordance with NIH guide for the care and use of animals (NIH publications No. 8023, revised 1978). The entire central nervous system (brain and ventral nerve cord) was dissected from 4 individuals, immediately snap frozen on dry ice, and stored in 1.5ml Eppendorf tubes at -80°C. Samples were then shipped on dry ice to University of Vermont Cancer Center DNA Analysis Facility (Burlington, VT), where RNA was extracted using Trizol RNA extraction (Invitrogen, Carlsbad, CA) and cleaned using the Qiagen RNeasy Mini column (Qiagen, Valencia, CA). Quality and concentration of cleaned RNA was determined using Nanodrop spectrophotometer (Thermo Scientific, Madison WI), Qubit Spectrofluorometer (Life Technologies, Carlsbad, CA) and Agilent Bioanalyzer 2100 (Agilent technologies, Santa Clara, CA).

2.1. Library construction and sequencing. A transcriptome library was constructed using Illumina TruSeq RNA Sample Prep LT version 2 (RS-122-2001/2002). 1 ug of each sample was PolyA enriched using magnetic beads and reverse transcribed into cDNA via Superscript II [Invitrogen]. The cDNA was then fragmented, end-repaired, adenylated, and tagged with sequence adaptors for source identification. The cDNA was sequenced using 12pM/flow cell lane with Illumina CBOT and HiSeq1000/1500.

2.2. Transcriptome de novo assembly. Transcriptome reads were de novo assembled using the CLC genomics workbench proprietary algorithm (CLC version 5.1.2, CLCBio, Boston, MA), as well as SOAP de novo

assembly software (Luo et al., 2012) (<http://soap.genomics.org.cn/soapdenovo.html>, version 1.02) with the default settings.

2.2. Transcriptome mining and gene identification. The transcriptome was mined with the CLC genomics workbench tBLASTn program (version 5.1.2), using *D. melanogaster* and other arthropod query sequences. The top hits from the *L. polyphemus* transcriptome were translated and subjected to a reciprocal BLASTp against the entire UniProtKB protein database to verify sequence homology. A second reciprocal BLASTp was also performed against *D. melanogaster* and *D. pulex* circadian proteins exclusively to compare scores to these model organisms.

2.3. Sequence analysis. Transcripts of interest were translated into protein sequences using CLC genomics workbench (version 5.1.2). These sequences were then aligned to respective orthologs of *D. melanogaster* and *D. pulex*, along with an ortholog of a chelicerate species when one was available using CLUSTAL Omega program (<http://www.ebi.ac.uk/Tools/msa/clustalo/>). Functional domains of all proteins of interest in *L. polyphemus*, *D. melanogaster*, *Daphnia*, and chelicerate species when present were identified using SMART Genome program (<http://smart.embl-heidelberg.de/>) (Letunic et al., 2015; Schultz et al., 1998)). The %Identity/%Similarity of the amino acid sequences these proteins and their respective functional domains were calculated using default settings of the SIAS program (<http://imed.med.ucm.es/Tools/sias.html>). Sequence alignments from CLUSTAL Omega were pasted into a word document and functional domains were color coded as follows: Light blue:bHLH, Green:PAS, Red:PAC, Grey:Period C, Dark green:Photolyase, Gold:FAD Binding 7, Dark blue:Timeless. Phylogenetic trees were constructed using Mega 5.2 (<http://www.megasoftware.net/>). Sequences, other than *L. polyphemus* proteins, were extracted from UniProt database and aligned using MUSCLE (<http://www.ebi.ac.uk/Tools/msa/muscle/>). After alignment a Maximum likelihood tree was generated using default setting and 500 Bootstraps. Tree branches were color coded to specify different clades on the trees: Red:Chelicerata, Blue:Non-chelicerate arthropods, Black:Vertebrates, Violet:Fungi, Green:Nematode. Diamonds were added at gene duplication events.

3. Results.

3.1. Top 5 BLAST hits from UniProtKB

In our analysis of the *L. polyphemus* (*Lp*) transcriptome we identified orthologs of all five core clock components (PER, TIM, CLK, CYC, npCRY2), including 3 copies of the *per* gene [*perioda* (*pera*), *periodb* (*perb*), and *periodc* (*perc*)] and 2 copies of the *cyc* gene [*cycle1* (*cyc1*) and *cycle2* (*cyc2*)]. Top BLAST hits for all core clock genes are reported in Table 1. A reciprocal BLASTp of all three *LpPER* paralogs against the UniProt KB Protein database showed closest homology to arthropod PER proteins, including orthologs from three insect species and one chelicerate species (*Ixodes scapularis*, Black legged tick). BLASTp of *LpTIM*, *LpCLK*, *LpCYC1*, *LpCYC2* and *LpnpCRY2* showed closest homology to insect orthologs. Top BLAST hits for all core clock components were significant, with e-values ranging from 10^{-114} to less than 10^{-180} , and BLAST scores ranging from 1004 to 2199.

We also identified orthologs of 15 accessory clock components (VRI, CWO, SLIMB, ARNT, CK I α , CK I ϵ , CK II α , CK II β , JET, SGG, PP1 α , PP1 β , PP2 α MTS, PP2 α WBT, PP2 α TWS), including duplications of both the *LpSLIMB* and *LpARNT* genes, for a total of 17 accessory genes. Only one known accessory clock gene (PDP1) was not found in the *L. polyphemus* transcriptome. Top BLAST hits for all accessory clock genes are reported in Table 1. A reciprocal BLAST of the *L. polyphemus* orthologs revealed nearly all clock genes (VRI, CWO, SLIMB1, SLIMB2, ARNT1, ARNT2, CKI α , JET, SGG, PP1 α , PP1 β , PP2 α -MTS, PP2 α -MTS, PP2 α -TWS) are most closely related to arthropod orthologs. However, CKI α , CKI ϵ , and CKII β all show highest levels of homology to vertebrate orthologs. The top 5 BLAST hits of 10 of the 17 accessory proteins included at least one known chelicerate ortholog, primarily from the tick family. Top BLAST hits for nearly all accessory clock components were significant, with e-values ranging from 10^{-31} to less than 10^{-180} , and BLAST scores ranging from 320 to 2138 (only *LpVRI*, *LpCWO*, and *LpJET* score below 1000).

We were able to identify partial sequences for the input protein pCRY1, the NPF receptor (NPFR) and the PDH receptor (PDHR), as well as a complete sequence for the output neuropeptide NPF. PDH was not identified in the transcriptome of *L. polyphemus*. Top BLAST hits for all input and output clock genes are reported in Table 1. The top BLAST hits for *LppCRY1* showed that *LppCRY1* is most closely related to insect pCRY1 sequences. Despite only being a partial sequence this protein is highly conserved and had an e-value of 10^{-132} , with a BLAST score of 1022. BLASTp of *LpNPF*, *LpNPFR*, and *LpPDHR* showed closest homology to arthropod orthologs, including several chelicerates. Top BLAST hits for most output clock components (excluding *LpNPF*) were significant, with e-values ranging from 10^{-60} to 10^{-150} , and BLAST scores ranging from 533 to 1022. Top BLAST hits for *LpNPF* showed e-values of 10^{-3} and BLAST scores of 84 and 86. The high e-values and low BLAST scores of the *LpNPF* BLAST are likely because NPF is a short peptide (96 amino acids long).

3.2. Comparisons with *D. melanogaster* and *D. pulex* clock genes

Top BLAST hits for all core clock genes against the *D. melanogaster* and *D. pulex* databases are reported in Table 2. All putative core clock components identified in this study, with the exception of npCRY2, (PERA, PERB, PERC, TIM, CLK, CYC1, and CYC2,) were found to have the highest homology to the orthologous proteins of *D. melanogaster* and *D. pulex*. *LpnpCRY2* most closely matched a 6-4 photolyase in *D. melanogaster*, which is the closest relative to npCRY2 on the *cryptochrome* gene family tree. This mis-match is the product of the fact that the genus *D. melanogaster* lost its npCRY2 ortholog during its evolution. However, *Apis mellifera* retained its npCRY2 ortholog, and the top BLAST hit for *LpCRY2* in *Apis* is npCRY2 (data not shown). Similarly, the closest match to *LpnpCRY2* in the *D. pulex* protein database was “CRY-M”, which is another name for npCRY2. The top BLAST hits for *D. melanogaster* and *D. pulex* also showed high levels of homology to our genes of interest, with e-values ranging from 10^{-68} to less than 10^{-180} and BLAST scores ranging from 653 to 2139.

Top BLAST hits for all accessory clock genes against the *D. melanogaster* and *D. pulex* databases are reported in Table 2. All putative accessory clock components identified in this study were found to have the highest homology to the predicted orthologous proteins of *D. melanogaster* and *D. pulex*. Although on several occasions we found that the closest match to the *L. polyphemus* genes was an uncharacterized protein (as seen when *LpCWO* was blasted against the *D. pulex* database), or the name of the hit simply referred to the gene family (e.g. when the *LpSLIMB1* and *LpSLIMB2* genes matched the “f-box/wd-repeat protein” in *D. pulex*). Generally, these problems were only found in some *D. pulex* orthologs, and so the *D. melanogaster* (and many of the *D. pulex*) matches give reliable gene annotations in the BLAST results. With the exception of *LpVRI*, *LpCWO* and *LpJET*, all accessory proteins show significant levels of homology to their respective BLAST hits, with e-values ranging from 10^{-110} to less than 10^{-180} and BLAST scores ranging from 842 to 2136.

Top BLAST hits for all input and output clock genes against the *D. melanogaster* and *D. pulex* databases are reported in Table 2. All putative input and output clock components identified in this study (pCRY1, NPF, NPFR and PDHR) were found to have the highest homology to the predicted orthologous proteins of *D. melanogaster* and *D. pulex*. The top BLAST hits for the *LppCRY1* showed pCRY1 and CRY-D (*D. melanogaster*-like *cryptochrome* also known as pCRY1) as the closest hits in our model systems. Similarly the top hits for all putative output genes (NPF, NPFR, and PDHR) for both *D. melanogaster* and *D. pulex* were the expected orthologs for both of these species. With the exception of *LpNPF*, all output proteins show significant levels of homology to their respective BLAST hits, with e-values ranging from 10^{-48} to 10^{-71} and BLAST scores ranging from 445 to 583. The high e-values and low BLAST scores of the *LpNPF* BLAST are likely because NPF is a short peptide (96 amino acids long), additionally the *D. pulex* NPF ortholog appears to have not yet been characterized as the top BLAST hit showed “Putative uncharacterized protein”.

3.3. Protein and domain alignment

The majority of the core clock components were more homologous to *D. pulex* core clock proteins than those of *D. melanogaster* (with the exception of *LpCYC1* and *LpCYC2* (Table 3; Figure 1 & 2; Supp Figure 1)).

The similarity between *Limulus* clock proteins and those in *D. pulex* and *D. melanogaster* were most pronounced when comparing the functional domains of these clock proteins (Table 3). Moreover, the levels of conservation within the functional domains were fairly comparable between *D. melanogaster* and *D. pulex*, suggesting that the difference in sequence similarity lies in the regions outside and between domains (Table 3). The two clock protein domains with lowest levels of conservation were the PeriodC domains of all three *LpPER* paralogs (Table 3; Figure 3 & 4; Supp Figure 2) and the TIMELESS domain of *LpTIM* (Table 3; Figure 5; Supp Figure 3) (Table 3). *LpnpCRY2* showed the highest level of conservation in the core clock proteins (Table 3; Figure 6). The full protein alignment of *LpCYC2* to *LpCYC1* shows a %Identity and %Similarity of 86% and 89%, respectively (Table 3; Figure 2). The three copies of the *period* genes also show high levels of homology to one another, with the alignments of the full *LpPERA* and *LpPERB* proteins showing a %identity and %similarity of 35% and 48%, *LpPERA* and *LpPERC* showing 37% and 49%, and *LpPERC* and *LpPERB* showing 65% and 72% (Figure 4). The duplicate *per* and *cyc* genes show a greater amount of similarity to their respective paralogs than they do to the respective orthologs of *D. melanogaster* and *D. pulex* (*per* (Figure 3 & 4; Supp Figure 2); *cyc* (Figure 1 & 2; Supp Figure 1)).

The sequence alignments of the accessory proteins were similar to those of the core proteins. For example, the sequence alignment of the duplicated *LpSLIMB* paralogs shows an overall %identity/%similarity value of 94/97, and the paralogs of the *LpARNT* show a value of 85/89. The paralogs of both genes share higher levels of sequence identity to one another than they do to *D. melanogaster* and *D. pulex* orthologs. Moreover, the majority of the *L. polyphemus* orthologs have higher sequence identity to *D. pulex* than to *D. melanogaster*, with the higher levels of conservation within the functional domains (Table 3). We found that *LpVRI*, *LpCWO*, and *LpJET* had the lowest levels of conservation in the accessory proteins, but higher levels of conservation in the functional domains of *LpVRI* and *LpCWO* (Table 3). Interestingly, we found that the functional domains within the *LpJET* appear to have less conservation than the non-domain regions (Table 3). Additionally, the kinases and phosphatases tended to be noticeably more conserved than the other accessory proteins, with the most conserved protein (PP1 α) having a %Identity/%Similarity of 88/91 for *D. melanogaster* and 90/93 for *D. pulex* (Table 3).

The sequence alignments for the input and output proteins showed high levels of homology for both *LpCRY1*, *LpNPFR*, and *LpPDHR*, although the sequence identity for *LpNPF* was much lower. All of the output proteins had higher levels of homology within the functional domains.

3.4. Phylogenetic trees

All the core clock components for *L. polyphemus* identified in this study (*PERA*, *PERB*, *PERC*, *TIM*, *CLK*, *CYC1*, *CYC2*, and *npCRY2*) clearly fall into the appropriate clades for each gene family (Figure 8, 9, 10 & 11). We also found that whenever the protein sequence for a chelicerate ortholog was available, the *L. polyphemus* ortholog(s) most closely associated with it. Furthermore, all the chelicerate proteins tended to nest within the clades consisting of arthropod orthologs, but always formed a distinct clade outside the insect and crustacean clades. Within the *period* gene tree the three paralogs of the *L. polyphemus period* gene probably originated from two novel duplication events that occurred after the formation of the subphylum chelicerata and after the divergence of *Ixodes scapularis* (Black legged tick, and a fellow chelicerate) and *L. polyphemus* (Figure 8). Furthermore, the *periodB/periodC* duplications probably occurred more recently than the duplication event involving *periodA* and the common ancestor of *periodB* and *periodC*. Similarly, the duplication of the *cyc* gene also appears to be specific to the subphylum chelicerata (Figure 10). However, without the sequences of any other chelicerate orthologs it is not possible to determine when in the evolution of the chelicerates this duplication occurred. Additionally the phylogenetic tree of the *cry* gene family showed that *LpCRY1* nests within the invertebrate-*cry1* clade of the *cryptochrome* gene family.

In the accessory gene group we found two duplicated genes *slimb* and *arnt*. Phylogenetic analysis of the *LpSLIMB* paralogs show that both copies of *LpSLIMB* nest within the *slimb* clade (Figure 12). Similarly phylogenetic analysis of the duplicate *LpARNT* paralogs show that these protein nest within the invertebrate *arnt* clade (Figure 10). Both of these duplication events appear to have occurred after the divergence of chelicerates, and have not been identified in any other species.

4. Discussion.

4.1. Identification of putative circadian proteins in the *L. polyphemus* de novo transcriptome assembly.

Many arthropods have been shown to exhibit robust physiological and behavioral rhythms (Palmer, 1973). However, the molecular basis of many of these rhythms has been elucidated in a few model species, such as *D. melanogaster* (Allada and Chung, 2010). While some progress has been made in understanding the architecture of these clocks in non-model insect and crustacean species (Rubin et al., 2006; Tomioka and Matsumoto, 2010; Yuan et al., 2007), little effort has been put forth to investigate the clock mechanisms in chelicerates, including *L. polyphemus*. Based on the data obtained in this study, it appears that horseshoe crabs possess endogenous clocks that may rely on many of the same molecular components as other, better understood, invertebrate circadian clocks. Moreover, given the fact that horseshoe crabs have been extant for at least 450 million years (Rudkin and Young, 2009), these data might also provide some insight into the evolution of endogenous clocks.

In this study we identified 29 orthologs of circadian clock related proteins in the *L. polyphemus* transcriptome, corresponding to insect core clock proteins (PER, TIM, CLK, CYC and npCRY2), accessory proteins (VRI, CWO, SLIMBS, ARNT, CKIa, CKIe, CKIIa, CKIIB, JETLAG, SHAGGY, PPT1a, PPT1B, PPT2a-MTS, PPT2a-WBT, and PPT2a-TWS), input proteins (pCRY1), and output proteins (NPF, its receptor (NPFR), and PDHR). Two of the top 5 BLAST hits for *LpCYC1*, and 1 top hit for *LpCYC2* were labeled as “ARNT” proteins. BLASTp of the *LpnpCRY2* showed closest homology to proteins labeled pCRY1 and 1 protein labeled as generic “CRY” protein. Despite the mislabeling of these genes, phylogenetic evidence suggests that these BLAST hits are, in fact, orthologs of npCRY2 and CYC, and nest within the appropriate clades of the *cryptochrome* and *bHLH* gene families, respectively (data not shown). The mislabeling of the CYC and npCRY2 genes appears to be the result of an inappropriate reference genome during genome annotation (using a mammalian reference instead of an insect). Only two proteins found in insect circadian systems were not identified in the *L. polyphemus* transcriptome (PDP1e, and PDH). Even though we were unable to identify PDH in the *L. polyphemus* transcriptome, the presence of its receptor (PDHR) may imply that *LpPDH* is likely expressed in the *L. polyphemus* CNS. The absence of *LpPDH* from the transcriptome is possibly due to the relatively small size of the preprohormone (*D. melanogaster* - 102 amino acids – Renn et al., 1999; Crayfish - deKleijn et al., 1993). Among the proteins that were identified, the full-length protein sequences were able to be determined for all but *LppCRY1* and *LpNPFR*, and *LpPDHR*.

The identification of both pCRY1 and npCRY2 orthologs allows us to infer some additional information regarding the architecture of the circadian clock in *L. polyphemus*, based on what is known about insect clocks. Insect clock systems have been categorized into three types: 1) type 1, which contains *cry1* but lacks *cry2* (such as in *D. melanogaster melanogaster* (Hardin, 2005)); 2) type 2, which contains both *cry1* and *cry2* (such as in the monarch butterfly, *Danaus plexipus* (Zhu et al., 2005)) and; 3) type 3, which contains *cry2* but lacks *cry1* (such as in the honeybee, *Apis mellifera*; (Yuan et al., 2007)). The presence of both *cry1* and *cry2* in the *L. polyphemus* transcriptome would classify *L. polyphemus* as having a type 2 clock system which is considered an “ancestral clock system” (Yuan et al., 2007), which is fitting because *L. polyphemus* is considered to be a “living fossil” (Rudkin and Young, 2009). Thus, the circadian system of *L. polyphemus* is most likely organized more like the butterfly and mosquito (Yuan et al., 2007; Zhu et al., 2005), than *D. melanogaster* (Hardin, 2005) or *Apis mellifera* (Yuan et al., 2007).

4.2. Conservation and evolution of circadian proteins. The conservation of the various circadian clock proteins ranged from poor to highly conserved. Nearly all genes showed higher levels of conservation between *D. melanogaster* and *D. pulex* than orthologs of either species did to *L. polyphemus* orthologs. These findings are consistent with the evolutionary relationships of these three subphyla, with insects and crustaceans being more closely related to one another than to chelicerata (Giribet and Edgecombe, 2012). Conservation within functional domains tended to be much higher, in general, indicating that the functions of these proteins within the circadian clock may be well conserved between insects and chelicerata, as well as Crustacea and chelicerata. As such, the proteins which makeup the circadian clock in *D. melanogaster* may be relatively well conserved across the entire phylum Arthropoda. Interestingly, several domains (PeriodC domain of all three *LpPER* paralogs, the Timeless domain of *LpTIM*, and the LRR domains of *LpJET*) did show surprisingly low levels of conservation. The lack of conservation in these functional domains may confer novel functions of these proteins in the circadian clock of *L. polyphemus*. For example, the PeriodC domain is a known DNA-binding domain, and the overabundance of mutations in this domain may facilitate new transcriptional targets in the clock cells. Similarly the Timeless and LRR domains are found to be involved in protein-protein interactions, and the high level of mutations in this domain could allow for previously unknown interactions, thus allowing for new clock mechanisms.

Most (72%) *L. polyphemus* proteins (19 out of the 26, [pCRY1, NPFR, and PDHR were excluded due to incomplete sequences]) showed higher homology to *D. pulex* orthologs than to *D. melanogaster* orthologs. Six proteins (CYC1, CYC2, CKI α , CKII α , JET, and PP2 α MTS) show higher levels of sequence identity to *D. melanogaster* orthologs, and one gene (CKII β) showed equal levels of sequence identity to both *D. melanogaster* and *D. pulex*. The finding that of 72% of genes share higher sequence identity with *D. pulex* may be due to the fact that *D. pulex* and *L. polyphemus* evolved in similar marine habitats and thus experienced similar environmental pressures. There were six *L. polyphemus* genes, which show higher homology to *D. melanogaster* orthologs than to *D. pulex*. Most of these genes show low levels of conservation in the *D. pulex* orthologs, which may suggest that these genes may have undergone rapid evolution in the crustacean lineage.

Several gene duplications were identified from the *L. polyphemus* transcriptome. The two duplications of the *period* gene in *L. polyphemus* appear to be a novel discovery in invertebrates, and while there have been several duplications of the *period* gene in vertebrates (Bae et al., 2001), phylogenetic evidence suggests that the duplications this study revealed are unique to the chelicerate lineage, and independent of the vertebrate duplication events. These three paralogs raise an interesting question regarding the possibility of functional divergence following the duplication. Such differences have been observed between two period alleles in *D. melanogaster*, where each allele has different thermokinetics and varies by latitude (Sawyer et al., 1997). More extreme functional divergences have also been observed following the duplications of the *period* genes during vertebrate evolution (Bae et al., 2001). The single duplication events of the *cycle*, *arnt*, and *slimb* genes also appear to be unique and previously unknown duplications. These results present intriguing opportunities for potential novel mechanisms and functions within the clock systems of *L. polyphemus*. Are these different protein isoforms differentially expressed in different cell-types? Could these duplicated genes have been co-opted to create a clock controlling circatidal rhythms in *L. polyphemus*? Since these duplications do not appear in other chelicerates, such as scorpions and ticks, it is tempting to consider that they might be part of a clock system controlling these rhythms.

The various gene duplications also support the recent discovery of a whole genome duplication event based on genetic mapping of the *L. polyphemus* genome (Nossa et al., 2014). This genome duplication event appears to have taken place roughly 300 million years ago (Nossa et al., 2014), well after the divergence of *Limulidae* from the other chelicerate families (~450 million years ago (Rudkin and Young, 2009)). This is consistent with our findings, as none of the observed duplications have been found in other species outside the family *Limulidae*. Given the evidence of at least one genome duplication event, what is even more intriguing

than the presence of these duplications is the absence of duplications of the other circadian clock genes. The absence of these duplicate genes following a genome duplication event can have one of two explanations: 1) these genes underwent a loss of function, resulting in the duplicate genes being deleted from the genome or rendered inert, or 2) these genes underwent a functional divergence, resulting in these genes not being expressed in the CNS. In either case we would not expect to find transcripts of these duplicates in our transcriptome assembly.

4.3. Implications and future research. This study provides the first insight into the molecular mechanisms underlying the biological clocks of *L. polyphemus*, a member of the subphylum chelicerata. The conservation of these proteins, particularly within functional domains, provides evidence that some or all of these proteins may be involved in the circadian clock, and possibly the circatidal clock, of *L. polyphemus*. Immunolabelling of putative circadian proteins has been instrumental in localizing circadian clock cells (Siwicki et al., 1988) and their output cells in a variety of insects (Sehadova et al., 2004; Shao et al., 2006) and this study provides the foundation for similar studies in *L. polyphemus*. This study also provides a resource to aid in the development of qPCR primers and probes for Southern blotting and *in situ* hybridization. Moreover, the identification of these genes may also allow us to begin to tease apart the mechanisms underlying circatidal/circalunidian clocks (Chabot et al., 2004; Chabot and Watson, 2010), the molecular bases of which are unknown even in model systems (Tessmar-Raible et al., 2011).

Biological clocks act as master regulators of the physiology and behavior of most organisms. The high level of conservation of circadian clock related genes in *Limulus* indicates that they may play a role in the biological clock system(s) of this species, thus it is likely that the basic architecture of the circadian clock found in insects dates back prior to the divergence of chelicerata on the arthropod phylogenetic tree.

Acknowledgments.

The authors thank Scott Tighe, Ramiro Barrantes, and Tim Hunter of the Vermont Cancer Center for help with the extraction and RNAseq runs and Kelley Thomas and Kazu Okamoto of the Hubbard Center for Genome Studies, UNH for help with the assembly. This work was supported by NSF IOS 0920342 to CCC and WHW and the New Hampshire IDEa Network of Biomedical Research Excellence (NH-INBRE) with an Institutional Development Award (IDEA) from the National Institute of General Medical Sciences of the National Institutes of Health: Grant number P20GM103506. The funders had no role in the design, data collection, and analysis, decision to publish, or preparation of the manuscript. The authors state that no competing interests exist.

5. References.

- Allada, R., Chung, B.Y., 2010. Circadian organization of behavior and physiology in *Drosophila*. *Annu Rev Physiol* 72, 605-624.
- Bae, K., Jin, X., Maywood, E.S., Hastings, M.H., Reppert, S.M., Weaver, D.R., 2001. Differential functions of mPer1, mPer2, and mPer3 in the SCN circadian clock. *Neuron* 30, 525-536.
- Barlow, R.B., 1983. Circadian rhythms in the *Limulus* visual system. *The Journal of Neuroscience* 3, 856-870.
- Barlow, R.B., Jr., Chamberlain, S.C., Levinson, J.Z., 1980. *Limulus* brain modulates the structure and function of the lateral eyes. *Science* 210, 1037-1039.
- Battelle, B.A., 2013. What the clock tells the eye: lessons from an ancient arthropod. *Integr Comp Biol* 53, 144-153.
- Chabot, C.C., C., R.-P.N., W.H., W., 2016. Circalunidian clocks control tidal rhythms of locomotion in the American horseshoe crab, *Limulus polyphemus*. *Marine and Freshwater Behaviour and Physiology*, 75-91.
- Chabot, C.C., Kent, J., Watson, W.H., 2004. Daily, circadian and tidal rhythms of locomotor activity in the horseshoe crab *Limulus polyphemus*. *Biological Bulliton*, 72-75.

Chabot, C.C., Watson, W.H., 2010. Circatidal rhythms of locomotion in the American horseshoe crab *Limulus polyphemus*: Underlying mechanisms and cues that influence them. *Curr Zool* 56, 499-517.

Chang, D.C., Reppert, S.M., 2003. A novel C-terminal domain of drosophila PERIOD inhibits dCLOCK:CYCLE-mediated transcription. *Curr Biol* 13, 758-762.

Christie, A.E., Fontanilla, T.M., Nesbit, K.T., Lenz, P.H., 2013. Prediction of the protein components of a putative *Calanus finmarchicus* (Crustacea, Copepoda) circadian signaling system using a de novo assembled transcriptome. *Comp Biochem Physiol Part D Genomics Proteomics* 8, 165-193.

de la Iglesia, H.O., Hsu, Y.W., 2010. Biological clocks and rhythms in intertidal crustaceans. *Front Biosci (Elite Ed)* 2, 1394-1404.

Depetris-Chauvin, A., Berni, J., Aranovich, E.J., Muraro, N.I., Beckwith, E.J., Ceriani, M.F., 2011. Adult-specific electrical silencing of pacemaker neurons uncouples molecular clock from circadian outputs. *Curr Biol* 21, 1783-1793.

Dunlap, J.C., 1999. Molecular bases for circadian clocks. *Cell* 96, 271-290.

Emery, P., So, W.V., Kaneko, M., Hall, J.C., Rosbash, M., 1998. CRY, a *Drosophila* clock and light-regulated cryptochrome, is a major contributor to circadian rhythm resetting and photosensitivity. *Cell* 95, 669-679.

Giribet, G., Edgecombe, G.D., 2012. Reevaluating the arthropod tree of life. *Annu Rev Entomol* 57, 167-186.

Hardin, P.E., 2005. The circadian timekeeping system of *Drosophila*. *Curr Biol* 15, R714-722.

Hermann, C., Yoshii, T., Dusik, V., Helfrich-Forster, C., 2012. Neuropeptide F immunoreactive clock neurons modify evening locomotor activity and free-running period in *Drosophila melanogaster*. *J Comp Neurol* 520, 970-987.

Lee, G., Bahn, J.H., Park, J.H., 2006. Sex- and clock-controlled expression of the neuropeptide F gene in *Drosophila*. *Proc Natl Acad Sci U S A* 103, 12580-12585.

Letunic, I., Doerks, T., Bork, P., 2015. SMART: recent updates, new developments and status in 2015. *Nucleic Acids Res* 43, D257-260.

Luo, R., Liu, B., Xie, Y., Li, Z., Huang, W., Yuan, J., He, G., Chen, Y., Pan, Q., Liu, Y., Tang, J., Wu, G., Zhang, H., Shi, Y., Liu, Y., Yu, C., Wang, B., Lu, Y., Han, C., Cheung, D.W., Yiu, S.M., Peng, S., Xiaoqian, Z., Liu, G., Liao, X., Li, Y., Yang, H., Wang, J., Lam, T.W., Wang, J., 2012. SOAPdenovo2: an empirically improved memory-efficient short-read de novo assembler. *Gigascience* 1, 18.

Naylor, E., 1996. Crab clockwork: the case for interactive circatidal and circadian oscillators controlling rhythmic locomotor activity of *Carcinus maenas*. *Chronobiol Int* 13, 153-161.

Nossa, C.W., Havlak, P., Yue, J.X., Lv, J., Vincent, K.Y., Brockmann, H.J., Putnam, N.H., 2014. Joint assembly and genetic mapping of the Atlantic horseshoe crab genome reveals ancient whole genome duplication. *Gigascience* 3, 9.

Palmer, J.D., 1973. Tidal rhythms: the clock control of the rhythmic physiology of marine organisms. *Biological Review* 48.

Palmer, J.D., Williams, B.G., 1986. Comparative studies of tidal rhythms II. The dual clock control of the locomotor rhythms of two decapod crustaceans. *Marine behaviour and physiology*, 269-278.

Renn, S.C.P., Park, J.H., Rosbash, M., Hall, J.C., Taghert, P.H., 1999. A pdf neuropeptide gene mutation and ablation of PDF neurons each cause severe abnormalities of behavioral circadian rhythms in *Drosophila*. *Cell* 99, 791-802.

Rubin, E.B., Shemesh, Y., Cohen, M., Elgavish, S., Robertson, H.M., Bloch, G., 2006. Molecular and phylogenetic analyses reveal mammalian-like clockwork in the honey bee (*Apis mellifera*) and shed new light on the molecular evolution of the circadian clock. *Genome Res* 16, 1352-1365.

Rudkin, D.M., Young, G.A., 2009. Horseshoe Crabs - An Ancient Ancestry Revealed. *Biology and Conservation of Horseshoe Crabs*, 25-44.

Sawyer, L.A., Hennessy, J.M., Peixoto, A.A., Rosato, E., Parkinson, H., Costa, R., Kyriacou, C.P., 1997. Natural variation in a *Drosophila* clock gene and temperature compensation. *Science* 278, 2117-2120.

Schultz, J., Milpetz, F., Bork, P., Ponting, C.P., 1998. SMART, a simple modular architecture research tool: Identification of signaling domains. *P Natl Acad Sci USA* 95, 5857-5864.

- Sehadova, H., Markova, E.P., Sehnal, F.S., Takeda, M., 2004. Distribution of circadian clock-related proteins in the cephalic nervous system of the silkworm, *Bombyx mori*. *J Biol Rhythm* 19, 466-482.
- Shao, Q.M., Sehadova, H., Ichihara, N., Sehnal, F., Takeda, M., 2006. Immunoreactivities to three circadian clock proteins in two ground crickets suggest interspecific diversity of the circadian clock structure. *J Biol Rhythm* 21, 118-131.
- Siwicki, K.K., Eastman, C., Petersen, G., Rosbash, M., Hall, J.C., 1988. Antibodies to the Period Gene-Product of *Drosophila* Reveal Diverse Tissue Distribution and Rhythmic Changes in the Visual-System. *Neuron* 1, 141-150.
- Stoleru, D., Peng, Y., Agosto, J., Rosbash, M., 2004. Coupled oscillators control morning and evening locomotor behaviour of *Drosophila*. *Nature* 431, 862-868.
- Strauss, J., Zhang, Q., Verleyen, P., Huybrechts, J., Neupert, S., Predel, R., Pauwels, K., Dircksen, H., 2011. Pigment-dispersing hormone in *Daphnia* interneurons, one type homologous to insect clock neurons displaying circadian rhythmicity. *Cell Mol Life Sci* 68, 3403-3423.
- Taghert, P.H., Shafer, O.T., 2006. Mechanisms of clock output in the *Drosophila* circadian pacemaker system. *J Biol Rhythm* 21, 445-457.
- Takekata, H., Numata, H., Shiga, S., Goto, S.G., 2014. Silencing the circadian clock gene Clock using RNAi reveals dissociation of the circatidal clock from the circadian clock in the mangrove cricket. *J Insect Physiol* 68, 16-22.
- Tauber, E., Last, K.S., Olive, P.J.W., Kyriacou, C.P., 2004. Clock gene evolution and functional divergence. *J Biol Rhythm* 19, 445-458.
- Tessmar-Raible, K., Raible, F., Arboleda, E., 2011. Another place, another timer: Marine species and the rhythms of life. *Bioessays* 33, 165-172.
- Tilden, A.R., McCoole, M.D., Harmon, S.M., Baer, K.N., Christie, A.E., 2011. Genomic identification of a putative circadian system in the cladoceran crustacean *Daphnia pulex*. *Comp Biochem Phys D* 6, 282-309.
- Tomioka, K., Matsumoto, A., 2010. A comparative view of insect circadian clock systems. *Cell Mol Life Sci* 67, 1397-1406.
- Watson, W.H., Bedford, L., Chabot, C.C., 2008. Rhythms of locomotion expressed by *Limulus polyphemus*, the American horseshoe crab: II. Relationship to circadian rhythms of visual sensitivity. *Biol Bull-US* 215, 46-56.
- Yuan, Q., Metterville, D., Briscoe, A.D., Reppert, S.M., 2007. Insect cryptochromes: Gene duplication and loss define diverse ways to construct insect circadian clocks. *Mol Biol Evol* 24, 948-955.
- Zhang, E.E., Kay, S.A., 2010. Clocks not winding down: unravelling circadian networks. *Nature review*, 764-776.
- Zhu, H.S., Yuan, Q., Briscoe, A.D., Froy, O., Casselman, A., Reppert, S.M., 2005. The two CRYs of the butterfly. *Current Biology* 15, R953-R954.

Table 1: Top 5 BLASTp hits from UniProt. † indicates partial sequence..

Proteins	UniProt KB Protein Database				E-value	BLAST score
Core Proteins	UniProt ID	Species	Protein description			
PERIODA (PERA)						
LpPERA	H7C8F3	<i>Apteronomobius asahinai</i>	PERIOD, isoform 1	1e-121	1,058	
	H7C8F4	<i>Apteronomobius asahinai</i>	PERIOD, isoform 2	2e-121	1,054	
	B7PDL3	<i>Ixodes scapularis</i>	Period circadian protein, putative	4e-118	1,013	
	Q8MMG2	<i>Blattella germanica</i>	Circadian clock protein PERIOD	1e-114	1,007	
	L0CAI9	<i>Rhyarobia maderae</i>	Period	2e-114	1,009	
PERIODB (PERB)						
LpPERB	B7PDL3	<i>Ixodes scapularis</i>	Period circadian protein, putative	6e-119	1,020	
	Q8MMG2	<i>Blattella germanica</i>	Circadian clock protein PERIOD	8e-118	1,031	
	A1EA98	<i>Blattella bisignata</i>	Circadian clock protein period	2e-115	1,008	
	H7C8F3	<i>Apteronomobius asahinai</i>	PERIOD, isoform 1	2e-114	1,008	
	H7C8F4	<i>Apteronomobius asahinai</i>	PERIOD, isoform 2	4e-114	1,004	
PERIODC (PERC)						
LpPERC	H7C8F3	<i>Apteronomobius asahinai</i>	PERIOD, isoform 1	4e-133	1,142	
	H7C8F4	<i>Apteronomobius asahinai</i>	PERIOD, isoform 2	3e-132	1,134	
	B7PDL3	<i>Ixodes scapularis</i>	Period circadian protein, putative	2e-129	1,095	
	L0CAI9	<i>Rhyarobia maderae</i>	Period	9e-129	1,115	
	A1EA98	<i>Blattella bisignata</i>	Circadian clock protein period	5e-124	1,071	
TIMELESS (TIM)						
LpTIM	J9Y3V3	<i>Clunio marinus</i>	Timeless	2e-180	1,464	
	G9M9V8	<i>Thermobia domestica</i>	TIMELESS	4e-178	1,458	
	H2D5T9	<i>Aedes albopictus</i>	Timeless	6e-165	1,370	
	E0D5C1	<i>Gryllus bimaculatus</i>	TIMELESS	5e-164	1,359	
	Q05E95	<i>Aedes aegypti</i>	TIMELESS	2e-161	1,349	
CLOCK (CLK)						
LpCLK	E2C765	<i>Harpegnathos saltator</i>	Circadian locomotor output cycles protein kaput	4e-164	1,276	
	H9KJ84	<i>Apis mellifera</i>	Uncharacterized protein	3e-161	1,264	
	E2AY34	<i>Camponotus floridanus</i>	Circadian locomotor output cycles protein kaput	3e-159	1,251	
	F4WUR0	<i>Acromyrmex echinator</i>	Circadian locomotor output cycles protein kaput	1e-156	1,233	
	H9HL46	<i>Atta cephalotes</i>	Uncharacterized protein	4e-156	1,223	
CYCLE 1 (CYC1)						
LpCYC1	E0D6T3	<i>Thermobia domestica</i>	CYCLE	<1e180	1,786	
	A0MH07	<i>Lutzomyia longipalpis</i>	Cycle	<1e180	1,622	
	E2A3F0	<i>Camponotus floridanus</i>	Aryl hydrocarbon receptor nuclear translocator	<1e180	1,643	
	A9XCF1	<i>Tribolium castaneum</i>	CYCLE	<1e180	1,621	
	E2B7K2	<i>Harpegnathos saltator</i>	Aryl hydrocarbon receptor nuclear translocator	<1e180	1,620	
CYCLE 2 (CYC2)						
LpCYC2	E0D6T3	<i>Thermobia domestica</i>	CYCLE	<1e180	1,674	
	E2A3F0	<i>Camponotus floridanus</i>	Aryl hydrocarbon receptor nuclear translocator	<1e180	1,606	
	H9K2D4	<i>Apis mellifera</i>	CYC	<1e180	1,562	
	A0MH07	<i>Lutzomyia longipalpis</i>	Cycle	<1e180	1,555	
	A9XCF1	<i>Tribolium castaneum</i>	CYCLE	<1e180	1,560	
CRYPTOCHROME 2 (npCRY2)						
LpnpCRY2	E2BGC2	<i>Harpegnathos saltator</i>	Cryptochrome-1	<1e180	2,213	
	L7YAB3	<i>Solenopsis invicta</i>	Cryptochrome	<1e180	2,210	
	A8QSC1	<i>Bombus impatiens</i>	Cryptochrome 2 protein	<1e180	2,202	
	L0C8K6	<i>Rhyarobia maderae</i>	Cryptochrome 2	<1e180	2,199	
	F4WVC9	<i>Acromyrmex echinator</i>	Cryptochrome-1	<1e180	2,199	
Accessory Proteins						
VRILLE (VRI)						
LpVRI	E0VHE1	<i>Pediculus humanus</i>	Transcriptional factor nf13/e4bp4, putative	1e-43	424	
	B7PEG8	<i>Ixodes scapularis</i>	Transcriptional factor nf13/e4bp4, putative	9e-43	417	
	Q699T4	<i>Antheraea pernyi</i>	Vrille	6e-42	410	
	Q1XD36	<i>Danaus plexippus</i>	Vrille	1e-41	408	
	J9K7G7	<i>Acyrtosiphon pisum</i>	Uncharacterized protein	4e-40	403	
CLOCK WORK ORANGE (CWO)						
LpCWO	B7PGM4	<i>Ixodes scapularis</i>	Putative uncharacterized protein	1e-57	528	
	B4LW47	<i>Drosophila virilis</i>	GJ23560	2e-42	430	
	Q7QFU3	<i>Anopheles gambiae</i>	AGAP003844-PA	2e-41	422	
	E9GEU4	<i>Daphnia pulex</i>	Putative uncharacterized protein	7e-41	395	
	B4JIN1	<i>Drosophila grimshawi</i>	GH19128	2e-40	415	
SUPERNUMERARY LIMBS1 (SLIMB1)						
LpSLIMB1	L7M0T0	<i>Rhipicephalus pulchellus</i>	Uncharacterized protein	<1e180	2,159	
	E9IHL0	<i>Solenopsis invicta</i>	Putative uncharacterized protein	<1e180	2,159	
	D2XMQ7	<i>Saccoglossus kowalevskii</i>	Beta-TCRP E3 ligase	<1e180	2,145	
	H9HC45	<i>Atta cephalotes</i>	Uncharacterized protein	<1e180	2,142	
	D6WA15	<i>Tribolium castaneum</i>	Supernumerary limbs	<1e180	2,138	
SUPERNUMERARY LIMBS2 (SLIMB2)						
LpSLIMB2	L7M0T0	<i>Rhipicephalus pulchellus</i>	Uncharacterized protein	<1e180	2,153	
	E9HMX3	<i>Daphnia pulex</i>	Putative uncharacterized protein	<1e180	2,150	
	E9IHL0	<i>Solenopsis invicta</i>	Putative uncharacterized protein	<1e180	2,147	
	H3JK44	<i>Strongylocentrotus purpuratus</i>	Uncharacterized protein	<1e180	2,135	
	G3IS74	<i>Cricetulus griseus</i>	F-box/WD repeat-containing protein 1A	<1e180	2,133	
ARYL HYDROCARBON RECEPTOR NUCLEAR TRANSPORTER1 (ARNT1)						
LpARNT1	C7B7E8	<i>Litopenaeus vannamei</i>	Hypoxia inducible factor 1 beta	<1e180	1,799	
	E0VXW6	<i>Pediculus humanus</i>	Putative uncharacterized protein	<1e180	1,805	
	L7MH74	<i>Rhipicephalus pulchellus</i>	Putative tango	<1e180	1,770	
	D6W6H5	<i>Tribolium castaneum</i>	Putative uncharacterized protein	<1e180	1,739	
	Q1JUI4	<i>Daphnia magna</i>	Aryl hydrocarbone receptor nuclear translocator	<1e180	1,723	
ARYL HYDROCARBON RECEPTOR NUCLEAR TRANSPORTER2 (ARNT2)						
LpARNT2	E0VXW6	<i>Pediculus humanus</i>	Putative uncharacterized protein	<1e180	1,900	

	C7B7E8	<i>Litopenaeus vannamei</i>	Hypoxia inducible factor 1 beta	<1e180	1,858
	Q1JU14	<i>Daphnia magna</i>	Aryl hydrocarbone receptor nuclear translocator	<1e180	1,850
	Q1JU15	<i>Daphnia magna</i>	Aryl hydrocarbone receptor nuclear translocator	<1e180	1,838
	E9FQM5	<i>Daphnia pulex</i>	Putative aryl hydrocarbon receptor nuclear translocator	<1e180	1,838
CASIEN KINASE Ia (CKIa)					
LpCKIa	Q8JG73	<i>Danio rerio</i>	Casein kinase 1alpha S	<1e180	1,570
	Q75WS8	<i>Carassius auratus</i>	Casein kinase I alpha S	<1e180	1,570
	Q8JGT0	<i>Danio rerio</i>	Casein kinase 1, alpha I	<1e180	1,568
	Q75WS9	<i>Carassius auratus</i>	Casein kinase I alpha	<1e180	1,568
	B5THN0	<i>Saccoglossus kowalevskii</i>	Casein kinase 1 protein catalytic subunit	<1e180	1,567
CASIEN KINASE Iε (CKIε)					
LpXKIε	F6YKD5	<i>Equus caballus</i>	Uncharacterized protein	<1e180	1,422
	G1SGL5	<i>Oryctolagus cuniculus</i>	Uncharacterized protein	<1e180	1,418
	G1U4F8	<i>Oryctolagus cuniculus</i>	Uncharacterized protein	<1e180	1,418
	M7AVG9	<i>Chelonia mydas</i>	Casein kinase I isoform epsilon	<1e180	1,416
	H0V5Y3	<i>Cavia porcellus</i>	Uncharacterized protein	<1e180	1,416
CASIEN KINASE IIα (CKIIα)					
LpCKIIα	E2AW17	<i>Camponotus floridanus</i>	Casein kinase II subunit alpha	<1e180	1,674
	F4WU82	<i>Acromyrmex echinatio</i>	Casein kinase II subunit alpha	<1e180	1,666
	B7PVH2	<i>Ixodes scapularis</i>	Mitogen-activated protein kinase, putative	<1e180	1,671
	E2BH98	<i>Harpegnathos saltator</i>	Casein kinase II subunit alpha	<1e180	1,669
	K7GIG0	<i>Pelodiscus sinensis</i>	Uncharacterized protein	<1e180	1,665
CASIEN KINASE IIβ (CKIIβ)					
LpCKIIβ	Q71U52	<i>Cyprinus carpio</i>	CK2 beta subunit	5e-144	1,057
	Q6DEU1	<i>Xenopus tropicalis</i>	Casein kinase 2, beta polypeptide	5e-144	1,057
	Q1LXD2	<i>Danio rerio</i>	Casein kinase 2 beta	5e-144	1,057
	M4AI87	<i>Xiphophorus maculatus</i>	Uncharacterized protein	5e-144	1,057
	J3SBX9	<i>Crotalus adamanteus</i>	Casein kinase II subunit beta	5e-144	1,057
JETLAG (JET)					
LpJET	K1P9I2	<i>Crassostrea gigas</i>	F-box only protein 37	1e-37	354
	C3Y7V1	<i>Branchiostoma floridae</i>	Putative uncharacterized protein	8e-37	356
	C3ZZX8	<i>Branchiostoma floridae</i>	Putative uncharacterized protein	6e-35	342
	R7VDT7	<i>Capitella teleta</i>	Uncharacterized protein	1e-32	326
	A7SQL5	<i>Nematostella vectensis</i>	Predicted protein	1e-31	320
SHAGGY (SGG)					
LpSGG	E2C155	<i>Harpegnathos saltator</i>	Protein kinase shaggy	<1e180	1,478
	E9H6Q6	<i>Daphnia pulex</i>	Glycogen synthase kinase 3 beta	<1e180	1,475
	E9I8X0	<i>Solenopsis invicta</i>	Putative uncharacterized protein	<1e180	1,469
	G3MI16	<i>Amblyomma maculatum</i>	Putative uncharacterized protein	<1e180	1,453
	F4X3V9	<i>Acromyrmex echinatio</i>	Protein kinase shaggy	<1e180	1,467
PROTEIN PHOSPHATASE 1α (PP1α)					
LpPP1α	K1PXC6	<i>Crassostrea gigas</i>	Serine/threonine-protein phosphatase	<1e180	1,631
	A7RVJ0	<i>Nematostella vectensis</i>	Serine/threonine-protein phosphatase	<1e180	1,627
	E9HKA6	<i>Daphnia pulex</i>	Serine/threonine-protein phosphatase	<1e180	1,624
	H2L9G2	<i>Oryzias latipes</i>	Serine/threonine-protein phosphatase	<1e180	1,620
	H9KP59	<i>Apis mellifera</i>	Serine/threonine-protein phosphatase	<1e180	1,618
PROTEIN PHOSPHATASE 1β (PP1β)					
LpPP1β	L7MB61	<i>Rhipicephalus pulchellus</i>	Serine/threonine-protein phosphatase	<1e180	1,470
	H2UK23	<i>Takifugu rubripes</i>	Serine/threonine-protein phosphatase	<1e180	1,458
	H3DIM0	<i>Tetraodon nigroviridis</i>	Serine/threonine-protein phosphatase	<1e180	1,457
	H2MDU5	<i>Oryzias latipes</i>	Serine/threonine-protein phosphatase	<1e180	1,457
	G3P7R9	<i>Gasterosteus aculeatus</i>	Serine/threonine-protein phosphatase	<1e180	1,457
PROTEIN PHOSPHATASE 2α-mts (PP2α-mts)					
LpPP2α-mts	L7M975	<i>Rhipicephalus pulchellus</i>	Serine/threonine-protein phosphatase	<1e180	1,336
	B7QGY5	<i>Ixodes scapularis</i>	Serine/threonine-protein phosphatase	<1e180	1,335
	E0VKT2	<i>Pediculus humanus</i>	Serine/threonine-protein phosphatase	<1e180	1,333
	R4G8Q4	<i>Rhodnius prolixus</i>	Putative serine/threonine protein phosphatase	<1e180	1,330
	E9IAY8	<i>Solenopsis invicta</i>	Serine/threonine-protein phosphatase	<1e180	1,334
PROTEIN PHOSPHATASE 2α-WBT (PP2α-WBT)					
LpPP2α-WBT	L7M3M4	<i>Rhipicephalus pulchellus</i>	Putative serine/threonine protein phosphatase...	<1e180	2,137
	H9J639	<i>Bombyx mori</i>	Uncharacterized protein	<1e180	2,073
	E2AI01	<i>Camponotus floridanus</i>	Serine/threonine-protein phosphatase 2A 56 kDa regulatory subunit	<1e180	2,045
	H9K2X9	<i>Apis mellifera</i>	Uncharacterized protein	<1e180	2,044
	E0VNG0	<i>Pediculus humanus</i>	Serine/threonine-protein phosphatase 2A 56 kDa regulatory subunit	<1e180	2,041
PROTEIN PHOSPHATASE 2α-TWS (PP2α-TWS)					
LpPP2α-TWS	H9KM79	<i>Apis mellifera</i>	Uncharacterized protein	<1e180	2,081
	I3RJJ3	<i>Scylla paramamosain</i>	Protein phosphatase 2A regulatory subunit B	<1e180	2,074
	G3MLH6	<i>Amblyomma maculatum</i>	Putative uncharacterized protein	<1e180	2,070
	L7M886	<i>Rhipicephalus pulchellus</i>	Putative serine/threonine protein phosphatase	<1e180	2,063
	F4WRB8	<i>Acromyrmex echinatio</i>	Protein phosphatase PP2A 55 kDa regulatory subunit	<1e180	2,055
Input Pathway Protein					
CRYPTOCHROME 1 (CRY1)†					
LpCRY1	A2A264	<i>Dianemobius nigrofasciatus</i>	Cryptochrome	4e-132	1,022
	K1PRK6	<i>Crassostrea gigas</i>	Cryptochrome-1	3e-120	941
	R7V1J3	<i>Capitella teleta</i>	Uncharacterized protein	5e-119	937
	Q2TJN5	<i>Danaus plexippus</i>	Cryptochrome	5e-115	908
	J9T2S0	<i>Mythinna separata</i>	Cryptochrome 1	4e-112	888
Output Pathway Proteins					
NEUROPEPTIDE F RECEPTOR (NPFR)†					
LpNPFR	T1KR44	<i>Tetranychus urticae</i>	Uncharacterized protein	6e-96	759
	T1IW42	<i>Strigamia maritima</i>	Uncharacterized protein	7e-88	707

	B0WJ64	<i>Culex quinquefasciatus</i>	G-protein coupled receptor	3e-78	642
	Q16SC4	<i>Aedes aegypti</i>	AAEL010626-PA	5e-78	636
	T1JTP7	<i>Tetranychus urticae</i>	Uncharacterized protein	1e-77	642
NEUROPEPTIDE F (NPF)					
<i>LpNPF</i>	T1KF83	<i>Tetranychus urticae</i>	Uncharacterized protein	2e-11	156
	A0A087UNM1	<i>Stegodyphus mimosarum</i>	Uncharacterized protein	2e-7	128
	T1JM02	<i>Strigamia maritima</i>	Uncharacterized protein	7e-7	125
	F6KM62	<i>Litopenaeus vannamei</i>	Preproneuropeptide F I	0.0000013	123
	F6KM64	<i>Melicertus marginatus</i>	Preproneuropeptide F I	0.0000013	123
PIGMENT DISPERSING HORMONE RECEPTOR (PDHR)†					
<i>LpPDHR</i>	K7JRN8	<i>Nasonia vitripennis</i>	Uncharacterized protein	2e-63	548
	U3U904	<i>Nilaparvata lugens</i>	Neuropeptide GPCR B2	1e-61	545
	Q7PQE3	<i>Anopheles gambiae</i>	AGAP003654-PA	8e-61	537
	A0A0P5AQT3	<i>Daphnia magna</i>	Class b secretin g-protein coupled receptor	6e-61	533
	A0A0P5CML9	<i>Daphnia magna</i>	Class b secretin g-protein coupled receptor	1e-60	533

Table 2: Top BLASTp hits for *Drosophila melanogaster* and *Daphnia pulex*. † indicates partial sequence. * indicates Protein name which originated from *D. pulex* genome map on Fleabase.

Proteins	Species name	Protein name	UniProt ID	Blast Score	E-value
CORE PROTEINS					
PERa	<i>Drosophila melanogaster</i>	Isoform PER-E	P07663	735	2.0e-78
	<i>Daphnia pulex</i>	PER	E9GW67	822	1.0e-90
PERb	<i>Drosophila melanogaster</i>	Isoform PER-E	P07663	653	7.0e-68
	<i>Daphnia pulex</i>	PER	E9GW67	701	4.0e-75
PERc	<i>Drosophila melanogaster</i>	Isoform PER-E	P07663	765	5.0e-82
	<i>Daphnia pulex</i>	PER	E9GW67	876	1.0e-97
TIM	<i>Drosophila melanogaster</i>	Timeless, isoform G	B7Z007	1,290	3.0e-152
	<i>Daphnia pulex</i>	Putative TIMELESS/TIM-1 protein	E9FZ81	956	2.0e-109
CLK	<i>Drosophila melanogaster</i>	Clk CLOCK jrk PAS1	O61735	1,048	1.0e-125
	<i>Daphnia pulex</i>	CLOCK	E9GKD1	1,033	2.0e-126
CYC1	<i>Drosophila melanogaster</i>	Protein cycle	O61734	1,220	6.0e-159
	<i>Daphnia pulex</i>	CYCLE	E9FRH8	1,500	<1.0e-180
CYC2	<i>Drosophila melanogaster</i>	Protein cycle	O61734	1,227	6.0e-160
	<i>Daphnia pulex</i>	CYCLE	E9FRH8	1,470	<1.0e-180
npCRY2	<i>Drosophila melanogaster</i>	phr6-4	Q8SXX5	1,296	3.0e-171
	<i>Daphnia pulex</i>	CRY-M	E9GDJ9	2,139	<1.0e-180
Accessory Proteins					
VRI	<i>Drosophila melanogaster</i>	Vri	Q7KTN9	306	1.0e-28
	<i>Daphnia pulex</i>	Vri†	E9HB85	280	1.0e-28
CWO	<i>Drosophila melanogaster</i>	cwo-RA	B7FNP5	312	1.0e-29
	<i>Daphnia pulex</i>	Putative uncharacterized protein	E9GEU4	395	2.0e-42
SLIMB1	<i>Drosophila melanogaster</i>	Slimb	Q9VDE3	2,114	<1.0e-180
	<i>Daphnia pulex</i>	f-box/wd-repeat protein*	E9HMX3	2,136	<1.0e-180
SLIMB2	<i>Drosophila melanogaster</i>	Slimb	Q9VDE3	2,107	<1.0e-180
	<i>Daphnia pulex</i>	f-box/wd-repeat protein*	E9HMX3	2,150	<1.0e-180
ARNT1	<i>Drosophila melanogaster</i>	Aryl hydrocarbon receptor nuclear translocator homolog	O15945	1,568	<1.0e-180
	<i>Daphnia pulex</i>	Putative aryl hydrocarbon receptor nuclear translocator	E9FQM5	1,723	<1.0e-180
ARNT2	<i>Drosophila melanogaster</i>	Aryl hydrocarbon receptor nuclear translocator homolog	O15945	1,680	<1.0e-180
	<i>Daphnia pulex</i>	Putative aryl hydrocarbon receptor nuclear translocator	E9FQM5	1,838	<1.0e-180
CKIε	<i>Drosophila melanogaster</i>	Dco dbt	O76324	1,282	6.0e-173
	<i>Daphnia pulex</i>	Casein kinase i alpha*	E9FS31	1,171	3.0e-157
CKIα	<i>Drosophila melanogaster</i>	Casein kinase I isoform alpha	O76324	1,348	<1.0e-180
	<i>Daphnia pulex</i>	Casein kinase I alpha*	E9HGM4	1,526	<1.0e-180
CKIIα	<i>Drosophila melanogaster</i>	Casein kinase II alpha subunit, isoform C	P08181	1,590	<1.0e-180
	<i>Daphnia pulex</i>	Casein kinase ii subunit alpha*	E9GCV0	1,659	<1.0e-180
CKIIβ	<i>Drosophila melanogaster</i>	Casein kinase II beta subunit, isoform C	P08182-3	1,010	6.0e-137
	<i>Daphnia pulex</i>	Casein kinase ii subunit beta*	E9GTE8	1,033	6.0e-142
JET	<i>Drosophila melanogaster</i>	Jetlag, isoform B	Q0E8T8	188	3.0e-15
	<i>Daphnia pulex</i>	f-box/leucine rich repeat protein*	E9G1Z9	155	2.0e-10
SGG	<i>Drosophila melanogaster</i>	Isoform G of Protein kinase shaggy	P18431	1,357	<1.0e-180
	<i>Daphnia pulex</i>	Glycogen synthase kinase 3 beta	E9H6Q6	1,475	<1.0e-180
PP1α	<i>Drosophila melanogaster</i>	Pp1alpha-96A	P48461	1,575	<1.0e-180
	<i>Daphnia pulex</i>	Serine/threonine-protein phosphatase	E9HKA6	1,624	<1.0e-180
PP1β	<i>Drosophila melanogaster</i>	Serine/threonine-protein phosphatase beta isoform	H5V895	1,386	<1.0e-180
	<i>Daphnia pulex</i>	Serine/threonine-protein phosphatase	E9G7U7	1,427	<1.0e-180
PP2a-MTS	<i>Drosophila melanogaster</i>	mts PP2A	P23696	1,313	<1.0e-180
	<i>Daphnia pulex</i>	Serine/threonine-protein phosphatase	E9G8K4	842	5.0e-110
PP2-WBT	<i>Drosophila melanogaster</i>	wdb	Q9VB23	1,986	<1.0e-180
	<i>Daphnia pulex</i>	Serine/threonine-protein phosphatase kda regulatory subunit*	E9G2F2	1,932	<1.0e-180
PP2-TWS	<i>Drosophila melanogaster</i>	tw5 aar Pp2A-85F	P36872	2,003	<1.0e-180
	<i>Daphnia pulex</i>	TWS	E9GU66	1,974	<1.0e-180
INPUT PATHWAY PROTEINS					
CRY1†	<i>Drosophila melanogaster</i>	Cryptochrome-1	O77059	754	1.0e-93
	<i>Daphnia pulex</i>	CRY-D	E9GSJ7	754	9.0e-94
OUTPUT PATHWAY PROTEINS					
NPFR†	<i>Drosophila melanogaster</i>	Isoform 6 of Neuropeptide F receptor	Q9VNM1	583	1.0e-68
	<i>Daphnia pulex</i>	NPFG-protein-coupled receptor	E9GBE9	593	3.0e-71
NPF	<i>Drosophila melanogaster</i>	Neuropeptide F	Q9VET0	84	5.0e-3
	<i>Daphnia pulex</i>	Putative uncharacterized protein	E9GJI3	86	2.0e-3
PDHR†	<i>Drosophila melanogaster</i>	PDF receptor	Q9W4Y2	445	1.4 e-48
	<i>Daphnia pulex</i>	Putative PDF receptor variant 2	E9FR28	532	6.3e-62

530

531

Table 3: %Identity/%Similarity for *L. polyphemus* proteins against *Drosophila melanogaster* and *Daphnia pulex*. † indicates partial sequence.

Proteins		<i>Drosophila melanogaster</i>		<i>Daphnia pulex</i>	
Core proteins	Length	Domains (%Identity/%Similarity)	Total	Domains (%Identity/%Similarity)	Total
PERA	1,001	PASa(40/53);PASb(44/63);PAC(66/82);PERIODC(N/A)	26/34	PASa(40/50);PASb(45/66);PAC(66/84);PERIODC(21/30)	30/38
PERB	1,021	PASa(37/56);PASb(44/55);PAC(75/86);PERIODC(N/A)	25/31	PASa(38/57);PASb(49/58);PAC(70/80);PERIODC(20/29)	32/39
PERC	1,047	PASa(38/56);PASb(45/65);PAC(68/82);PERIODC(N/A)	25/32	PASa(43/57);PASb(49/61);PAC(73/82);PERIODC(22/31)	31/39
TIM	1,100	TIMELESS(35/45)	27/38	TIMELESS(25/40)	36/45
CLK	654	bHLH(67/82);PASa(52/61);PASb(78/82);PAC(75/93)	32/40	bHLH(71/84);PASa(55/66);PASb(76/85);PAC(84/91)	43/50
CYC1	623	bHLH(75/81);PASa(75/87);PASb(53/73);PAC(57/66)	62/68	bHLH(78/85);PASa(87/94);PASb(57/66);PAC(57/70)	31/37
CYC2	632	bHLH(76/81);PASa(76/88);PASb(53/66);PAC(52/66)	60/66	bHLH(76/85);PASa(85/93);PASb(59/68);PAC(57/70)	31/38
npCRY2	549	6-4 Photolyase(N/A);FAD Binding 7(N/A)	N/A	6-4 Photolyase(64/76);FAD Binding 7(84/91)	70/79
Accessory Proteins					
VRI	439	BRLZ(69/80)	20/28	BRLZ(68/75)	42/43
CWO	536	bHLH(82/86);ORANGE(32/39)	16/22	bHLH(88/95);ORANGE(34/46)	23/25
SLIMB1	533	b-TrCP D(55/65);FBOX(85/93);WD40(92/92);WD40(84/98);WD40(97/97);WD40(94/94);WD40(97/100);WD40(92/97);WD40(95/95)	73/78	b-TrCP D(53/65);FBOX(75/85);WD40(92/95);WD40(84/95);WD40(87/87);WD40(95/97);WD40(95/100);WD40(89/97);WD40(100/100)	74/80
SLIMB2	528	b-TrCP D(57/68);FBOX(88/93);WD40(92/92);WD40(84/89);WD40(97/97);WD40(95/95);WD40(97/100);WD40(92/97);WD40(92/95)	74/79	b-TrCP D(57/68);FBOX(78/85);WD40(92/92);WD40(84/95);WD40(87/87);WD40(95/97);WD40(95/100);WD40(89/97);WD40(97/100)	76/82
ARNT1		bHLH(76/81);PASa(74/90);PASb(74/83);PAC(61/75)	49/57	bHLH(76/81);PASa(74/88);PASb(75/78);PAC(84/93)	52/60
ARNT2		bHLH(89/91);PASa(76/90);PASb(75/84);PAC(57/70)	49/57	bHLH(89/91);PASa(78/88);PASb(74/78);PAC(84/91)	54/59
CK Ia	402	STKc(78/84)	79/84	STKc(64/71)	46/52
CK Iε		STKc(75/81)	53/58	STKc(73/77)	64/67
CK IIα	329	STKc(92/96)	88/92	STKc(94/98)	86/90
CK IIβ	198	CKIIβ(90/93)	84/86	CKIIβ(93/95)	84/86
JET	234	LRR(12/35);LRR(15/41);LRR(31/50)	41/48	LRR(15/38);LRR(33/48);LRR(31/46)	14/19
SGG	364	STKc(77/85)	50/57	STKc(83/88)	69/75
PP1α	330	PP2Ac(94/97)	88/91	PP2Ac(96/99)	90/93
PP1β	274	PP2Ac(86/88)	76/79	PP2Ac(88/89)	81/82
PP2αMTS	296	PP2Ac(79/82)	80/84	PP2Ac(54/65)	51/61
PP2αWBT	494	B56(84/91)	71/78	B56(81/89)	74/82
PP2αTWS	443	WD40(85/97);WD40(90/93);WD40(93/98);WD40(93/95);WD40(87/90);WD40(93/94);WD40(87/92)	73/79	WD40(79/87);WD40(88/100);WD40(93/98);WD40(80/85);WD40(87/92);WD40(87/89);WD40(82/89)	84/91
Input Pathway Proteins					
CRY1	313†	FAD Binding 7(47/55)	N/A	FAD Binding 7(45/60)	N/A
Output Pathway Proteins					
NPFR	213†	7tM GPCR Srsx(53/66)	N/A	7tM GPCR Srsx(49/61)	N/A
NPF	95	PAH(36/54)	23/34	PAH(41/54)	29/37
PDHR	231†	7TM-2(40/57)	N/A	7TM-2(46/63)	N/A

[illegible]

Figure 1: bHLH, PASa, PASb and PAC domains region of putative *L. polyphemus* CYCLE1 (CYC1) protein. Alignment of *L. polyphemus* (Lim) CYC1 to *Daphnia pulex* (Daph) CYC and *Drosophila melanogaster* (Dros) CYC using CLUSTAL Omega. Symbols immediately below sequence indicates functional similarity of residues between protein orthologs. “*” Indicates identical residues, “:” Indicates residues with strongly similar properties, “.” Indicates residues with weakly similar properties. Blue shows bHLH domain, light green shows PAS domains, Red shows PAC domain. For full alignment see Supplemental Figure 4.

CYCLE 1 (CYC1) vs. CYCLE 2 (CYC2)

```

L. polyphemus ___CYC1      MDVGMGLDFNMTLKRKINEINDASDMEDED---MKVQKLESEQTKRHHSEIEKRRRD
L. polyphemus ___CYC2      MDIDMGLDFNMDTLKRKIMDIXXXXXXXXXXXXXXXXXXXXXXXXXXXXXXRRHHSEIEKRRRD
                             **: *****:***** :* *****

L. polyphemus ___CYC1      KMNTYITELSSMVPCKAMSRKLDKLTVLRMAVQHMKTLRINSYTEGHYKPSFLSDEELK
L. polyphemus ___CYC2      KMNTYISELSSMVPCKAMSRKLDKLTVLRMAVQHMKTLRINSYTEGHYKPSFLSDEELK
                             *****:*****:*****:*****:*****:*****:*****

L. polyphemus ___CYC1      YLILEAAEGFLFVVSQCDRCRLFVSESVSQILNYSQGDLLGQSWFDILHPKDIKVKKEQL
L. polyphemus ___CYC2      YLILEAAEGFLFVVSQCDRCRLFVSESVSQILNYSQGDLLGQSWFDILHPKDIKVKKEQL
                             *****:*****:*****:*****:*****:*****:*****

L. polyphemus ___CYC1      SSSDILPKERLIDAKTMLPVKTDMPGQSRLCPGSRRSFFCRMKCRSMPTVKEEADTTTG
L. polyphemus ___CYC2      SSSDILPKERLIDAKTMLPVKTDMPGQSRLCPGSRRSFFCRMKCRSLATVKEEADTTTG
                             *****:*****:*****:*****:*****:*****:*****

L. polyphemus ___CYC1      CHRKRKSQS-SDRKYLVIHCTGYLKSAPAKINLQEETDSDGESCNLSCLVAVGRVHPEI
L. polyphemus ___CYC2      CHKRKSQSSTDRKYLVIHCTGYLKSAPAKINLQEETDSDGESCNLSCLVAVGRVHPEI
                             **:*****:*****:*****:*****:*****:*****:*****

L. polyphemus ___CYC1      LHFVFPQGLEVRPFEFVSRHAMDCKFLYVDORATIMLGYPQELLGTSFYEYCHQDDIS
L. polyphemus ___CYC2      LHFVFPQGLEVRPFEFVSRHAMDCKFLYVDORATIMLGYPQELLGTSFYEYCHQDDIS
                             ****:*****:*****:*****:*****:*****:*****

L. polyphemus ___CYC1      HLADSHKQVLTNERLTIQYRFRTEKGSFICMQSKWKNPKNPWTKFFEYLVAVNPIIET
L. polyphemus ___CYC2      HLADSHKQVLTANEKITIQYRFRTEKGSFVCMQSTWKNFKNPWTKFFEYLVAVNSSVPY
                             *****:*****:*****:*****:*****:*****:*****

L. polyphemus ___CYC1      SHDPSSAPDVAVVESNATLEEMLNNSNSLDAVLSAFPTSSTASTSDTIQKFLSTRVG
L. polyphemus ___CYC2      SRDSCSAANLAVVESNATLEEMLNNSNSLDAVLVFPSTSDASTSDTIQKFLSTRVG
                             **:*.**.:*****:*****:*****:*****:*****:*****

L. polyphemus ___CYC1      ASKIGRQIADAEAMEVQTRDSSASNSFPVPFDNNVGLPNRTQLLVNSSIEDQTNLIPGPS
L. polyphemus ___CYC2      ACKIGQQIADAEAMEVQTRDSSASNSFPVPFDNNVGLPNRTQLLVNSSIENQATAVPGPS
                             *:*****:*****:*****:*****:*****:*****:*****

L. polyphemus ___CYC1      MASMTFVEFQHLFVPMNGDVRSVSPSQSCHSNQSQYSGTFHNSSVSDPDIDIMDTLMGRD
L. polyphemus ___CYC2      TAPMTTVEFNHVPVMNGDACPHSPSQSCHSNQSQYSGTFHNSSVSDPDIDFMDTLMGRD
                             * ** *:*****:*****:*****:*****:*****:*****

L. polyphemus ___CYC1      MLSVSYQNNSECNDEAAMAVIMSLEADAGLG
L. polyphemus ___CYC2      IISSTYQS-SNEGNDFAAMAVIMSLEADAGLG
                             **:*.**.:*****:*****:*****:*****:*****:*****

```

Figure 2: Putative *L. polyphemus* CYCLE 1 (CYC1) protein aligned against putative CYCLE 2 (CYC2) using CLUSTAL Omega. Symbols immediately below sequence indicates functional similarity of residues between protein orthologs. “*” indicates identical residues, “:” indicates residues with strongly similar properties, “.” indicates residues with weakly similar properties. Light green shows bHLH domain, Blue shows PAS domain, Red shows PAC domain.

PERIODa (PASa, PASb and PAC domains)

```

Dros_PER      TGVGAAGAAGTCGRGERVVKEDSFCCVISMHDGIVLYITTPSIDVGLYPRDMWLGRSFDIFV
Lim_PerA     -S--LSLKAVKHELDTKVKEGGFSV:ISLLDGTTHVTSSVTFALGYFEDFVLGGQCLTNFI
Tx_PER       -S--SAFKDLKNESECVKSGDGFCLAVSLQDGTIVYVSSSTIPILGYSKDMLLQGCTMFLN
Daph_PER     LK--KTFARHLEEVPTVKSKEAFIAATISLDQGVIVVYTPSLTEKLGYPVDMWCGRSLLDFI
              :      :      :      :      :      :      :      :      :      :
Dros_PER      HLKDRATFASQITTTGTLIAESR-----GSVP-----KDAKSTFCVMLLR
Lim_PerA     HPKDRVTFLSHLTCEGLNRFVEKKN-----GGFG-----NGQQTAFYCRIRQ
Tx_PER       YPRDRTTFANHLSGGLNRFENDAK-----GMCH-----NRSQSTFLCLRRLQ
Daph_PER     HPKDRLAFTNQITSKILASLDKDDSSSGYSFSSDEGLFPRGSMIGSSSSPPNMLCCRRLM
              :      :      :      :      :      :      :      :      :      :
Dros_PER      YRGLKSGGFGVIGRPVSVEFFRLGLTFREAPEEARP----DNY-----MVS
Lim_PerA     YNBLK-MGFEISNNKKPQYKFPSFTFHLKDLVNDTDSVE-----E
Tx_PER       YQSLK--FGYGISDKVKQYKFPQMSVYIKDVIDVDSVE-----N
Daph_PER     YRGLRTSGFGIVDKKTSYLPFKIILNLEKITLPRESPESQNDKETVAEGSSVPKNPEEDDS
              *.*:      :      :      :      :      :      :      :      :
Dros_PER      NGTNMMLVICATPIKSSYKVPDFEILSQKSPKTAIRHTATGIISHIVSAAVSA.GYLQDIL
Lim_PerA     SNHSSCLIATIPVICISAYKVPEEVPAM--TTPSTRHTSSCFESHIDERAVPIGYLPQDI
Tx_PER       ANIAMLIVITAVISITAYRVPNSLPAM--TCEPTRHTASCFESHVDLSAAVLYGLYLPQDI
Daph_PER     DSAEYVLLAYAVPISTAYKNEDPKNST--GETGLRHISANCLFEVDMSSVPIGLHLPQDI
              :      :      :      :      :      :      :      :      :      :
Dros_PER      LGRSIMDFYHHEDLSVMKETYETVMMKKGQIAGASFCSKPYRFLIQNGCVLLETETWTSFV
Lim_PerA     LIGKSVPEFFHYSQDLPLQKRDVLYELVINE-----QGHSFWSKPYRFAVLNGCFILETETWSCFI
Tx_PER       LGHSVDFEYFCMEDLSQLQDLQYELVINE-----QGHSFWSKPYRFAKNGSVILETETWSCFI
Daph_PER     LGTSALDFYHPNDLPBLKKIYDSVIGR-----QKGSLRSKPYNFAENGCGVILQTDWTQCFV
              :      :      :      :      :      :      :      :      :      :
Dros_PER      NPWSRKLEFVVGHERVVGQKQCNVFEAAPTCKLKI--SEEAQMSNRTRIKEDIVKRLAE
Lim_PerA     NPWSRKLEFVVGHERVVGQKQCNVFEAGNREELVA--SSEDCMKRSQRIQDDIRHILSQ
Tx_PER       NPWTRKLEFVVGQFVRVIGRKTIPNVFMEASRDNPETRPPEFVLNNNRHLQEQTKDILSQ
Daph_PER     NPWTKKLEFVAGHERVVGKGSNDPVDVFNPEGCVPFVDSPTAAENTKKLHAEICITLGT
              ***.*****.*****.*****.*****.*****.*****.*****.*****.

```

Figure 3: PASa, PASb and PAC domains region of putative *L. polyphemus* PERIOD (PER) protein. Alignment of *L. polyphemus* (Lim) PER to *Daphnia pulex* (Daph) PER, *D. melanogaster* (Dros) PER, *Ixodes scapularis* (Ix) PER using CLUSTAL Omega. Symbols immediately below sequence indicates functional similarity of residues between protein orthologs. “*” Indicates identical residues, “:” Indicates residues with strongly similar properties, “.” Indicates residues with weakly similar properties. Light green shows PAS domain, Red shows PAC domain. For full alignment see Supplemental Figure 1.

563

TIMELESS (TIMELESS domain)

```

Dros_TIM      MSRVRLHNNHIWNNQNFQKSVMDWLLATPQ-LYSAFSSLGCLEGDTYVVRNPALATLE
Lim_TIM       -----MVQESGMNMLKVKESELE
Ix_TIM        -----
Daph_TIM-1    -----MDHWIKLSGEVPCLSLIPLGSFHNKCYCTDPECLNNLK

Dros_TIM      EINYKLTIEDQTLRTFRRAIGFGQNVRSDLIPLLENAK-DDAVLESVIRILVNLTVPVEC
Lim_TIM       TMEQELHQBELGSRNVRLSFGMSCVVKDLVPILVHEKYDKEIFKVAVRLLVNLTLPLEC
Ix_TIM        -----
Daph_TIM-1    EINAKLSCEEPNTRPLRRAICFFNVLNKDLIPILSSKDQPTIFRTTIKLLAELTTPTEC

Dros_TIM      LFSVDVMYRTDV--GRHTIFELNKLLYTSKEAFTEARSTKSVV-EYMKHILESDEP--KL
Lim_TIM       LVPVDIASKTSS--GKSTIIQLHHCLEGRKAPLDVRSTAABI-KSICKPFLMEKEGSSS
Ix_TIM        -----M-RVFVLLFGQDQEPSEL
Daph_TIM-1    LTCTDASSNRSTSQRTVTHELSQLLYSTKEATLEQPNATKVVIDHTEELL--EKKTTI
                                     :  :  :
Dros_TIM      SPHKCDQINNCLLLLRNLLHIPETHAHCVMFMMS-M-----
Lim_TIM       KESGSSFVNNCLLLLRNLLHIPDKIDTVKYGVDESVE-----
Ix_TIM        TENDLEGLSNCLVLLRNLLHVPDKSESAKFGISENTY-----
Daph_TIM-1    SREDCECVQHSLLLRNLLHVPQRPRTVVDVESATSSAARQPRDAQT PHQVFAQNGSS
      . . . . . * : * : * : * : * : . . . . .

Dros_TIM      ---PEGISMQNTILWNLFIQSIDKLLYLMTCPORAFWGVMTVOLIALYKDOHSGSDS
Lim_TIM       ---LEWQSIQNKLVVNLVQGLDEALLLVNNKKEKWTIAIVOLIALYKDOHISKIQ
Ix_TIM        ---DDWLSVHNKLIWNLFVHGLDGVLLMLNSEYKVNWAVVVQIIALLYKDQDVCSLQ
Daph_TIM-1    CTTADCNSEQENQRLLWNLFQRTLDRLLTNTLTSPQKGDWTVTITQIVAFYKDRHFEDMK
      . . . : * * * : * * : * : . . . : * : . . . : * : * : * : . . .

Dros_TIM      SPMLTSDPTSDSSDNGSNGRCMGGGMREGTAATLOEVSRRKQ
Lim_TIM       QLITLVSSGSSESSGEETESDTSK-----HSSATLSS-----ESY-NKKSIHSHNSDSGFT
Ix_TIM        HLISNSTSTSESEDDIESNTSK-----HSMHHTSS-----NDSVQQQSLNKLVGGT---
Daph_TIM-1    KLMEAPPTFESSDEHSDAINNTPIVNRAD-----AHVTAYSPSKNF
      :      : * * : : . :      :

```

564

565 Figure 5: TIMELESS domain region of putative *L. polyphemus* TIMELESS (TIM) protein.566 Alignment of *L. polyphemus* (Lim) TIM to *Daphnia pulex* (Daph) TIM and *D.*567 *melanogaster*(Dros) TIM using CLUSTAL Omega. Symbols immediately below sequence

568 indicates functional similarity of residues between protein orthologs. “*” indicates identical

569 residues, “:” indicates residues with strongly similar properties, “.” Indicates residues with

570 weakly similar properties. **Dark Blue** shows TIMELESS domain. For full alignment see

571 Supplemental Figure 2.

572

CRYPTOCHROME 2

[illegible]

573

Figure 6: Putative *L. polyphemus* CRYPTOCHROME 2 (CRY2) protein. Alignment of *L. polyphemus* (Lim) CRY2 to *Daphnia pulex* (Daph) CRY-M and *Apis mellifera* (Apis) CRY2 using CLUSTAL Omega. Symbols immediately below sequence indicates functional similarity of residues between protein orthologs. “*” Indicates identical residues, “:” Indicates residues with strongly similar properties, “.” Indicates residues with weakly similar properties. **Dark green** shows Photolyase domain, **Gold** shows FAD Binding 7 domain.

CLOCK (bHLH, PASa, PASb and PAC domains)

[illegible]

Figure 7: bHLH, PASa, PASb and PAC domains region of putative *L. polyphemus* CLOCK (CLK) protein. Alignment of *L. polyphemus* (Lim) CLK to *Daphnia pulex* (Daph) CLK, *D. melanogaster* (Dros) CLK, *Ixodes scapularis* (Ix) CLK using CLUSTAL Omega. Symbols immediately below sequence indicates functional similarity of residues between protein orthologs. “*” Indicates identical residues, “:” Indicates residues with strongly similar properties, “.” Indicates residues with weakly similar properties. Blue shows bHLH domain, Light green shows PAS domain, Red shows PAC domain. For full alignment see Supplemental Figure 3.

589
 590 Figure 8: *L. polyphemus* PERIOD phylograms. *L. polyphemus* homologs are marked with an “*”,
 591 chelicerates shown in red, while all other arthropods are shown in blue, and all vertebrates are
 592 labeled in black, the root is a fungi ortholog and is labeled in violet. Major clades are bracketed
 593 and labeled. Branch lengths correspond to distance bar on bottom left corner. See Supplemental
 594 figure 5 for Cladogram.

Figure 9: *L. polyphemus* TIMELESS phylograms. *L. polyphemus* homologs are marked with an “*”, chelicerates shown in red, while all other arthropods are shown in blue, and all vertebrates are labeled in black, the root is a fungi ortholog and is labeled in violet. Major clades are bracketed and labeled. Branch lengths correspond to distance bar on bottom left corner. See Supplemental figure 6 for Cladogram.

Figure 10: *L. polyphemus* CLOCK/CYCLE phylograms. *L. polyphemus* homologs are marked with an “*”, chelicerates shown in red, while all other arthropods are shown in blue, and all vertebrates are labeled in black, the root is a fungi ortholog and is labeled in violet. Major clades are bracketed and labeled. Branch lengths correspond to distance bar on bottom left corner. See Supplemental figure 7 for Cladogram.

Figure 11: *L. polyphemus* CRYPTOCHROME phylograms. *L. polyphemus* homologs are marked with an “*”, chelicerates shown in red, while all other arthropods are shown in blue, and all vertebrates are labeled in black, the root is a fungi ortholog and is labeled in violet. Major clades are bracketed and labeled. Branch lengths correspond to distance bar on bottom left corner. See Supplemental figure 8 for Cladogram.

Figure 12: *L. polyphemus* SLIMB phylograms. *L. polyphemus* homologs are marked with an “*”, chelicerates shown in red, while all other arthropods are shown in blue, and all vertebrates are labeled in black, the root is a fungi ortholog and is labeled in violet. Major clades are bracketed and labeled. Branch lengths correspond to distance bar on bottom left corner. See Supplemental figure 9 for Cladogram.