**Forensic and population genetic analysis of Xinjiang Uyghur
population on 21 short tandem repeat loci of 6-dye
GlobalFiler™ PCR Amplification kit**

Abstract

Estimating the allele frequencies and forensic statistical parameters of commonly used short tandem repeat (STR) loci of the Uyghur population, which is the fifth largest group in China, provides a more precise reference database for forensic investigation.

The 6-dye GlobalFiler™ Express PCR Amplification kit incorporates 21 autosomal STRs, which have been proven that could provide reliable DNA typing results and enhance the power of discrimination. Here we analyzed the GlobalFiler STR loci on 1,962 unrelated individuals from Chinese Uyghur population of Xinjiang, China. No significant deviations from Hardy-Weinberg equilibrium and linkage disequilibrium were detected within and between the GlobalFiler STR loci. SE33 showed the greatest power of discrimination in Uyghur population, whereas TPOX showed the lowest. The combined power of discrimination was 99.999999999999999999998746%. No significant difference was observed between Uyghur and the other two Uyghur populations at all tested STRs, as well as Dai and Mongolian. Significant differences were only observed between Uyghur and other Chinese populations at TH01, as well as Central-South Asian at D13S317, East Asian at TH01 and VWA. The phylogenetic analysis showed that Uyghur is genetically close to Chinese populations, as well as East Asian and Central-South Asian.

1. Introduction

Uyghur is one of ethnic populations of Xinjiang Autonomous Region in northwest China. According to the 2010 census, the population of Uyghur had reached 10,069,346, who becomes the fifth largest group in China. Previous genetic [1-4] and physical anthropology [5] studies have showed that Uyghur had a substantial admixture of East-Asian and European ancestries, making Uyghur has a great concern by population genetics and epidemiological investigations [1, 6].

The 6-dye GlobalFiler™ Express PCR Amplification kit was designed to incorporate 21 commonly used autosomal STR loci (D8S1179, D21S11, D7S820, CSF1PO, D3S1358, TH01, D13S317, D16S539, D2S1338, D19S433, VWA, TPOX, D18S51, D5S818, FGA, D12S391, D1S1656, D2S441, D10S1248, D22S1045 and SE33) and three gender determination loci (Amelogenin, Yindel and DYS391) [7], which have been proven that could provide reliable DNA typing results and enhance the power of discrimination [8, 9]. In this study, to estimate the allele frequencies and forensic statistical parameters of 21 GlobalFiler STR loci on Chinese Uyghur population, we typed a total of 1,962 Uyghur samples. We compared allele frequencies of each STR locus between Uyghur and other populations. We also constructed Neighbor-Joining tree to reveal the genetic relationships among these populations.

2. Materials and methods

2.1. Samples and experiments

A total of 1,962 unrelated Uyghur blood donors were sampled from Xinjiang after

acquiring their informed consent.

All DNA samples were amplified using GlobalFiler™ Express kit (Thermo Fisher Scientific Company, Carlsbad, USA) in the GeneAmp PCR System 9700 (Thermo Fisher Scientific Company) according to manufacturer's recommendation. A total of 24 loci were amplified, including 21 autosomal STR loci and three gender determination loci. The PCR products were separated by capillary electrophoresis in an ABI PRISM 3730xL Genetic Analyzer (Thermo Fisher Scientific Company), and the GeneMapper® ID-X Software v1.4 (Thermo Fisher Scientific Company) was used for genotype assignment. DNA typing and assignment of nomenclature were based on the ISFG recommendations [10-12].

2.2. Statistical analyses

The observed heterozygosity (H_o) and expected heterozygosity (H_e) were estimated using Arlequin v3.5 [13]. The exact test of Hardy-Weinberg equilibrium (HWE) and linkage disequilibrium (LD) were also performed using Arlequin v3.5 [13]. Match probability (MP), power of discrimination (PD) and power of exclusion (PE) were estimated using Modified-Powerstats [14]. The exact test of population differentiation was performed between Uyghur and other populations using Arlequin v3.5 [13]. Unbiased estimate of F_{st} was calculated as described by Weir and Hill [15], and the negative values were set to zero. We also constructed Neighbor-Joining tree using Mega 6 software [16].

3. Results and discussions

3.1 HWE and LD test

No significant deviation from HWE was observed after Bonferroni correction ($P > 0.002381$) (see Table S1), suggesting the genotype frequencies of the GlobalFiler STR loci can be estimated by their allele frequencies in Uyghur population in the present study. We also found no significant deviations from LD between pairwise STR loci after Bonferroni correction ($P > 0.0002381$) in Uyghur population (see Table S2), that means match probability can be calculated by multiplying the genotype frequencies across 21 STR markers.

3.2 Allele frequencies and forensic parameters

[illegible]

3.3 Exact test of population differentiation and genetic distance

The exact test of population differentiation was performed between Uyghur and other populations from previous reports. Overall, the allele frequency data of the 20 populations were obtained, including seven CEPH-HGDP continental groups [17, 18], the populations from South Portuguese [19] and United Arab Emirates [20], as well as

eleven Chinese populations [21, 22]. CEPH-HGDP contains a total of 51 globally distributed populations, but the sample size of each population is not enough to provide accurate estimate of the allele frequencies [17]. Therefore, the allele frequencies of seven continental groups were analyzed, including African (AFR), European (EUR), NorthAfrican/Middle Eastern (ME), Central-South Asian (S ASN), East Asian (E ASN), Native American (AME) and Oceanian (OCE). The other two populations, South Portuguese and United Arab Emirates were both typed by the GlobalFiler kit, who were also compared with the Uyghur population of the present study. The other Chinese populations from previously reports include two Uyghur, six Han and the other three ethnic populations.

As shown in Table S3, no significant difference was observed between Uyghur in the present study and the other two Uyghur populations at all tested STR loci after Bonferroni correction ($p = 0.05/358 = 0.0001397$, 358 was the number of test), as well as Dai and Mongolian. Significant differences were only observed at TH01 between Uyghur and other Chinese populations, including six Han populations (Jilin, Qinghai, Gansu, Shandong, Jiangsu and Guangdong) and Tibetan. Uyghur only showed significant differences between S ASN at D13S317, as well as E ASN at TH01 and VWA. We observed significant differences between Uyghur and other populations at multiple STR loci (see details in Table S3). The results showed that Chinese Uyghur population had the similar allele frequencies of the GlobalFiler STRs with Chinese populations, as well as East Asian and Central-South Asian.

To measure the genetic distance between Uyghur and other populations, we

estimated pairwise Fst based on all and fifteen shared STR loci, respectively. The Neighbor-Joining trees showed that Uyghur is genetically close to Chinese populations, as well as East Asian and Central-South Asian (Figure 1 and Table S4-5).

4. Conclusions

In conclusion, we report the allele frequencies and forensic statistical parameters of the GlobalFiler STR loci in Chinese Uyghur population. The results may serve as a forensic database reference of Chinese populations.

This study follows the new guidelines for publication of population data requested by the journal [23, 24].

Acknowledgments

This study was supported by grants from the National Science Foundation of China (31271338), Project of Chinese Ministry of Education (113022A) and National High Technology Research and Development Program (2012AA021802).

Conflict of interest

The authors declare that they have no conflict of interest.

Figure legends

Figure 1. The Neighbor-Joining trees constructed with pairwise Fst. AFR, AME, ASN, E ASN, EUR, ME and OCE are African, American, Central-South Asian, East Asian, European, North African/Middle Eastern and Oceanian, respectively. (A) The Neighbor-Joining tree among ten populations typed on all 21 STR loci in GlobalFiler™ kit. (B) The Neighbor-Joining tree among 21 populations including Chinese and world populations. The Fst were estimated based on fifteen shared STR

loci of these populations.

References

- [1] Xu SH, Jin L: **A genome-wide analysis of admixture in Uyghurs and a high-density admixture map for disease-gene discovery.** *Am J Hum Genet* 2008, **83**(3):322-336.
- [2] Xu SH, Huang W, Qian J, Jin L: **Analysis of genomic admixture in Uyghur and its implication in mapping strategy.** *Am J Hum Genet* 2008, **82**(4):883-894.
- [3] Lou HY, Li SL, Jin WF, Fu RQ, Lu DS, Pan XW, Zhou HG, Ping Y, Jin L, Xu SH: **Copy number variations and genetic admixtures in three Xinjiang ethnic minority groups.** *Eur J Hum Genet* 2015, **23**(4):536-542.
- [4] Xu SH, Jin WF, Jin L: **Haplotype-Sharing Analysis Showing Uyghurs Are Unlikely Genetic Donors.** *Mol Biol Evol* 2009, **26**(10):2197-2206.
- [5] Tan JZ, Peng QQ, Li JX, Guan YQ, Zhang LP, Jiao Y, Yang YJ, Wang SJ, Jin L: **Characteristics of dental morphology in the Xinjiang Uyghurs and correlation with the EDARV370A variant.** *Sci China-Life Sci* 2014, **57**(5):510-518.
- [6] Li J, Lou HY, Yang X, Lu DS, Li SL, Jin L, Pan XW, Yang WJ, Song MS, Mamatyusupu D *et al*: **Genetic architectures of ADME genes in five Eurasian admixed populations and implications for drug safety and efficacy.** *J Med Genet* 2014, **51**(9):614-622.
- [7] Wang DY, Gopinath S, Lagacé RE, Norona W, Hennessy LK, Short ML, Mulero JJ: **Developmental validation of the GlobalFiler® Express PCR Amplification Kit: A 6-dye multiplex assay for the direct amplification of reference samples.** *Forensic Sci Int-Genet* 2015, **19**:148-155.
- [8] Flores S, Sun J, King J, Budowle B: **Internal validation of the GlobalFiler (TM) Express PCR Amplification Kit for the direct amplification of reference DNA samples on a high-throughput automated workflow.** *Forensic Sci Int-Genet* 2014, **10**:33-39.
- [9] Martin P, de Simon LF, Luque G, Farfan MJ, Alonso A: **Improving DNA data exchange: Validation studies on a single 6 dye STR kit with 24 loci.** *Forensic Sci Int-Genet* 2014, **13**:68-78.
- [10] Bar W, Brinkmann B, Budowle B, Carracedo A, Gill P, Lincoln P, Mayr W, Olaisen B: **DNA recommendations - Further report of the DNA Commission of the ISFH regarding the use of short tandem repeat systems.** *International Journal of Legal Medicine* 1997, **110**(4):175-176.
- [11] Olaisen B, Bar W, Brinkmann B, Budowle B, Carracedo A, Gill P, Lincoln P, Mayr WR, Rand S: **DNA recommendations 1997 of the International Society for Forensic Genetics.** *Vox Sanguinis* 1998, **74**(1):61-63.
- [12] Gettings KB, Aponte RA, Vallone PM, Butler JM: **STR allele sequence variation: Current knowledge and future issues.** *Forensic Sci Int-Genet* 2015, **18**:118-130.
- [13] Excoffier L, Lischer HEL: **Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows.** *Molecular Ecology Resources* 2010, **10**(3):564-567.
- [14] Zhao F, Wu X, Cai G, Xu C: **The application of Mdived-Powerstates software in forensic biostatistics (In Chinese).** *Chinese Journal of Forensic Medicine* 2003, **18**(5):297-298.
- [15] Weir BS, Hill WG: **Estimating F-statistics.** *Annual Review of Genetics* 2002, **36**:721-750.
- [16] Tamura K, Stecher G, Peterson D, Filipski A, Kumar S: **MEGA6: Molecular Evolutionary**

Genetics Analysis Version 6.0. *Mol Biol Evol* 2013, 30(12):2725-2729.

- [17] Phillips C, Fernandez-Formoso L, Garcia-Magarinos M, Porras L, Tvedebrink T, Amigo J, Fondevila M, Gomez-Tato A, Alvarez-Dios J, Freire-Aradas A *et al*: **Analysis of global variability in 15 established and 5 new European Standard Set (ESS) STRs using the CEPH human genome diversity panel. *Forensic Sci Int-Genet* 2011, 5(3):155-169.**
- [18] Phillips C, Fernandez-Formoso L, Gelabert-Besada M, Garcia-Magarinos M, Amigo J, Carracedo A, Lareu MV: **Global population variability in Qiagen Investigator HDplex STRs. *Forensic Sci Int-Genet* 2014, 8(1):36-43.**
- [19] Almeida C, Ribeiro T, Oliveira AR, Porto MJ, Santos JC, Dias D, Dario P: **Population data of the GlobalFiler® Express loci in South Portuguese population. *Forensic Sci Int-Genet* 2015, 19:39-41.**
- [20] Alhmoudi OA, Jones RJ, Tay GK, Alsafar H, Hadi S: **Population genetics data for 21 autosomal STR loci for United Arab Emirates (UAE) population using next generation multiplex STR kit. *Forensic Sci Int-Genet* 2015, 19:190-191.**
- [21] Yuan L, Liu H, Liao Q, Xu X, Chen W, Hao S: **Genetics analysis of 38 STR loci in Uyгур population from Southern Xinjiang of China. *International Journal of Legal Medicine* 2015 Oct 5. [Epub ahead of print].**
- [22] Li L, Xu J, Liu X, Liu W, Xia M, et al: **Population data of 15 short tandem repeat loci in 1084 individuals from six Han and four ethnic populations in China. *Forensic Sci Int-Genet* 2015, 19:e146–e147.**
- [23] Carracedo A, Butler JM, Gusmao L, Linacre A, Parson W, Roewer L, Schneider PM: **New guidelines for the publication of genetic population data. *Forensic Sci Int-Genet* 2013, 7(2):217-220.**
- [24] Carracedo A, Butler JM, Gusmao L, Linacre A, Parson W, Roewer L, Schneider PM: **Update of the guidelines for the publication of genetic population data. *Forensic Sci Int-Genet* 2014, 10:A1-A2.**

Figure