1 Risk of encountering ticks and tick-borne pathogens in a rapidly growing metropolitan area in
2 the U.S. Great Plains

3

4 Bruce H Noden¹, Scott R. Loss², Courtney Maichak¹, Faithful Williams¹

5 ¹Department of Entomology and Plant Pathology, Oklahoma State University, Stillwater, OK
6 74074 USA.

7 Maichak email: cmaicha@ostatemail.okstate.edu

8 Williams email: faithfu@ostatemail.okstate.edu

9 ²Department of Natural Resource Ecology and Management, Oklahoma State University,
10 Stillwater, OK 74078 USA. Email: scott.loss@okstate.edu

11 Corresponding author: Dr. Bruce Noden: bruce.noden@okstate.edu

Abstract

The prevalence of tick-borne diseases has increased dramatically in many urban areas of the U.S., yet little is known about the ecology of ticks and tick-borne pathogens in relation to characteristics of North American urban and suburban landscapes. This study aimed to begin identification of the risk of encountering ticks and tick-borne pathogens within a rapidly expanding metropolitan area in the U.S. Great Plains region. Ten sites across Oklahoma City, Oklahoma were selected for tick sampling based on presence of tick habitat and level of urbanization intensity. Sampling was conducted using CO₂ traps and flagging in June, July and October 2015. A total of 552 ticks were collected from eight of the ten sampled greenspaces. The majority of ticks collected in summer were *Amblyomma americanum* (N=534 (97.8%)), followed by *Dermacentor variabilis* (N=10 (1.8%)) and *Amblyomma maculatum* (N=2 (0.3%)). *Ixodes scapularis* adult females (N=4) and nymphal *A. americanum* (N=2) were also collected in October 2015. Tick species diversity was highest in sites with >15% of the surrounding landscape composed of undeveloped land. *Rickettsia* sp. (including *R. amblyommii* and ‘*Candidatus R. andeanae*’), *Ehrlichia chaffeensis* and/or *E. ewingii* were detected in tick pools from all eight sites where ticks were found. Our data suggest that the risk of encountering ticks and tick-borne pathogens exists throughout the Oklahoma City metropolitan area and that tick populations are likely influenced by urbanization intensity. Continued research is needed to clarify the full range of abiotic and biotic features of urban landscapes that influence the risk of encountering ticks and transmitting tick-borne diseases.

Keywords: Rickettsiae, Ehrlichiae, Greenspace, Great Plains, *Amblyomma americanum*, urban ecology

Introduction

The emergence and increased prevalence of vector-borne diseases in U.S. urban areas is a major public health concern (Bonnefoy et al. 2008; LaDeau et al. 2015). In particular, the prevalence of ticks and tick-borne diseases has increased dramatically in many U.S. urban areas (Salgo et al. 1988; Maupin et al. 1991; Jobe et al. 2007; Rydzewski et al. 2012; Blanton et al. 2014). For example, reported U.S. human cases and ticks infected with Lyme disease (caused by *Borrelia burgdorferi*), spotted fever group (SFG) rickettsiosis (caused by various *Rickettsia* spp.) and ehrlichiosis (caused by *Ehrlichia chaffeensis* and *E. ewingii*) are known to occur in urban areas (Salgo et al. 1988; Maupin et al. 1991; Blanton et al. 2014). In addition, pathogens like the *Ehrlichia muris*-like agent (Pritt et al. 2009), *Borrelia miyamotoi* (Krause et al. 2013), Heartland virus (McMullan et al. 2012), and Bourbon virus (Kosoy et al. 2015) are either emerging or just now being detected (Telford and Goethert 2004), and these pathogens are likely to affect urban areas in the future. Thus, urban and suburban residents are being exposed, perhaps more than ever, to the risk of several tick-borne diseases in or near their own backyards. Nonetheless, few studies in the Great Plains region have investigated tick populations and tick-borne pathogen prevalence in relation to abiotic and biotic characteristics of urban landscapes.

In this regard, we hypothesize there are strong relationships between the ecology of tick-borne pathogens and urban areas in the U.S. because: (1) the abiotic conditions (e.g., moisture and temperature) and biotic conditions (e.g., local vegetation and landscape-scale patterns of greenspace) that drive tick distributions in rural areas vary predictably with varying urbanization intensity (McDonnell et al. 1997; Pickett et al. 2001; Kalnay and Cai 2003; McKinney et al. 2008), (2) populations and communities of other arthropods, including ants, bees, and butterflies,

vary predictably across urban landscapes (Vergnes et al. 2012; Casner et al. 2014; Fortel et al. 2014; Savage et al. 2014), (3) populations of vertebrates that are common hosts for ticks—including birds and mammals—vary predictably across urban landscapes (Blair 1996; Loss et al. 2009; Thamm et al. 2009; Hamer et al. 2012), and (4) relationships between ticks and characteristics of urban areas are beginning to be found in other parts of the world (Elfving et al. 2010; Hornok et al. 2013; reviewed by Rizzoli et al. 2014 and Uspensky 2014).

A high prevalence of tick-borne disease transmission occurs in Oklahoma, including some of the highest rates of spotted fever group (SFG) rickettsiosis (Drexler et al. 2016), ehrlichiosis (Heitmann et al. 2016), tularemia (*Francisella tularensis*) (CDC 2015), and recently, one of the three fatal cases of the rare Heartland virus (OSDH, 2014a) and the second U.S. case of Bourbon virus (KFOR News 2015). Notably, historical mapping suggests increased incidence and westward expansion in Oklahoma for both SFG rickettsiosis and ehrlichiosis with Oklahoma City geographically on the edge of this increased disease risk (Noden, unpublished data). The Oklahoma City metropolitan region therefore makes an ideal field laboratory to investigate the risk of tick-borne pathogen transmission to humans and companion canines. The city is among the top-10 fastest growing metropolitan areas in the U.S. (CNN 2014), has 7,000 acres of parkland (along with >10,000 acres in adjacent suburbs) (City of Oklahoma City 2015), and has a climate favorable for outdoor recreation and tick activity for 10 months of the year (Talley et al. 2014).

This study aimed to begin evaluation of the risk of exposure to ticks and tick-borne pathogens across a gradient of urbanization intensity in the rapidly expanding Oklahoma City metropolitan

area. Our specific objectives were to quantify tick abundance and species diversity, as well as potential risk of exposure to tick-borne pathogens, in greenspaces with differing levels of surrounding undeveloped land.

Methods and Materials

Study site selection. Ten sites throughout the Oklahoma City metropolitan area were sampled for ticks in June, July, and October of 2015. Using Google Earth, a larger pool of 87 candidate sites were first chosen that were publicly accessible and included substantial ground- and shrub-layer vegetation and trees that provided sufficient habitat for ticks (vegetation features were determined by “ground truthing” with the Google Street View feature). Using ArcGIS 10.1 (Environmental Systems Research Institute, Redlands, California) and land cover data from the National Land Cover Database (NLCD; Fry et al. 2011), we calculated (in 1,000 m buffers around each site) percentage of impervious surface (i.e., buildings and pavement), percentage of high density development, and percentage of undeveloped land (defined to include NLCD cover classes for forest (all types), shrub/scrub, grass/herbaceous, pasture/hay, and woody wetland). Ten sites were then chosen among the candidate sites such that each of two classes of urban development intensity were represented (as based on the percentage of surrounding undeveloped land, Class 1: $\geq 15\%$ undeveloped land; Class 2: $\leq 15\%$ undeveloped land). Due to access and other logistical issues, an equivalent number of sites in each class could not be sampled; instead, three selected sites were in Class 1 and seven were in Class 2. Once chosen, sites were visited to ensure that favorable tick habitat was present.

Tick sampling: All 10 sites were sampled for ticks once between June 23 and July 2, 2015, a period close to the end of the peak seasonal period of foraging for many tick species in our study area. All sampling occurred between 9am and 12pm (i.e., before temperatures became too hot for tick activity). At each site, six CO₂ traps were set for one hour and 1 or 2 persons also flagged during this period. Details of these sampling methods are described in Barrett et al. (2015). CO₂ traps were placed 10-50m apart along a transect that was defined prior to site visits using Google Earth. For flagging, one or two individuals flagged vegetation along paths and in other areas heavily used by people and their dogs. Using both types of collection techniques, we were able to detect actively questing ticks. All ticks were stored in 70% EtOH and identified in the laboratory using standard keys (Keirans & Litwak 1989; Keirans & Durden 1998).

One additional sampling event occurred at one site (Martin Park Nature Center) in October 2015 in the same locations used for trapping and flagging during the summer. The objective of this sampling event was to detect *Ixodes scapularis* adults, which are primarily active in Oklahoma during October and November. The only difference for this sampling event was that flagging was conducted by 10 individuals for one hour each.

Pathogen testing: Pools of collected ticks were tested for *Rickettsia* spp. and *Ehrlichia chaffeensis* and *E. ewingii* DNA using modified PCR protocols (Salazar 2015; Mitcham 2015). Adult ticks were grouped into pools of one to five ticks, and nymphs were grouped into pools of up to 25 ticks. After washing in de-ionized water and 70% ethanol, individual adult ticks were bisected with one half used for DNA extraction and the other half stored at -80°C. Nymphs were not bisected. Individual adults or pools of nymphs were heated at 80-90°C for fifteen minutes in 2mL vials (SARSTEDT) with 100µL of DNAzol® Direct sample processing reagent. After heating, zirconia/silica beads (BioSpec Products) were added and the tubes were placed in a

Mini-Beadbeater-16 (BioSpec Products) for three minutes. After bead-beating, resulting supernatant was collected and stored at -20° C until DNA testing. Prior to PCR testing, pools of DNA were created from up to five adults or 25 nymphs from the extracted samples of individual adults detailed above. Pooled samples of DNA were tested for rickettsial and ehrlichial DNA by PCR using protocols developed by Dawson et al. (1996) and modified by Salazar (2015). Pooled samples of *Amblyomma americanum*, *Dermacentor variabilis*, and *Amblyomma maculatum* were screened by end-point PCR for the presence of *Rickettsia* spp. using the 17kd pan-specific rickettsia primers (TZ15/TZ16) (Tzianabos et al. 1989) and confirmed using the citrate synthase (gltA) primers (CS-78/CS-323) described in Labruna et al. (2004). A nested PCR assay for *Ehrlichia* spp. specific to *E. chaffeensis* or *E. ewingii* (Dawson et al. 1996) was also used to screen all pools of ticks. Fifteen *Rickettsia* positive samples and nine positive samples of each *Ehrlichia* species were chosen for sequencing using both primer sets. The positive bands were gel extracted using a PureLink™ Quick Gel Extraction Kit (Invitrogen) and then sequenced at the Oklahoma State University Core Facility. These sequences were then searched in the nucleotide BLAST database to verify the primers amplified the targets.

Results

A total of 552 ticks consisting of 4 species were collected from eight of the ten sites (Table 1), including from sites in both development intensity classes (Figure 1). The majority of ticks collected during the summer were *A. americanum* (N=534 (97.8%)) followed by *D. variabilis* adults (N=10 (1.8%)) and *A. maculatum* adults (N=2 (0.3%)). Of the 534 *A. americanum* collected, 333 (62.4%) were nymphs. *Ixodes scapularis* adult females (N=4) and *A. americanum* nymphs (N=2) were collected in Martin Park Nature Center during the October sampling event. Tick species diversity was highest in sites with surrounding undeveloped land

greater than 15% (Table 1), and the vast majority of ticks were collected from sites with relatively high percentages of surrounding undeveloped land, including Martin Park Nature Center (N = 457 ticks; 83.7% of all ticks collected; 20% surrounding undeveloped land) and Trosper Park (N=57 ticks; 10.1% of all ticks collected; 64% surrounding undeveloped land) (Figure 1). However, smaller numbers of ticks were also found in highly urbanized sites with very little surrounding undeveloped land, including Trinity School (N=9 ticks; 0% surrounding undeveloped land) and Dolese Youth Park (N=1 tick; 2% surrounding undeveloped land).

At least one pathogen or group of pathogens (*Rickettsia* sp., *E. chaffeensis* or *E. ewingii*) was detected at each of the eight sites where ticks were collected. For *Rickettsia* spp., 54 (72%) pools of *A. americanum*, six (66.7%) pools of *D. variabilis* and one (50%) pool of *A. maculatum* tested positive (Table 2). For *E. chaffeensis*, 20 (26.7%) pools of *A. americanum* tested positive, and 14 (18.7%) pools of this species were positive for *E. ewingii*. One pool (11.1%) of *D. variabilis* tested positive for *E. chaffeensis* and one pool (11.1%) tested positive for *E. ewingii*. Of the 75 pools of *A. americanum*, fourteen (18.7%) were positive for *Rickettsia* spp. and *E. chaffeensis*, six (8.0%) were positive for *Rickettsia* spp. and *E. ewingii*, and two (2.7%) were positive for *E. chaffeensis* and *E. ewingii*. Six pools (8.0%) were positive for *Rickettsia* spp. and both *Ehrlichia* species. The only questing *A. americanum* male recovered by flagging in Dolese Youth Park (a park with 1.63% surrounding undeveloped land) was infected with *E. ewingii*. Additionally, of the four *A. americanum* ticks collected along trails in Stars and Stripes Park (a park with 2.6% surrounding undeveloped land), two were infected with *Rickettsia* spp. and one with *E. chaffeensis*.

Of the fifteen positive *Rickettsia* pools that were chosen for sequencing using gltA and 17kDa primers (samples from 5 parks included 7 pools of *A. americanum* adults, 3 pools of *A.*

170 *americanum* nymphs, 4 pools of individual *D. variabilis* adults and 1 pool consisting of an
 171 individual male *A. maculatum*), all *A. americanum* (10/10) and *D. variabilis* (4/4) samples were
 172 99-100% identical to known *Rickettsia amblyommii* sequences (Genbank sequence
 173 numbers CP012420.1 and CP003334.1). *Rickettsia*-positive *D. variabilis* samples were also 89-
 174 96% identical to *R. montanensis* (Genbank sequence no. CP003340.1). The rickettsial species in
 175 *A. maculatum* was 100% identical to known sequences of ‘*Candidatus Rickettsia andeanae*’
 176 (Genbank sequence nos. KT153033.1 and GU395295.1). Of the nine positive *E. chaffeensis* pools
 177 (samples from 4 parks of *A. americanum* adults (pools=7) and nymphs (pools=2)), all were 99-
 178 100% identical to known sequences of *E. chaffeensis* (Genbank sequence no. NR074500.1). The
 179 nine positive *E. ewingii* pools (samples from 4 parks of *A. americanum* adults (pools=7) and
 180 nymphs (pools=2)) were 99% identical to known sequences of *E. ewingii* (Genbank sequence
 181 no. NR044747.1).

182 **Discussion**

183 Our results demonstrate the presence of ticks across varying levels of urbanization intensity in the
 184 Oklahoma City metropolitan area, from greenspaces near the city core and surrounded by a
 185 highly urbanized landscape to parks in outer suburban areas surrounded by a relatively
 186 undeveloped landscape. Likewise, tick-borne pathogens were detected across the entire gradient
 187 of urbanization intensity: three tick species were collectively infected with at least three different
 188 pathogens of medical and veterinary importance. We also note a potential trend for increasing
 189 tick abundance in association with increasing amounts of surrounding undeveloped land. Across
 190 the United States, including in Oklahoma, the highest-risk areas for tick-borne pathogen exposure
 191 are generally considered to be outside of urban and suburban areas in grasslands, shrublands,
 192 woodlands, and agricultural areas. However, our results indicate that research in the U.S. should

also address the ecology of ticks and tick-borne pathogens, and the behaviors that influence tick-borne disease risk in humans and pets, in urban and suburban areas.

Although ticks are well-studied in urban areas of Europe (Hornok et al. 2014; reviewed by Rizzoli et al. 2014 and Uspensky 2014), very few U.S. studies have been designed with the *a priori* objective of systematically assessing tick populations and tick-borne pathogen risk in urban areas. Studies have investigated the presence of *B. burgdorferi*-infected ticks in Chicago (Jobe et al. 2007; Rydzewski et al. 2012) and New York City (Maupin et al. 1991; Daniels et al. 1997), and only two studies have quantified the risk of encountering ticks infected with non-Lyme pathogens (Salgo et al. 1998 in New York City; Blanton et al. 2014 in Little Rock, Arkansas). Two studies have evaluated tick populations in relation to urban characteristics. One study focused on several pathogens in ticks sampled from birds across an urbanization gradient in Chicago but did not explicitly assess how specific characteristics of the urban landscape were related to tick populations and risk of exposure to tick-borne pathogens (Hamer et al. 2012a,b). The other study sampled *I. scapularis* in a single residential area of Westchester County, NY, and related presence of infected ticks to four broad vegetation types (woods, ecotone, ornamental vegetation, and lawns) (Maupin et al. 1991). To date, no study in the U.S. Great Plains region has systematically addressed the biotic and abiotic factors driving risk of encountering ticks and transmitting tick-borne pathogens in urban areas. Interestingly, this major geographical gap seems to exist for tick ecology research in general (Loss et al. submitted).

Notably, all but one of the 15 *Rickettsia* positive pools aligned with *R. amblyommii*, with the exception being one pool of a single *A. maculatum* that aligned with ‘C.R. andeanae’. ‘C. R. andeanae’ has not been associated with infections in humans (Paddock et al. 2015). However, *Rickettsia amblyommii*, which is normally associated with *A. americanum* in Oklahoma (Heise et

al. 2010, Barrett et al. 2014), is increasingly being recognized as a probable causal agent for a
 milder form of spotted fever group rickettsiosis in humans (Apperson et al. 2008; Dahlgren et al.
 2016). Interestingly, all four pools of individual *D. variabilis* aligned at a higher percentage for
 known sequences of *R. amblyommi* than for *R. montanensis* using primers for two different genes.
 While this has been reported previously (Smith et al. 2010; Fritzen et al. 2011), *D. variabilis* is
 most typically associated with *R. rickettsii* and *R. montanensis* (Smith et al. 2010; Nadolny et al.
 2014). The majority of pools that were positive for *E. chaffeensis* and *E. ewingii* were *A.*
americanum, the main vector species for these *Ehrlichia* species in the U. S. (Paddock & Yabsley
 2007). Interestingly, both *Ehrlichia* species were each found in one pool consisting of an
 individual *D. variabilis*; while typically not associated with this tick species, these *Ehrlichia* have
 been previously reported in *D. variabilis* (Steiert & Gilfoy, 2002; Fritzen et al. 2011). The reasons
 for these somewhat unexpected pathogen/tick species associations is uncertain; however, future
 long-term tick studies will help clarify how prevalent these pathogens are in different tick species
 and whether other rickettsial species are present in the urban areas of the Great Plains.

The active host-seeking behavior displayed by the tick species collected, along with the
 identification of three known pathogens from 8 of the 10 urban parks sampled, suggests a high
 risk of tick attachment and the potential transmission of tick-borne disease for companion
 canines along edges of paths and manicured grass areas in urban greenspaces. While active
 infections with *R. amblyommi* have been reported in canines (Barrett et al. 2014; Barrett & Little,
 2016), the effect of this rickettsial species on canines is unknown. The pathogenesis of *E.*
chaffeensis and *E. ewingii* in canines, however, is well-established (Breitschwerdt et al. 1998;
 Little 2010). Given that we found tick-borne pathogens in most urban sites sampled, including
 those with small numbers of ticks collected, dogs frequenting urban parks in our study region

may acquire infected ticks and bring them into the home, thus posing an additional indirect health risk to humans. Thus, even sites perceived by the public to have a low risk of acquiring ticks could have an associated risk of tick-borne pathogen transmission to humans or pets.

Notably, all sites with greater than 15% surrounding undeveloped land had a sampling rate of at least one adult tick per trap-day, while most sites with fewer ticks sampled were in highly urbanized landscapes with less than 15% undeveloped land. Although this pattern is preliminary and requires further investigation, the finding is suggestive that there could be predictable characteristics of urban landscapes that could be assessed in a geospatial analytical framework to enhance predictions about tick encounter risk in U.S. urban areas. While intensity of development in the landscape surrounding greenspaces might be a factor influencing tick abundance and diversity (Rizzoli et al. 2014; Uspensky 2014; LaDeau et al. 2015; Löhms & Balbus 2015); we have no data to suggest this factor affects the risk for encountering tick-borne pathogens.

Indeed, ticks harboring at least one pathogen were recovered in all 8 sites where ticks were recovered. While further research into tick-host interactions is needed in urban areas, both surrounding undeveloped land and vegetation within greenspaces is likely to influence populations of white-tailed deer (*Odocoileus virginiana*), a key reservoir for many tick-borne pathogens and major transporter of ticks (Paddock & Yabsley, 2007; Walter et al. 2011). Thus, even with small tick populations, pathogen transmission cycles could persist in small fragments of undeveloped land in otherwise heavily urbanized areas.

In conclusion, we provide results of the first study of ticks and tick-borne pathogens across a major U.S. urban area in the Great Plains region. Although data are based on single sampling events at 10 sites over a two week period—factors which limit our inference due to expected variation in tick activity patterns with varying weather and environmental factors—we

nonetheless demonstrate the strong likelihood that urban residents and pets are at risk of encountering multiple ticks and tick-borne pathogens in a wide variety of urban greenspaces. Based on our results suggesting possible links between urban landscape characteristics and tick abundance and diversity, we recommend further studies into the biotic and abiotic factors influencing spatiotemporal variation in tick populations in urban areas. Given the large proportion of the global and U.S. human population in urban areas, these studies will have substantial importance for future public health, medical, and veterinary efforts to prevent and treat tick-borne pathogens.

Acknowledgements

The authors would like to thank Jennifer Siegmann, Anna Noden, and Ethan Noden for assistance with summer tick collections and the members of the Medical and Veterinary Entomology course at Oklahoma State University for assistance in tick collections in October 2015. Additionally, special thanks to Jennifer Salazar for technical assistance for processing and testing tick pools. Funding for this project was provided by NIFA/USDA Hatch Grant funds through the Oklahoma Agricultural Experiment Station (OKL-02909 and OKL-02915).

References

- Apperson, C.S., Engber, B., Nicholson, W.L., Mead, D.G., Engel, J., Yabsley, M.J., Dail, K., Johnson, J., Watson, D.W., 2008. Tick-borne diseases in North Carolina: is "*Rickettsia amblyommii*" a possible cause of rickettsiosis reported as Rocky Mountain spotted fever? *Vector Borne Zoonotic Disease* 8, 597-606.
- Barrett, A.W., Little, S.E., 2016. Vector-borne infections in tornado-displaced and owner-relinquished dogs in Oklahoma, USA. *Vector Borne Zoonotic Diseases* 16, 428-430.
- Barrett, A., Little, S.E., Shaw, E., 2014. "*Rickettsia amblyommii*" and *R. montanensis* infection in dogs following natural exposure to ticks. *Vector Borne and Zoonotic Diseases* 14, 20-25.

286 Barrett, A.W., Noden, B.H., Gruntmeir, J.M., Johnson, E.M., Mitcham, J.R., Martin, J.E., Little,
287 S.E., 2015. County scale distribution of *Amblyomma americanum* in Oklahoma: addressing local
288 deficits in tick maps based on passive reporting. *Journal of Medical Entomology* 52, 269-73.

289 Blanton, L.S., Walker, D.H., Bouyer, D. H., 2014. Rickettsiae and ehrlichiae within a city park:
290 is the urban dweller at risk? *Vector-Borne and Zoonotic Diseases* 14, 168-170.

291 Blair, R.B., 1996. Land use and avian species diversity along an urban gradient. *Ecological*
292 *Applications* 6, 506-519.

293 Bonnefoy, X., Kampen, H., Sweeney, K., 2008. Public health significance of urban pests. *World*
294 *Health*
295 *Organization*. Online: http://www.euro.who.int/__data/assets/pdf_file/0011/98426/E91435.pdf.
296 (Accessed March 8, 2016)

297 Breitschwerdt, E.B., Hegarty, B.C., Hancock, S.I., 1998. Sequential evaluation of dogs naturally
298 infected with *Ehrlichia canis*, *Ehrlichia chaffeensis*, *Ehrlichia equi*, *Ehrlichia ewingii*, or
299 *Bartonella vinsonii*. *Journal of Clinical Microbiology* 36, 2645-2651.

300 Casner, K.L., Forister, M.L., O'Brien, J.M., Thorne, J., Waetjen, D., Shapiro, A.M., 2014.
301 Contribution of urban expansion and a changing climate to decline of a butterfly fauna.
302 *Conservation Biology* 28, 773-782.

303 CDC. 2015. Tickborne Diseases of the U.S. Online: <http://www.cdc.gov/ticks/diseases/>
304 (Accessed March 8, 2016).

305 City of Oklahoma City, 2015. Oklahoma City Department of Parks, Recreation, and Cultural
306 Services. Online: <http://www.okc.gov/parks/> (Accessed March 8, 2016).

307 CNN, 2014. 10 fastest growing cities. Online:
308 http://money.cnn.com/gallery/real_estate/2014/03/27/fastest-growing-cities/9.html (Accessed
309 March 8, 2016).

310 Dahlgren, F.S., Heitman, K.N., Drexler, N.A., Massung, R.F., Behravesh, C.B., 2015. Human
311 granulocytic anaplasmosis in the United States from 2008 to 2012: a summary of national
312 surveillance data. *American Journal of Tropical Medicine and Hygiene* 93, 66-72.

313 Dahlgren, F.S., Paddock, C.D., Springer, Y.P., Eisen, R.J., Behravesh, C.B. 2016. Expanding
314 range of *Amblyomma americanum* and simultaneous changes in the epidemiology of Spotted
315 Fever group Rickettsiosis in the United States. *American Journal of Tropical Medicine and*
316 *Hygiene* 94, 35-42.

317 Daniels, T.J., Falco, R.C., Schwartz, I., Varde, S., Robbins, R.G., 1997. Deer ticks (*Ixodes*
318 *scapularis*) and the agents of Lyme disease and human granulocytic ehrlichiosis in a New York
319 City park. *Emerging Infectious Diseases* 3, 353-355.

320 Dawson, J.E., Biggie, K.L., Warner, C.K., Cookson, K., Jenkins, S., Levine, J.F., Olson, J.G.
 321 1996. Polymerase chain reaction evidence of *Ehrlichia chaffeensis*, an etiologic agent of human
 322 ehrlichiosis, in dogs from southeast Virginia. American Journal of Veterinary Research 57,
 323 1175-1179.

324 Drexler, N.A., Dahlgren, F.S., Heitman, K.N., Massung, R.F., Paddock, C.D., Behravesh, C.B.,
 325 2016. National Surveillance of Spotted Fever Group Rickettsioses in the United States, 2008-
 326 2012. American Journal of Tropical Medicine and Hygiene 94, 26-34.

327 Elfving, K., Olsen, B., Bergstorm, S., Waldenstrom, J., Lundkvist, A., Sjostedt, A., Mejlom, H.,
 328 Nilsson, K., 2010. Dissemination of Spotted Fever Rickettsia agents in Europe by migrating
 329 birds. PLoS ONE 5, e8572.

330 Fortel, L., Henry, M., Guilbaud, L., Guirao, A.L., Kuhlmann, M., Mouret, H., Rollin, O.,
 331 Vaissière, B.E., 2014. Decreasing abundance, increasing diversity and changing structure of the
 332 wild bee community (Hymenoptera: Anthophila) along an urbanization gradient. PLoS ONE 9,
 333 e104679.

334 Fritzen, C.M., Huang, J., Westby, K., Freye, J.D., Dunlap, B., Yabsley, M.J., Schardein, M.,
 335 Dunn, J.R., Jones, T.F., Moncayo, A.C., 2011. Infection prevalences of common tick-borne
 336 pathogens in adult lone star ticks (*Amblyomma americanum*) and American dog ticks
 337 (*Dermacentor variabilis*) in Kentucky. American Journal of Tropical Medicine and Hygiene 85,
 338 718-723.

339 Fry, J., Xian, G., Jin, S., Dewitz, J., Homer, C., Yang, L., Barnes, C., Herold, N., Wickham, J.,
 340 2011. Completion of the 2006 National Land Cover Database for the Conterminous United
 341 States. PE&RS 77, 858-864.

342 Hamer, S.A., Goldberg, T.L., Kitron, U.D., Brawn, J.D., Anderson, T.K., Loss, S.R., Walker,
 343 E.D., Hamer, G.L., 2012a. Wild birds and urban ecology of ticks and tickborne pathogens,
 344 Chicago, Illinois, USA, 2005-2010. Emerging Infectious Diseases 18, 1589-1595.

345 Hamer, S.A., Lehrer, E., Magle, S.B., 2012b. Wild birds as sentinels for multiple zoonotic
 346 pathogens along an urban to rural gradient in greater Chicago, Illinois. Zoonoses and Public
 347 Health 59, 355-364.

348 Harris, R.M., Couturier, B.A., Sample, S.C., Coulter, K.S., Casey, K.K., Schlager, R., 2016.
 349 Expanded geographic distribution and clinical characteristics of *Ehrlichia ewingii* infections,
 350 United States. Emerging Infectious Diseases 22, 862-865.

351 Heise, S.R., Elshahed, M.S., Little, S.E., 2010. Bacterial diversity in *Amblyomma americanum*
 352 (Acari: Ixodidae) with a focus on members of the genus Rickettsia. Journal of Medical
 353 Entomology 47, 258-268.

354 Heitman, K.N., Dahlgren, F.S., Drexler, N.A., Massung, R.F., Behravesh, C.B., 2016. Increasing
355 incidence of Ehrlichiosis in the United States: A summary of national surveillance of *Ehrlichia*
356 *chaffeensis* and *Ehrlichia ewingii* infections in the United States, 2008-2012. American Journal
357 of Tropical Medicine and Hygiene 94, 52-60.

358 Hornok, S., Csorgo, T., de la Fuente, J., Gyuranecz, M., Privigyei, C., Meli, M.R., Kreizinger,
359 Z., Gonczi, E., Fernandez de Mera, I.G., Hofmann-Lehmann, R., 2013. Synanthropic birds
360 associated with high prevalence of tick-borne Rickettsiae and with the first detection of
361 *Rickettsia aeschlimannii* in Hungary. Vector-Borne and Zoonotic Diseases 13, 77-83.

362 Jobe, D.A., Nelson, J.A., Adam, M.D., Martin Jr, S.A., 2007. Lyme disease in urban areas,
363 Chicago. Emerging Infectious Diseases 13, 1799-1800.

364 Kalnay, E., Cai, M., 2003. Impact of urbanization and land-use change on climate. Nature 423,
365 528-531.

366 KFOR News, 2015. Rare illness caused by a tick diagnosed found in Oklahoma, only second case
367 in U.S. Online: [http://kfor.com/2015/05/27/rare-illness-caused-by-a-tick-diagnosed-found-in-](http://kfor.com/2015/05/27/rare-illness-caused-by-a-tick-diagnosed-found-in-oklahoma-only-second-case-in-u-s/)
368 [oklahoma-only-second-case-in-u-s/](http://kfor.com/2015/05/27/rare-illness-caused-by-a-tick-diagnosed-found-in-oklahoma-only-second-case-in-u-s/). (Accessed March 6, 2016).

369 Keirans, J.E., Litwak, T.R., 1989. Pictorial key to the adults of hard ticks, Family Ixodidae
370 (Ixodida: Ixodoidea), east of the Mississippi River. Journal of Medical Entomology 26, 435-448.

371 Keirans, J.E., Durden, L.A., 1998. Illustrated key to nymphs of the tick Genus *Amblyomma*
372 (Acari: Ixodidae) found in the United States. Journal of Medical Entomology 35, 489-495.

373 Kosoy, O.I., Lambert, A.J., Hawkinson, D.J., Pastula, D.M., Goldsmith, C.S., Hunt, D.C.,
374 Staples, J.E., 2015. Novel thogotovirus associated with febrile illness and death, United States,
375 2014. Emerging Infectious Diseases 21, 760-764.

376 Krause, P.J., Narasimhan, S., Wormser, G.P., Rollend, L., Fikrig, E., Lepore, T., Barbour, A.,
377 Fish, D., 2013. Human *Borrelia miyamotoi* infection in the United States. New England Journal
378 of Medicine 368, 291-293.

379 Labruna, M.B., Whitworth, T., Horta, M.C., Bouyer, D.H., McBride, J.W., Pinter, A., Popov, V.,
380 Gennari, S.M., Walker, D.H., 2004. Rickettsia species infecting *Amblyomma cooperi* ticks from
381 an area in the state of São Paulo, Brazil, where Brazilian spotted fever is endemic. Journal of
382 Clinical Microbiology 42, 90-98.

383 LaDeau, S.L., Allan, B.F., Leisnham, P.T., Levy, M.Z., 2015. The ecological foundations of
384 transmission potential and vector-borne disease in urban landscapes. Functional Ecology 29,
385 889-901.

386 Little, S.E., 2010. Ehrlichiosis and anaplasmosis in dogs and cats. Veterinary Clinics of North
387 America Small Animal Practice 40, 1121-1140.

388 Löhmus, M., Balbus, J., 2015. Making green infrastructure healthier infrastructure. *Infection*
 389 *Ecology & Epidemiology* 27, 30082.

390 Loss, S.R., Hamer, G.L., Goldberg, T.L., Ruiz, M.O., Kitron, U.D., Walker, E.D., Brawn, J.D.,
 391 2009. Nestling passerines are not important hosts for amplification of West Nile virus in
 392 Chicago, Illinois. *Vector Borne and Zoonotic Diseases* 9, 13-18.

393 Maupin, G.O., Fish, D., Zultowsky, J., Campos, E.G., Piesman, J., 1991. Landscape ecology of
 394 Lyme disease in a residential area of Westchester County, New York. *American Journal of*
 395 *Epidemiology* 133, 1105-1113.

396 McDonnell, M.J., Pickett, S.T., Groffman, P., Bohlen, P., Pouyat, R.V., Zipperer, W.C.,
 397 Parmelee, R.W., Carreiro, M.M., Medley, K., 1997. Ecosystem processes along an urban-to-rural
 398 gradient. *Urban Ecosystems* 1, 21-36.

399 McKinney, M.L., 2008. Effects of urbanization on species richness: a review of plants and
 400 animals. *Urban Ecosystems*, 11, 161-176.

401 McMullan, L.K., Folk, S.M., Kelly, A.J., MacNeil, A., Goldsmith, C.S., Metcalfe, M.G., Batten,
 402 B.C., Albarrino, C.G., Zaki, S.R., Rollin, P.E., Nicholson, W.L., Nichol, S.T., 2012. A new
 403 phlebovirus associated with severe febrile illness in Missouri. *New England Journal of Medicine*
 404 367, 834-841.

405 Mitcham, Jessica. 2016. Updated ecology and distribution of tick species in Oklahoma. Master's
 406 thesis. Oklahoma State University.

407 Nadolny, R.M., Wright, C.L., Sonenshine, D.E., Hynes, W.L., Gaff, H.D., 2014. Ticks and
 408 spotted fever group rickettsiae of southeastern Virginia. *Ticks Tick Borne Disease* 5, 53-57.

409 Oklahoma State Department of Health (OSDH). 2014a. Oklahoma State Health Department
 410 Confirms first case and death of Heartland Virus. Online:
 411 [http://www.ok.gov/health/Organization/Office_of_Communications/News_Releases/2014_News](http://www.ok.gov/health/Organization/Office_of_Communications/News_Releases/2014_News_Releases/Oklahoma_State_Health_Department_Confirms_First_Case_and_Death_of_Heartland_Virus.html)
 412 [_Releases/Oklahoma_State_Health_Department_Confirms_First_Case_and_Death_of_Heartland](http://www.ok.gov/health/Organization/Office_of_Communications/News_Releases/2014_News_Releases/Oklahoma_State_Health_Department_Confirms_First_Case_and_Death_of_Heartland_Virus.html)
 413 [_Virus.html](http://www.ok.gov/health/Organization/Office_of_Communications/News_Releases/2014_News_Releases/Oklahoma_State_Health_Department_Confirms_First_Case_and_Death_of_Heartland_Virus.html) (Accessed March 6, 2016).

414 Oklahoma State Department of Health (OSDH). 2014b. Tickborne and Mosquitoborne Disease
 415 Prevention. Online:
 416 [http://www.ok.gov/health/Disease,_Prevention,_Preparedness/Acute_Disease_Service/Disease_I](http://www.ok.gov/health/Disease,_Prevention,_Preparedness/Acute_Disease_Service/Disease_Information/Tickborne_and_Mosquitoborne_Diseases/index.html)
 417 [nformation/Tickborne_and_Mosquitoborne_Diseases/index.html](http://www.ok.gov/health/Disease,_Prevention,_Preparedness/Acute_Disease_Service/Disease_Information/Tickborne_and_Mosquitoborne_Diseases/index.html) (Accessed March 8, 2016).

418 Paddock, C.D., Denison, A.M., Dryden, M.W., Noden, B.H., Lash, R.R., Abdelghani, S.S.,
 419 Evans, A.E., Kelly, A.R., Hecht, J.A., Karpathy, S.E., Ganta, R.R., Little, S.E., 2015. High
 420 prevalence of "*Candidatus Rickettsia andeanae*" and apparent exclusion of *Rickettsia parkeri* in

421 adult *Amblyomma maculatum* (Acari: Ixodidae) from Kansas and Oklahoma. Ticks and Tick
422 Borne Disease 6, 297-302.

423 Paddock, C.D., Yabsley, M.J., 2007. Ecological havoc, the rise of white-tailed deer, and the
424 emergence of *Amblyomma americanum*-associated zoonoses in the United States. Current Topics
425 in Microbiology and Immunology 315, 289-324.

426 Pickett, S.T., Cadenasso, M.L., Grove, J.M., Nilon, C.H., Pouyat, R.V., Zipperer, W.C.,
427 Costanza, R., 2001. Urban ecological systems: linking terrestrial ecological, physical, and
428 socioeconomic components of metropolitan areas. Annual Review of Ecology and
429 Systematics 32, 99-122.

430 Pritt, B.S., Sloan, L.M., Johnson, D.K., Munderloh, U.G., Paskewitz, S.M., McElroy, K.M.,
431 McFadden, J.D., Binnicker, M.J., Neitzel, D.F., Liu, G., Nicholson, W.L., Nelson, C.M.,
432 Franson, J.J., Martin, S.A., Cunningham, S.A., Steward, C.R., Bogumill, K., Bjorgaard, M.E.,
433 Davis, J.P., McQuiston, J.H., 2011. Emergence of a new pathogenic Ehrlichia species, Wisconsin
434 and Minnesota, 2009. New England Journal of Medicine 365, 422-429.

435 Rizzoli, A., Silaghi, C., Obiegala, A., Rudolf, I., Hubálek, Z., Földvári, G., Plantard, O.,
436 Vayssier-Taussat, M., Bonnet, S., Spitalská, E., Kazimírová, M., 2014. *Ixodes ricinus* and its
437 transmitted pathogens in urban and peri-urban areas in Europe: new hazards and relevance for
438 public health. Frontiers in Public Health 2, 251.

439 Rydzewski, J., Mateus-Pinilla, N., Warner, R.E., Nelson, J.A., Velat, T.C., 2012. *Ixodes*
440 *scapularis* (Acari: Ixodidae) distribution surveys in the Chicago metropolitan region. Journal of
441 Medical Entomology 49, 955-959.

442 Salazar, J.L. 2015. Detection of tick-borne pathogens in lab reared tick colonies and wild
443 populations. Master's thesis. Oklahoma State University.

444 Salgo, M.P., Telzak, E.E., Currie, B., Perlman, D.C., Litman, N., Levi, M., Nathenson, G.,
445 Benach, J.L., Al-Hafidh, R., Casey, J., 1988. A focus of Rocky Mountain spotted fever within
446 New York City. New England Journal of Medicine 318, 1345-1348.

447 Savage, A.M., Hackett, B., Guénard, B., Youngsteadt, E.K., Dunn, R.R., 2014. Fine-scale
448 heterogeneity across Manhattan's urban habitat mosaic is associated with variation in ant
449 composition and richness. Insect Conservation and Diversity 8, 216-228.

450 Smith, M.P., Ponnusamy, L., Jiang, J., Ayyash, L.A., Richards, A.L., Apperson, C.S., 2010.
451 Bacterial pathogens in ixodid ticks from a Piedmont County in North Carolina: prevalence of
452 rickettsial organisms. Vector Borne Zoonotic Diseases 10, 939-952.

- Steiert, J.G., Gilfoy, F., 2002. Infection rates of *Amblyomma americanum* and *Dermacentor variabilis* by *Ehrlichia chaffeensis* and *Ehrlichia ewingii* in southwest Missouri. *Vector Borne Zoonotic Diseases* 2, 53-60.
- Talley, J.L., Jaworski, D.C., Noden, B.H., Kocan, K.M., & Little, L. 2014. Common ticks of Oklahoma and tick-borne diseases. Oklahoma Cooperative Extension Fact Sheet (Epp-7001). Online: <http://www.okrangelandswest.okstate.edu/files/wildlife%20pdfs/EPP-7001.pdf>. (Accessed March 8, 2016).
- Telford, S.R. 3rd, Goethert, H.K., 2004. Emerging tick-borne infections: rediscovered and better characterized, or truly 'new'? *Parasitology* 129 Suppl, S301-327.
- Thamm, S., Kalko, E.K., Wells, K., 2009. Ectoparasite infestations of hedgehogs (*Erinaceus europaeus*) are associated with small-scale landscape structures in an urban-suburban environment. *Ecohealth* 6, 404-413.
- Tzianabos, T., Anderson, B.E., McDade, J.E., 1989. Detection of *Rickettsia rickettsii* DNA in clinical specimens by using polymerase chain reaction technology. *Journal of Clinical Microbiology* 27, 2866-2868.
- Uspensky, I., 2014. Tick pests and vectors (Acari: Ixodoidea) in European towns: Introduction, persistence and management. *Ticks and Tick Borne Disease* 5, 41-47.
- Vergnes, A., Viol, I.L., Clergeau, P., 2012. Green corridors in urban landscapes affect the arthropod communities of domestic gardens. *Biological Conservation* 145, 171-178.
- Walter, W.D., Beringer, J., Hansen, L.P., Fischer, J.W., Millspaugh, J.J., Vercauteren, K.C., 2011. Factors affecting space use overlap by white-tailed deer in an urban landscape, *International Journal of Geographical Information Science* 3, 379-392.

Figure captions

Figure 1. Location of 10 sampling sites across Oklahoma City, Oklahoma, USA, in June and July 2015; sites were divided two classes (indicated by open and closed circles) based on the amount of undeveloped land within 1,000 m and size of site labels indicate mean number of total ticks/trap-day collected from each site (trap-days = total number of traps set at each site during the sampling period).

Figure 1

Table 1

Table 1. Total ticks collected by species and site and sampling technique, including for sites with $\geq 15\%$ and $\leq 15\%$ undeveloped area in the surrounding landscape (i.e., within 1,000 m of sites), across Oklahoma City, Oklahoma, USA, in June and July 2015*.

Sites sampled	# Traps	# Flags	Total Ticks: CO ₂ traps^	Total Ticks / Trap	Total Ticks: Flags^	Total Ticks / Flag	% Undeveloped
Class 1 ($\geq 15\%$ undeveloped area)							
Trosper Park Archery	6	2	14	2.3	41	20.5	63.80
<i>A. americanum</i>			12 (75%)		40 (98%)		
<i>D. variabilis</i>			2 (25%)		1 (2%)		
<i>A. maculatum</i>			-		-		
Martin Park Nature Center	6	2	394	65.7	66	33	19.92
<i>A. americanum</i>			388 (98.5%)		63 (95%)		
<i>D. variabilis</i>			2 (0.5%)		2 (3%)		
<i>A. maculatum</i>			-		1 (2%)		
<i>I. scapularis</i> (Oct)			4 (1%)		-		
Bluff Creek Park	6	2	3	0.5	9	4.5	17.30
<i>A. americanum</i>			3 (100%)		6 (67%)		
<i>D. variabilis</i>			-		3 (33%)		
Class 2 ($\leq 15\%$ undeveloped area)							
Edwards Park	6	2	1	0.2	4	2.0	10.78
<i>A. americanum</i>			1 (100%)		1 (25%)		
<i>D. variabilis</i>			-		2 (50%)		
<i>A. maculatum</i>			-		1 (25%)		
Lakeshore Park	6	0	0	0.0	0	-	6.72
Lake Hefner Park	6	0	3	0.5	0	0.0	6.0
<i>A. americanum</i>			3 (100%)		-		
Stars and Stripes Park	6	1	2	0.3	2	2.0	2.40
<i>A. americanum</i>			2 (100%)		2 (100%)		
Dolese Youth Park	6	2	0	0.0	1	0.5	1.63
<i>A. americanum</i>			-		1 (100%)		

Trinity School	6	1	9	1.5	0	0.0	0.00
<i>Am. americanum</i>			9 (100%)		-		
Smitty park	4	0	0	0.0	0	-	0.00

* Martin Park Nature Center and Bluff Creek were also sampled once in October 2015.

^ Total ticks = All ticks collected in one park. Under each total is the breakdown of tick species collected with appropriate percentages based solely on each park collection.

Table 2. Tick-borne pathogens (documented using PCR analysis) in ticks sampled from 10 sites across Oklahoma City, Oklahoma, OK, in June and July 2015.

Tick species	Stage	Total no. ticks tested^	Rickettsia spp.* No. positive pools/No. pools	E. chaffeensis No. positive pools/No. pools	E. ewingii No. positive pools/No. pools
<i>A. americanum</i>	Adult	199	36/48	15/48	7/48
	Nymph	332	18/27	5/27	7/27
<i>D. variabilis</i>	Adult	10	6/9	1/9	1/9
<i>A. maculatum</i>	Adult	2	1/2	0/2	0/2

^ 6 ticks collected in Fall 2015 were not tested for pathogens.

* *R. amblyommii* was identified in 10 pools of *A. americanum* (7 adult and 3 nymph pools) and 4 pools of *D. variabilis* while ‘*Candidatus R. andeanae*’ was identified in one *A. maculatum* pool consisting of a single male.