Running Title:

Theta Burst Stimulation Effects on Suprahyoid Motor Cortex

Article Title:

Effects of Theta Burst Stimulation on Suprahyoid Motor Cortex Excitability in
Healthy Subjects

Authors:

Tuo LIN^{a, #}, Guangqing XU^{b, #}, Lisheng JIANG^a, Feng LIU^c, Yue LAN^{a, *}

^aDepartment of Rehabilitation Medicine, Guangzhou First People's Hospital,
Guangzhou Medical University, Guangzhou, China,

^bDepartment of Rehabilitation Medicine, The First Affiliated Hospital, Sun Yat-sen
University, Guangzhou, China,

^cDepartment of Geriatrics, Guangzhou First People's Hospital, Guangzhou Medical
University, Guangzhou, China,

[#]These authors contributed equally to this work.

*Corresponding Author

Email:bluemooning@163.com

Tel: +86 18988991916

This work was completed at the Guangzhou First People's Hospital.

Funding

This research was supported by grants 81371441, 81372109, 81572230 from the
National Science Foundation of China, grants 2013B051000036 and
2014B020212001 from Guangdong Provincial Science and Technology Program,

grants 2016201604030036 from Guangzhou Municipal Science and Technology Program, and the National Key Clinical Specialist Construction Programs of China.

Abstract

Background: Continuous theta burst stimulation (cTBS) and intermittent TBS (iTBS) are powerful patterns of repetitive transcranial magnetic stimulation (rTMS), with substantial potential for motor function rehabilitation post-stroke. However, TBS of suprahypoid motor cortex excitability has not been investigated. This study investigated TBS effects on suprahypoid motor cortex excitability and its potential mechanisms in healthy subjects.

Methods: Thirty-five healthy subjects (23 females; mean age = 21.66 ± 1.66 years) completed three TBS protocols on separate days, separated by at least one week. A stereotaxic neuronavigation system facilitated accurate TMS positioning. Left and right suprahypoid motor evoked potentials (SMEP) were recorded using single-pulse TMS from the contralateral suprahypoid motor cortex before stimulation (baseline) and 0, 15, and 30 min after stimulation. The SMEP latency and amplitude were analyzed via repeated measures analysis of variance.

Results: cTBS suppressed ipsilateral suprahypoid motor cortex excitability and activated the contralateral suprahypoid motor cortex. iTBS facilitated ipsilateral suprahypoid motor cortex excitability; however, it did not affect the contralateral excitability. iTBS eliminated the inhibitory effect caused by cTBS applied to the contralateral suprahypoid motor cortex. TBS had no significant effect on the latencies

of bilateral SMEP. TBS effects on suprahyoid motor cortex excitability lasted a minimum of 30 min.

Conclusions: TBS effectively regulates suprahyoid motor cortex excitability. Suppression of excitability in one hemisphere leads to further activation of the corresponding contralateral motor cortex. iTBS reverses the inhibitory effect induced by cTBS of the contralateral suprahyoid motor cortex.

Keywords: theta burst stimulation; suprahyoid motor cortex; suprahyoid motor evoked potential; swallowing

Abbreviations

AMT, active motor threshold; cTBS, continuous theta burst stimulation; DTI, diffusion tensor imaging; EMG, electromyography; FDI, first dorsal interosseous; fMRI, functional magnetic resonance imaging; LTD, long-term depression; LTP, long-term potentiation; MEPs, motor evoked potentials; MSO, maximum stimulator output; iTBS, intermittent theta burst stimulation; rTMS, repetitive transcranial magnetic stimulation; SMEP, suprahyoid motor evoked potential; UES, upper esophageal sphincter.

1. Introduction

Dysphagia is a severe disability after stroke, with an incidence that ranges between 30-70% [1]. Dysphagia has been associated with an increased risk of dehydration, malnutrition, aspiration pneumonia, and mortality [1-3]. Aspiration

pneumonia is one of the major causes of death in dysphagic stroke patients [2]. Hyoid-throat complex movement forward and upward comprises the precondition for the opening of the upper esophageal sphincter (UES), which guarantees the safety and effectiveness of swallowing. Insufficient forward and upward movement of the hyoid-throat complex contributes to repeatedly occurring aspiration and aspiration pneumonia, whereas the suprahyoid muscles play an important role in the forward and upward movement of the hyoid-throat complex [4, 5].

Traditional treatments for dysphagia have focused on behavioral exercises and compensatory strategies, such as bolus modification, swallow posture changes, and protective maneuvers [6]. Although these treatments can improve swallowing safety to some extent, they cannot promote recovery of the damaged swallow-related neural networks in dysphagic stroke patients [7]. Furthermore, a recent systematic review identified limited evidence to support the medical efficacy of common dysphagia therapies and interventions [8]. Thus, the development of novel effective rehabilitation treatments is urgently needed.

Repetitive transcranial magnetic stimulation (rTMS) is a non-invasive stimulation technique with the capacity to modulate cortical excitability and has exhibited substantial potential in dysphagia treatment. The application of high-frequency rTMS to the affected hemisphere [9] or low-frequency rTMS to the intact hemisphere [10, 11] induced an improvement in the swallowing function of unilateral stroke dysphagic patients. The potential mechanism may be related to re-balanced interhemispheric interactions because it has been demonstrated that an

interhemispheric imbalance is evident in unilateral stroke dysphagic patients and it hinders motor function recovery in stroke patients [12-14]. However, the understanding of rTMS mechanisms is inadequate, as the swallowing function also improves via the application of high-frequency rTMS to the intact hemisphere [15]. To date, no standard rTMS treatment protocols have been developed because of insufficient understanding of rTMS mechanisms. Furthermore, patient tolerance to these traditional rTMS techniques is low because of the long stimulation time and strong stimulation intensity.

Theta-burst stimulation (TBS), including continuous TBS (cTBS) and intermittent TBS (iTBS), is a powerful pattern of rTMS, which rapidly regulates motor cortical excitability [16]. Compared with traditional rTMS, TBS produces a robust and long-lasting effect with a low intensity after only 20-190 s of application in humans. Thus, it represents a safe and good rTMS option for motor function rehabilitation in stroke patients. Moreover, it has been confirmed that TBS is beneficial to the recovery of paretic hand function and aphasia in chronic stroke patients [17, 18].

However, few studies have examined the effects of TBS on the swallowing motor system. Mistry et al. [19, 20] evaluated the effects of TBS on cortical properties in the human pharyngeal motor system, and the results indicated that only iTBS to the hemisphere with stronger pharyngeal projections induced increased responses in the contralateral weaker projection 60-90 min post-iTBS [19]. No significant cTBS effect was detected on the pharyngeal motor cortex [20]. Moreover, Mistry et al. reported

limited TBS effects on the pharyngeal motor system. However, the effects of TBS on the suprahyoid motor cortex have not previously been investigated. Furthermore, no previous studies have used a neuronavigation system to facilitate an accurate TMS position. It has been established that the suprahyoid muscles were innervated by the bilateral motor cortex, which was thought to be in a state of interhemispheric balance. We hypothesized that when the motor cortex excitability of one hemisphere decreased, the motor cortex excitability of the other hemisphere would increase, and vice versa. Therefore, the present study aimed to identify the effects of neuronavigated TBS on suprahyoid motor cortex excitability and investigate its potential mechanisms in healthy subjects.

2. Materials and methods

2.1 Participants

Thirty-five healthy volunteers (23 females; age range = 20-28 years; mean age = 21.66 ± 1.66 years) participated in this study. All subjects were right handed according to the Edinburgh Handedness Inventory [21]. The exclusion criteria included a history of swallowing problems, epilepsy, or brain, ear, nose, or throat surgery; significant medical disorders; pregnancy; a cardiac pacemaker; metal in the head or eyes; or the use of medication that acts on the neuromuscular system. The experiments were approved by the clinical research ethical committee of the Third Affiliated Hospital of Sun Yat-sen University. Written informed consent was obtained from all participants prior to participation in the experiments.

2.2 Neuronavigation

The motor cortices of the suprahyoid muscles in each hemisphere were targeted with a TMS neuronavigation system (SofTaxis, E.M.S., Bologna, Italy) that used a graphic user interface and a three dimensional (3D) optical digitizer (NDI, Polaris Vicra, Ontario, Canada) to precisely position the coil over the cortical site. This approach enabled a high degree of reproducibility and accuracy throughout the experimental protocols. The procedure involved the computation of an estimated volume of head magnetic resonance images (MRIs) for all subjects because MRIs were unavailable. The estimated MRIs, which referred to the Talairach space, were calculated using a warping 3D procedure, which operated on a template MRI volume on the basis of a set of approximately 40-50 points digitized from the subject's scalp. The digitized points were used to compute a subsequent set of reference points that were analogous to a set of points pre-localized on the scalp of the template. The warping procedure was performed using these two corresponding sets of reference points [22, 23].

2.3 Electromyographic (EMG) recordings

Suprahyoid muscle EMG responses were detected with two pairs of bipolar silver-silver chloride electrodes (Yiruide, Wuhan, China). Each pair was positioned on the suprahyoid muscle surface, 1 cm lateral to the midline. One pair was placed over the left suprahyoid muscles, and the other pair was placed over the right suprahyoid muscles. The interelectrode distance was 2 cm for each pair of electrodes, measured from the center of the electrodes [24]. A pair of electrodes was placed on

the surface of the right (dominant hand in all subjects) first dorsal interosseous (FDI) in a belly-tendon montage. All electrodes were connected to an EMG recording system (Yiruide, Wuhan, China). The MEP signals were amplified and filtered with a bandpass set between 2Hz-10 kHz. The response signals were processed through a 50/60 Hz noise eliminator (Yiruide, Wuhan, China) to remove unwanted electrical interference. The MEP recordings were digitized with a sample rate of 100 kHz and stored for offline analysis, using data analysis software (Yiruide, Wuhan, China).

2.4 TMS

Magnetic stimulation was performed using a figure-of-eight coil (external loop diameters, 70 mm) connected to a Rui Chi magnetic stimulator (Yiruide CCY-IA, Wuhan, China), which produced a maximum stimulator output (MSO) of 3.0 Tesla. The stimulating coil was held tangentially to the skull with the coil handle pointing backwards and laterally 45° away from the anterior-posterior axis. The exact position was adjusted according to the results of the online neuronavigation, such that the coil wing center was oriented perpendicularly to the target point, to deliver the maximum magnetic power.

2.5 TBS

TBS was performed using a Rui Chi magnetic stimulator (Yiruide CCY-IA, Wuhan, China) connected to a figure of eight coil with a 70 mm outer diameter (MSO of 3.0 tesla). The magnetic stimulus had a biphasic waveform. When performing TBS, the coil was held in an identical orientation to single pulse TMS. TBS was applied over the “hot spot” of the suprahypoid motor cortex. TBS was applied over the suprahypoid motor cortex. The TBS pattern consisted of bursts that contained three

pulses at 50 Hz repeated at 5 Hz. In the iTBS paradigm, a 2 s train of TBS was repeated every 10 s for approximately 190 s (600 pulses). In the cTBS paradigm, a 40 s train of uninterrupted TBS was administered (600 pulses) [16]. All TBS interventions were delivered at 80% of the active motor threshold (AMT) of the FDI.

2.6 Experimental procedure

2.6.1 Determination of FDI hot spot and AMT

In every protocol, the subjects were relaxed and seated in a comfortable chair. The cortex was visualized through the subject's estimated MRI created in advance via the neuronavigation system. To locate the optimal stimulus site for the left or right FDI, the contralateral hand motor cortex around the anatomical cortex area was stimulated with 70% of the maximum output. During the stimulation, the coil was held tangential to the scalp, and the direction of the induced current (computed and visualized using the navigation software) was held perpendicular to the anatomically defined central sulcus. The motor 'hot spot' was determined as the site at which TMS consistently elicited the largest MEPs from the contralateral FDI. This 'hot spot' position was marked on the subject's estimated MRI, and the 3D coordinates were acquired and stored to ensure constant coil placement throughout the experiment. The AMT was determined at the hot spot, which was defined as the lowest stimulus intensity at which 5 of 10 consecutive stimuli elicited reliable MEPs ($\geq 200 \mu\text{V}$ in amplitude) during a weak voluntary isometric contraction of the contralateral FDI [25]. To ensure a low-level and consistent contraction of the FDI during the assessment of AMT, the subject was asked to pinch a one-dollar coin with the thumb and forefinger to ensure that it would not fall off.

2.6.2 Determination of the suprahyoid hot spot

Similar to the method of determination of the FDI hot spot, the suprahyoid motor cortex around the anatomical cortex area was stimulated with the stimulator set to 70% of the maximum output, and the suprahyoid ‘hot spot’ was defined as the site at which TMS consistently elicited the largest MEPs from the contralateral suprahyoid muscles. If the MEPs were very large or small during the location of the hotspot, the TMS intensity was adjusted (increased or decreased) to produce discernible MEPs [24]. Bilateral ‘hot spot’ positions were marked on the subject’s estimated MRI, and the 3D coordinates were stored via the neuronavigation system to ensure constant coil placement throughout the experiment.

2.7 Experimental protocols

To elucidate the neurophysiological effects of TBS on the suprahyoid motor cortex excitability, all subjects completed three successive research protocols on separate days at least one week apart.

In protocol A, the subjects were stimulated by cTBS to the left suprahyoid motor cortex hot spot with the intensity of 80% AMT of the right FDI.

In protocol B, the subjects were stimulated via iTBS to the left suprahyoid motor cortex hot spot with an intensity of 80% AMT of the right FDI.

In protocol C, the subjects were stimulated via cTBS to the left suprahyoid motor cortex hot spot with an intensity of 80% AMT of the right FDI, followed by iTBS to the right suprahyoid motor cortex hot spot with an intensity of 80% AMT of the left FDI immediately following the completion of cTBS.

For each protocol, to assess the bilateral suprahyoid motor cortex excitability, the MEPs of both sides of the suprahyoid muscles were recorded via single-pulse TMS from the contralateral hemispheric motor hot spot before stimulation (baseline) and 0, 15, and 30 min after stimulation based on previous TMS studies. Thus, the left SMEP reflected the right suprahyoid motor cortex excitability and vice versa. To avoid inadvertent facilitation of the cortically evoked responses, all subjects remained as relaxed as possible and minimized swallowing, coughing or speaking during the study. Ten SMEPs for each side of the suprahyoid muscles were recorded per time point.

2.8 Data analysis

The peak-to-peak amplitude of the SMEP was used as a measure of motor cortex excitability. The MEP latency comprised the interval between the stimulus onset and the MEP onset. The individual SMEPs were reviewed with signal software (Yiruide CCY-IA, Wuhan, China), and ten traces were analyzed per time point. The average latency and amplitude of the SMEPs at each time point were determined as the final value. For every participant, the latencies of the left or right SMEPs recorded at each time point had minimal differences; the SMEPs with clearly different latencies were discarded prior to the calculation of the mean value, as well as the SMEPs recorded when the participant was coughing, swallowing, speaking, or moving his or her head.

2.9 Statistical analysis

Statistical tests were conducted using SPSS version 19.0 (IBM Corporation, Somers, NY). The data are presented as the mean $\pm$ standard deviation (SD) for the SMEP latency and amplitude. The left and right SMEP latencies and amplitudes were

independently analyzed for each TBS protocol using repeated measures one-way analysis of variance (ANOVA) with time (four levels: baseline, 0, 15, and 30 min post-TBS) as the within-subject factor. Sphericity was assumed according to Mauchly's test, and the Greenhouse-Geisser correction was used when sphericity was violated. A P -value of ≤ 0.05 was used to indicate statistical significance. We also used a Bonferroni correction to adjust for multiple comparisons (six comparisons were corrected) and considered P -values ≤ 0.0083 ($0.05/6$) to be statistically significant.

3. Results

The subjects did not report side effects during or after the experiment. Less than one percent of MEP trials were discarded prior to the calculation of the mean value. The SMEP elicited by single-pulse TMS from the contralateral suprahypoid motor cortex is shown in Figure 1. The FDI AMTs (expressed as the mean % of the MSO) accessed in every protocol were similar (Table 1), as well as the stimulation intensities used in every protocol (Table 2).

3.1 Effects of cTBS on suprahypoid motor cortical excitability

Following cTBS to the left suprahypoid motor cortex, the left SMEP amplitudes were significantly different across time points ($F_{(2.185, 74.277)} = 10.930$, $P < 0.001$) (Table 3). Multiple comparisons indicated that the SMEP amplitudes were significantly increased at 0 min ($P = 0.004$, 95% confidence interval [CI] -77.20 to -11.20), 15 min ($P = 0.005$, 95% CI = -90.92 to -11.89), and 30 min ($P = 0.001$, 95%

CI = -97.43 to -19.88) compared with the baseline. The right SMEP amplitudes were also significantly different across time points ($F_{(2.253, 76.596)} = 22.537$, $P < 0.001$). The SMEP amplitudes were significantly smaller at 0 min ($P < 0.001$, 95% CI = 28.77 to 85.12), 15 min ($P < 0.001$, 95% CI = 38.04 to 86.82), and 30 min ($P < 0.001$, 95% CI = 26.75 to 88.51) compared with the baseline (Fig. 2).

3.2 Effects of iTBS on suprahyoid motor cortical excitability

Following iTBS to the left suprahyoid motor cortex, the right SMEP amplitudes were significantly different across time points ($F_{(2.015, 68.514)} = 12.384$, $P < 0.001$) (Table 3). Multiple comparisons indicated that the SMEP amplitudes were significantly increased at 0 min ($P < 0.001$, 95% CI = -66.77 to -23.48), 15 min ($P < 0.001$, 95% CI = -70.33 to -25.22), and 30 min ($P = 0.001$, 95% CI = -90.89 to -18.88) compared with the baseline. No iTBS effect was identified on the left SMEP amplitudes ($F_{(2.185, 74.277)} = 0.470$, $P > 0.05$; Fig. 3).

3.3 Effects of cTBS followed by iTBS on suprahyoid motor cortical excitability

After cTBS to the left suprahyoid motor cortex followed by iTBS to the right suprahyoid motor cortex, the left SMEP amplitudes were significantly different across time points ($F_{(3, 102)} = 21.211$, $P < 0.001$) (Table 3). Multiple comparisons indicated that the left SMEP amplitudes were significantly increased at 0 min ($P < 0.001$, 95% CI = -62.61 to -21.05), 15 min ($P < 0.001$, 95% CI = -66.36 to -25.98), and 30 min ($P < 0.001$, 95% CI = -72.18 to -30.57) compared with the baseline. The right SMEP amplitudes were also significantly different across time points ($F_{(2.193, 74.552)} = 5.531$, $P = 0.005$). However, multiple comparisons indicated there was no difference between

the amplitudes at different time points (Fig. 4).

3.4 Effects of TBS on the latencies of the left and right SMEPs

The latencies of the left and right SMEPs recorded in every protocol are shown in Table 4. There was no significant difference in the latencies of the left or right SMEPs in every protocol.

4. Discussion

To the best of our knowledge, the current study provides the first evidence regarding TBS effects on suprahyoid motor cortex excitability in healthy subjects. The findings indicated that after the left suprahyoid motor cortex was stimulated by cTBS, the ipsilateral suprahyoid motor cortex excitability was suppressed, whereas the contralateral suprahyoid motor cortex excitability was enhanced. Moreover, after the left suprahyoid motor cortex was stimulated by iTBS, the ipsilateral suprahyoid motor cortex excitability was facilitated; however, the contralateral suprahyoid motor cortex excitability was not influenced. Furthermore, the bilateral suprahyoid motor cortex excitability was up-regulated after the left suprahyoid motor complex was stimulated by cTBS, followed by stimulation of the right suprahyoid motor cortex by iTBS. TBS had no effect on the MEP latencies of the bilateral suprahyoid muscles. The effect of TBS on the suprahyoid motor cortex excitability lasted a minimum of 30 min.

4.1 Mechanisms of cTBS effects on suprahyoid motor cortical excitability

In previous studies, cTBS rapidly suppressed human primary motor cortex

excitability for a minimum of 30 min [16, 26]. The mechanism may be related to GABAergic activity based on evidence using magnetic resonance spectroscopy that indicated cTBS of M1 increases the GABA concentration [27]. In accordance with previous studies, the current findings indicate cTBS effectively inhibited the ipsilateral suprahyoid motor cortex excitability. Interestingly, the reduction in the excitability was associated with a significant increase in the contralateral suprahyoid motor cortex excitability. Previous studies have confirmed that cortical activity depends on the balance between excitatory and inhibitory influences (for example, see [28-30]). We speculate that when the motor cortex excitability of the suprahyoid muscles was suppressed, the balance was broken; thus, the extent of the transcallosal inhibition to the contralateral motor cortex decreased. Consequently, the contralateral suprahyoid motor cortex was further activated. Corroborating evidence has originated from Bajwa et al. [31] who investigated the effects of 1 Hz rTMS on the corticospinal motor system and demonstrated that the suppression of the hand motor cortex in one hemisphere with 1 Hz rTMS facilitated the hand motor cortex in the contralateral hemisphere in healthy subjects.

However, in contrast to our results, Mistry et al. [20] did not identify an inhibitory effect of cTBS on the pharyngeal motor cortex in healthy subjects. This discrepancy may be attributed to different stimulation positions. In the present study, cTBS was applied to the left suprahyoid motor cortex, whereas Mistry et al. applied cTBS to the dominant pharyngeal motor cortex. According to previous research, the pharyngeal motor cortex exhibited a clear lateralization; however, the motor cortex of

the suprahyoid muscles presented less lateralization [32]. Thus, a single train of cTBS with the same intensity may not sufficiently suppress the dominant pharyngeal motor cortex excitability.

4.2 Mechanisms of iTBS effects on suprahyoid motor cortical excitability

In the current study, iTBS was applied to the left suprahyoid motor cortex, and the results demonstrated an increased excitability of the ipsilateral suprahyoid motor cortex, with no significant change in the contralateral suprahyoid motor cortex excitability. It is likely that the circuitry in the bilateral suprahyoid motor cortex is similar to the circuitry between the bilateral pharyngeal motor cortex, which is more easily activated than suppressed [33-36]. Moreover, high-frequency rTMS has been demonstrated to suppress the activity of the intracortical inhibitory circuitry and increase the excitability of the excitatory circuitry [37]. We hypothesized that when the suprahyoid motor cortex excitability in one hemisphere improves, the excitability of the homologous area in the contralateral hemisphere would decrease via an increase in interhemispheric inhibition.

Surprisingly, the contralateral motor cortex excitability did not change in the current study. Gow et al. [34] determined that ipsilateral and contralateral pharyngeal motor cortex excitability was up-regulated by 5 Hz rTMS to the dominant pharyngeal motor cortex in healthy subjects, with a greater increase in the cortico-pharyngeal excitability on the ipsilateral versus contralateral side. This finding may be related to the iTBS intensity, which was limited to 80% of the active motor threshold of the hand motor cortex according to international safety guidelines for the use of repetitive

TMS in humans [38]. Inconsistent with the current findings, a prior study demonstrated that iTBS of the hemisphere with stronger pharyngeal projections induced increased responses only in the contralateral weaker projection 60-90 min post-iTBS, with no change in the stronger hemisphere responses; furthermore, iTBS of the weaker projections had no significant effect on either hemisphere [19]. However, the small sample size may account for these previous findings. Additionally, the application of iTBS to different areas may comprise an alternative explanation for this inconsistency. Further investigation is needed to clarify the mechanism that drives these changes in cortical excitability.

4.3 Mechanisms of cTBS followed by iTBS effects on suprahyoid motor cortical excitability

In protocol A, following cTBS administration to the left suprahyoid motor cortex, the ipsilateral motor cortex excitability decreased, whereas the contralateral motor cortex excitability increased. Questions arose as to why the left suprahyoid motor cortex excitability did not decrease to a further degree but rather abolished the previous inhibitory effect and even tended to increase after iTBS was applied to the right suprahyoid motor cortex (multiple comparisons by the Bonferroni method indicated there was no difference in the four time points, whereas multiple comparisons by the LSD method indicated the left suprahyoid motor cortex excitability significantly increased at 15 and 30 min after the stimulation), as well as how iTBS could reverse virtual-lesion-induced cortical suppression via cTBS. Metaplasticity and homeostatic plasticity may explain the current results. TBS is

thought to invoke long-term depression (LTD) or long-term potentiation (LTP) at synapses [16, 39]. According to the sliding threshold idea proposed in the Bienenstock-Cooper-Munroe rule, high levels of synaptic activity indicate circumstances that increase the possibility to induce LTP. However, compensatory changes cause the suppression of LTP and an enhanced possibility of LTD. Conversely, low levels of synaptic activity shift the threshold toward greater LTP and less LTD [40-42]. Thus, synaptic plasticity is activity-dependent. This type of activity-dependent modulatory plasticity has been referred to as “metaplasticity” [43]. Metaplasticity maintains synapses within a dynamic functional range and prevents them from entering states of saturated LTP or LTD, which thus reach homeostasis [44, 45].

Transcranial stimulation techniques have successfully demonstrated homeostatic BCM-like metaplasticity mechanisms in the human motor cortex, which manage the threshold for inducing LTP-like or LTD-like plasticity [46-49]. Previous studies have confirmed that the excitability-enhancing pre-conditioning of M1 increased the efficacy of a subsequent excitability-reducing session of low-frequency rTMS [46, 49]. Murakami et al. [50] investigated metaplasticity via interactions of priming TBS with LTP/LTD-like plasticity induced by subsequent test TBS. The findings demonstrated that non-primed iTBS increased the MEP amplitude, but non-primed cTBS decreased the MEP amplitude. Furthermore, the pairing of identical protocols (iTBS-primed iTBS, cTBS-primed cTBS) resulted in the suppression of the test TBS effects on the MEP amplitude, whereas the pairing of different protocols

(cTBS-primed iTBS, iTBS-primed cTBS) enhanced the test TBS effects on the MEP amplitude. In the present study, a primed cTBS to the left motor cortex of the suprahyoid muscles led to excitability-reducing preconditioning of the ipsilateral motor cortex (i.e., the threshold toward greater LTP) and excitability-enhancing pre-conditioning of the contralateral motor cortex (i.e., the threshold toward greater LTD). Therefore, we speculated that when iTBS was applied to the right suprahyoid motor cortex after cTBS application to the left region, the increase in the ipsilateral motor cortex excitability was restrained, whereas the increase in the contralateral corresponding motor cortex excitability was facilitated, and the interhemispheric interactions between the suprahyoid motor cortices would eventually restore balance. Our speculation is supported by the findings of Jefferson et al. [51] who demonstrated that 5 Hz rTMS completely reversed a virtual lesion induced by 1 Hz rTMS in the contralateral pharyngeal motor cortex. This finding indicates that the effect of rTMS on motor cortex is not fixed and is affected by the excitability state of the motor cortex before and during stimulation.

4.4 TBS effect on the suprahyoid MEP latency

The suprahyoid MEP latency reflects the conduction time from the suprahyoid motor cortex to the suprahyoid muscles. For most unilateral post-stroke dysphagic patients, no MEP or longer latency and smaller amplitude MEPs were detected from the affected hemisphere [12, 13]. Dysphagia is related to the degree of damage to projections from the affected hemisphere [12]. Consistent with previous studies, the current findings demonstrated there was no TBS effect on the bilateral SMEP

latencies. Based on the evidence demonstrated by Gow et al. [34], we speculated that the neuroplastic effects induced by TBS occurred at a cortical level, but not a subcortical level.

4.5 Advantage of the neuronavigation system

In most previous studies, the target point was marked on a cap taped over the subject's head. However, this method cannot ensure that the cap position is exactly the same as the previous time. Furthermore, it does not take into account the substantial inter-subject variability in cortex morphology. In the present study, a neuronavigation system was applied to facilitate an accurate TMS position. Through the 3D reconstruction of the head and brain anatomy in real time on a computer screen, the neuronavigation system allows the visualization of the location and orientation of the coil over the head, as well as the resulting electric field density in the cortex [52, 53]. A comparison of the navigated and non-navigated TMS for motor cortex mapping indicated that all stimulation points with navigated TMS were less than 1.8 mm from the target, whereas the points were up to 60.2 mm away from the target with non-navigated TMS [52]. In addition, this approach allows the stimulation to be repeated to previously determined targets, which is crucial to ensure constant coil placement throughout the experiment. Thus, the neuronavigation system greatly improves the accuracy and repeatability of TMS, which therefore improves the reliability of the results.

4.6 Rationale for targeting the left hemisphere

The lateralization of function proposes that swallowing is predominantly mediated

by the right or left hemisphere, which is known as the ‘dominant’ hemisphere. In a prospective study of unilateral stroke patients (n=79), Shanahan et al. [54] identified ‘swallowing apraxia’ in a small subset of left hemisphere damaged (LHD) patients but not in right hemisphere damaged (RHD) individuals. Hamdy et al. [32] determined that the pharyngeal motor cortex exhibited a clear lateralization; however, the motor cortex of the suprahyoid muscles presented less lateralization. Plowman [55] mapped the swallowing musculature in the human motor cortex of 20 healthy adult volunteers and determined that the majority of subjects lateralized to the left hemisphere for the pharynx, right suprahyoid and left suprahyoid muscle sites. Therefore, in the present study, we chose the left hemisphere as the target, which will make it feasible for us to further confirm the TBS effects via functional magnetic resonance imaging.

4.7 Heterogeneity in the responses followed by TBS

In every protocol, heterogeneity in the responses followed by TBS was identified in no more than five subjects. After the left suprahyoid motor cortex was stimulated by cTBS, the left suprahyoid motor cortex excitability was facilitated rather than suppressed in five subjects. After the left suprahyoid motor cortex was stimulated by iTBS, the left suprahyoid motor cortex excitability was inhibited rather than enhanced in three subjects. And the responses to cTBS followed by iTBS were also inconsistent, with five subjects reacting in precisely the opposite way. This heterogeneity is likely associated with contractions of the suprahyoid muscles, such as coughing, swallowing, speaking, or head movements that occurred prior to the stimulation of TBS. Evidence indicates that prior tonic voluntary contraction or prior phasic movements may

influence the response to TBS. Without prior tonic voluntary contraction or prior phasic movements, cTBS elicited inhibitory and iTBS elicited facilitatory after-effects. However, following a short period of voluntary contraction or phasic movements, the after-effects of both cTBS and iTBS were reversed [56, 57]. These findings demonstrated that prior behavioral engagement of a cortical region may influence the response to a subsequent period of TBS in a homeostatic manner.

4.8 Study limitations and future directions

There are several limitations that should be considered in the interpretation of the findings. First, all subjects involved in this study were healthy, young adults. Future research should be conducted in different age groups, as well as post-stroke dysphagic patients to obtain a systematic and comprehensive conclusion. Second, the MEP recording was only maintained for 30 minutes after stimulation. The long-term effects of TBS on suprahyoid motor cortex excitability in healthy and dysphagic patients require further investigation. Future research should use functional magnetic resonance imaging (fMRI) and diffusion tensor imaging (DTI) to confirm the TBS effects on the swallowing neural network and focus on identifying the optimal TBS treatment protocols in post-stroke dysphagia.

4.9 Clinical significance of the current study

For unilateral stroke patients, the affected cortical excitability is decreased, whereas the intact cortical excitability is increased. Furthermore, there is an abnormally high interhemispheric inhibition from the intact hemisphere to the affected hemisphere, which further reduces the affected cortical excitability and

prevents the recovery of motor function. The modulation of excitability in motor regions of the intact and affected hemispheres contributes to improvements in motor function [14]. Thus, the current findings are of physiological and clinical significance and suggest that cTBS and iTBS have therapeutic potential to treat post-hemispheric stroke dysphagia when cTBS is applied to the intact hemisphere or iTBS is applied to the affected hemisphere to restore the excitatory and inhibitory balance of interhemispheric interactions. What is noteworthy is that the effects of TBS are likely influenced by preconditioning cortical excitability changes. Using TBS-primed TBS effects to purposefully direct homeostatic metaplasticity to increase or decrease the probability of induction of LTP-/LTD-like plasticity may also be beneficial for the rehabilitation of unilateral hemispheric stroke dysphagia. Additional studies with large sample sizes are required to confirm the effects of TBS on aged and post-stroke dysphagic patients, as well as the optimal TBS protocol, stimulation duration, and intensity required to facilitate the recovery of swallowing function in post-stroke dysphagic patients.

5. Conclusions

In summary, this study is the first investigation to identify TBS effects on neuroplastic excitability changes in the human suprahyoid motor cortex in healthy subjects. The findings have demonstrated that TBS effectively regulates the suprahyoid motor cortex excitability, with cTBS downregulating it and iTBS upregulating it. Inhibition of the suprahyoid motor cortex excitability in one

hemisphere can lead to further activation of the corresponding motor cortex in the other hemisphere. Furthermore, iTBS may abolish the inhibitory effect caused by cTBS applied to the contralateral suprahyoid motor cortex. The current study provides critical information to help clarify the TBS effects on suprahyoid motor cortex excitability and the potential regulatory mechanism in healthy subjects. These findings are important for the therapeutic application of TBS in neurogenic dysphagia. Additional studies are required to identify the optimal TBS protocols in the rehabilitation of post-stroke dysphagia.

Acknowledgements

We thank all subjects to participate in the experiments.

References

- [1] Martino R, Foley N, Bhogal S, Diamant N, Speechley M, Teasell R. Dysphagia after stroke: incidence, diagnosis, and pulmonary complications. *Stroke; a journal of cerebral circulation*. 2005;36:2756-2763.
- [2] Horner J, Massey EW, Riski JE, Lathrop DL, Chase KN. Aspiration following stroke: clinical correlates and outcome. *Neurology*. 1988;38:1359-1362.
- [3] Horner J, Buoyer FG, Alberts MJ, Helms MJ. Dysphagia following brain-stem stroke. Clinical correlates and outcome. *Archives of neurology*. 1991;48:1170-1173.
- [4] Nam HS, Beom J, Oh BM, Han TR. Kinematic effects of hyolaryngeal electrical stimulation therapy on hyoid excursion and laryngeal elevation. *Dysphagia*. 2013;28:548-556.
- [5] Beom J, Kim SJ, Han TR. Electrical Stimulation of the Suprahyoid Muscles in Brain-injured Patients with Dysphagia: A Pilot Study. *Annals of rehabilitation medicine*. 2011;35:322-327.
- [6] Speyer R, Baijens L, Heijnen M, Zwijnenberg I. Effects of therapy in oropharyngeal dysphagia by speech and language therapists: a systematic review. *Dysphagia*. 2010;25:40-65.
- [7] Rofes L, Vilardell N, Clave P. Post-stroke dysphagia: progress at last. *Neurogastroenterology and motility : the official journal of the European Gastrointestinal Motility Society*. 2013;25:278-282.

- [8] Foley N, Teasell R, Salter K, Kruger E, Martino R. Dysphagia treatment post stroke: a systematic review of randomised controlled trials. *Age and ageing*. 2008;37:258-264.
- [9] Khedr EM, Abo-Elfetoh N, Rothwell JC. Treatment of post-stroke dysphagia with repetitive transcranial magnetic stimulation. *Acta neurologica Scandinavica*. 2009;119:155-161.
- [10] Verin E, Leroi AM. Poststroke dysphagia rehabilitation by repetitive transcranial magnetic stimulation: a noncontrolled pilot study. *Dysphagia*. 2009;24:204-210.
- [11] Kim L, Chun MH, Kim BR, Lee SJ. Effect of repetitive transcranial magnetic stimulation on patients with brain injury and Dysphagia. *Annals of rehabilitation medicine*. 2011;35:765-771.
- [12] Khedr EM, Abo-Elfetoh N, Ahmed MA, Kamel NF, Farook M, El Karn MF. Dysphagia and hemispheric stroke: a transcranial magnetic study. *Neurophysiologie clinique = Clinical neurophysiology*. 2008;38:235-242.
- [13] Gallas S, Moirrot P, Debono G, Navarre I, Denis P, Marie JP, Verin E. Mylohyoid motor-evoked potentials relate to swallowing function after chronic stroke dysphagia. *Neurogastroenterology and motility : the official journal of the European Gastrointestinal Motility Society*. 2007;19:453-458.
- [14] Hummel FC, Cohen LG. Non-invasive brain stimulation: a new strategy to improve neurorehabilitation after stroke? *Lancet neurology*. 2006;5:708-712.
- [15] Park JW, Oh JC, Lee JW, Yeo JS, Ryu KH. The effect of 5Hz high-frequency rTMS over contralesional pharyngeal motor cortex in post-stroke oropharyngeal dysphagia: a randomized controlled study. *Neurogastroenterology and motility : the official journal of the European Gastrointestinal Motility Society*. 2013;25:324-e250.
- [16] Huang YZ, Edwards MJ, Rounis E, Bhatia KP, Rothwell JC. Theta burst stimulation of the human motor cortex. *Neuron*. 2005;45:201-206.
- [17] Talelli P, Greenwood RJ, Rothwell JC. Exploring Theta Burst Stimulation as an intervention to improve motor recovery in chronic stroke. *Clinical neurophysiology : official journal of the International Federation of Clinical Neurophysiology*. 2007;118:333-342.
- [18] Szaflarski JP, Vannest J, Wu SW, DiFrancesco MW, Banks C, Gilbert DL. Excitatory repetitive transcranial magnetic stimulation induces improvements in chronic post-stroke aphasia. *Medical science monitor : international medical journal of experimental and clinical research*. 2011;17:CR132-139.
- [19] Mistry S, Michou E, Rothwell J, Hamdy S. Remote effects of intermittent theta burst stimulation of the human pharyngeal motor system. *The European journal of neuroscience*. 2012;36:2493-2499.
- [20] Mistry S, Verin E, Singh S, Jefferson S, Rothwell JC, Thompson DG, Hamdy S. Unilateral suppression of pharyngeal motor cortex to repetitive transcranial magnetic stimulation reveals functional asymmetry in the hemispheric projections to human swallowing. *The Journal of physiology*. 2007;585:525-538.

- [21] Oldfield RC. The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia*. 1971;9:97-113.
- [22] Capotosto P, Babiloni C, Romani GL, Corbetta M. Differential contribution of right and left parietal cortex to the control of spatial attention: a simultaneous EEG-rTMS study. *Cerebral cortex*. 2012;22:446-454.
- [23] Babiloni C, Vecchio F, Miriello M, Romani GL, Rossini PM. Visuo-spatial consciousness and parieto-occipital areas: a high-resolution EEG study. *Cerebral cortex*. 2006;16:37-46.
- [24] Plowman-Prine EK, Triggs WJ, Malcolm MP, Rosenbek JC. Reliability of transcranial magnetic stimulation for mapping swallowing musculature in the human motor cortex. *Clinical neurophysiology : official journal of the International Federation of Clinical Neurophysiology*. 2008;119:2298-2303.
- [25] Rossini PM, Berardelli A, Deuschl G, Hallett M, Maertens de Noordhout AM, Paulus W, Pauri F. Applications of magnetic cortical stimulation. *The International Federation of Clinical Neurophysiology. Electroencephalography and clinical neurophysiology. Supplement*. 1999;52:171-185.
- [26] Huang YZ, Chen RS, Rothwell JC, Wen HY. The after-effect of human theta burst stimulation is NMDA receptor dependent. *Clinical neurophysiology : official journal of the International Federation of Clinical Neurophysiology*. 2007;118:1028-1032.
- [27] Stagg CJ, Wylezinska M, Matthews PM, Johansen-Berg H, Jezzard P, Rothwell JC, Bestmann S. Neurochemical effects of theta burst stimulation as assessed by magnetic resonance spectroscopy. *Journal of neurophysiology*. 2009;101:2872-2877.
- [28] Chen R. Interactions between inhibitory and excitatory circuits in the human motor cortex. *Experimental brain research*. 2004;154:1-10.
- [29] Gerloff C, Cohen LG, Floeter MK, Chen R, Corwell B, Hallett M. Inhibitory influence of the ipsilateral motor cortex on responses to stimulation of the human cortex and pyramidal tract. *The Journal of physiology*. 1998;510 (Pt 1):249-259.
- [30] Di Lazzaro V, Oliviero A, Profice P, Insola A, Mazzone P, Tonali P, Rothwell JC. Direct demonstration of interhemispheric inhibition of the human motor cortex produced by transcranial magnetic stimulation. *Experimental brain research*. 1999;124:520-524.
- [31] Bajwa S, BERPohl F, Rigonatti SP, Pascual-Leone A, Boggio PS, Fregni F. Impaired interhemispheric interactions in patients with major depression. *The Journal of nervous and mental disease*. 2008;196:671-677.
- [32] Hamdy S, Aziz Q, Rothwell JC, Crone R, Hughes D, Tallis RC, Thompson DG. Explaining oropharyngeal dysphagia after unilateral hemispheric stroke. *Lancet*. 1997;350:686-692.
- [33] Hamdy S, Aziz Q, Rothwell JC, Hobson A, Thompson DG. Sensorimotor modulation of human cortical swallowing pathways. *The Journal of physiology*. 1998;506 (Pt 3):857-866.

- [34] Gow D, Rothwell J, Hobson A, Thompson D, Hamdy S. Induction of long-term plasticity in human swallowing motor cortex following repetitive cortical stimulation. *Clinical neurophysiology : official journal of the International Federation of Clinical Neurophysiology*. 2004;115:1044-1051.
- [35] Singh S, Mistry S, Jefferson S, Davies K, Rothwell J, Williams S, Hamdy S. A magnetic resonance spectroscopy study of brain glutamate in a model of plasticity in human pharyngeal motor cortex. *Gastroenterology*. 2009;136:417-424.
- [36] Michou E, Mistry S, Jefferson S, Singh S, Rothwell J, Hamdy S. Targeting unlesioned pharyngeal motor cortex improves swallowing in healthy individuals and after dysphagic stroke. *Gastroenterology*. 2012;142:29-38.
- [37] Wu T, Sommer M, Tergau F, Paulus W. Lasting influence of repetitive transcranial magnetic stimulation on intracortical excitability in human subjects. *Neuroscience letters*. 2000;287:37-40.
- [38] Rossi S, Hallett M, Rossini PM, Pascual-Leone A, Safety of TMS/SCG. Safety, ethical considerations, and application guidelines for the use of transcranial magnetic stimulation in clinical practice and research. *Clinical neurophysiology : official journal of the International Federation of Clinical Neurophysiology*. 2009;120:2008-2039.
- [39] Di Lazzaro V, Pilato F, Saturno E, Oliviero A, Dileone M, Mazzone P, Insola A, Tonali PA, Ranieri F, Huang YZ, Rothwell JC. Theta-burst repetitive transcranial magnetic stimulation suppresses specific excitatory circuits in the human motor cortex. *The Journal of physiology*. 2005;565:945-950.
- [40] Bienenstock EL, Cooper LN, Munro PW. Theory for the development of neuron selectivity: orientation specificity and binocular interaction in visual cortex. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 1982;2:32-48.
- [41] Stanton PK. LTD, LTP, and the sliding threshold for long-term synaptic plasticity. *Hippocampus*. 1996;6:35-42.
- [42] Narayanan R, Johnston D. The h current is a candidate mechanism for regulating the sliding modification threshold in a BCM-like synaptic learning rule. *Journal of neurophysiology*. 2010;104:1020-1033.
- [43] Abraham WC, Bear MF. Metaplasticity: the plasticity of synaptic plasticity. *Trends in neurosciences*. 1996;19:126-130.
- [44] Abraham WC, Tate WP. Metaplasticity: a new vista across the field of synaptic plasticity. *Progress in neurobiology*. 1997;52:303-323.
- [45] Muller-Dahlhaus F, Ziemann U. Metaplasticity in Human Cortex. *The Neuroscientist : a review journal bringing neurobiology, neurology and psychiatry*. 2015;21:185-202.
- [46] Iyer MB, Schleper N, Wassermann EM. Priming stimulation enhances the depressant effect of low-frequency repetitive transcranial magnetic stimulation. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2003;23:10867-10872.

- [47] Lang N, Siebner HR, Ernst D, Nitsche MA, Paulus W, Lemon RN, Rothwell JC. Preconditioning with transcranial direct current stimulation sensitizes the motor cortex to rapid-rate transcranial magnetic stimulation and controls the direction of after-effects. *Biological psychiatry*. 2004;56:634-639.
- [48] Muller JF, Orekhov Y, Liu Y, Ziemann U. Homeostatic plasticity in human motor cortex demonstrated by two consecutive sessions of paired associative stimulation. *The European journal of neuroscience*. 2007;25:3461-3468.
- [49] Siebner HR, Lang N, Rizzo V, Nitsche MA, Paulus W, Lemon RN, Rothwell JC. Preconditioning of low-frequency repetitive transcranial magnetic stimulation with transcranial direct current stimulation: evidence for homeostatic plasticity in the human motor cortex. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 2004;24:3379-3385.
- [50] Murakami T, Muller-Dahlhaus F, Lu MK, Ziemann U. Homeostatic metaplasticity of corticospinal excitatory and intracortical inhibitory neural circuits in human motor cortex. *The Journal of physiology*. 2012;590:5765-5781.
- [51] Jefferson S, Mistry S, Michou E, Singh S, Rothwell JC, Hamdy S. Reversal of a virtual lesion in human pharyngeal motor cortex by high frequency contralesional brain stimulation. *Gastroenterology*. 2009;137:841-849, 849 e841.
- [52] Julkunen P, Saisanen L, Danner N, Niskanen E, Hukkanen T, Mervaala E, Kononen M. Comparison of navigated and non-navigated transcranial magnetic stimulation for motor cortex mapping, motor threshold and motor evoked potentials. *NeuroImage*. 2009;44:790-795.
- [53] Ahdab R, Ayache SS, Brugieres P, Goujon C, Lefaucheur JP. Comparison of "standard" and "navigated" procedures of TMS coil positioning over motor, premotor and prefrontal targets in patients with chronic pain and depression. *Neurophysiologie clinique = Clinical neurophysiology*. 2010;40:27-36.
- [54] Shanahan T, Logemann J, Colangelo L. Swallowing function and lesion site in stroke patients. *ASHA*. 1995;37:90.
- [55] Plowman EK. Reproducibility of transcranial magnetic stimulation for mapping swallowing musculature in the human motor cortex. unpublished. 2005.
- [56] Gentner R, Wankerl K, Reinsberger C, Zeller D, Classen J. Depression of human corticospinal excitability induced by magnetic theta-burst stimulation: evidence of rapid polarity-reversing metaplasticity. *Cerebral cortex*. 2008;18:2046-2053.
- [57] Iezzi E, Conte A, Suppa A, Agostino R, Dinapoli L, Scontrini A, Berardelli A. Phasic voluntary movements reverse the aftereffects of subsequent theta-burst stimulation in humans. *Journal of neurophysiology*. 2008;100:2070-2076.

Figure Legend

Fig. 1. SMEP waveform of one representative participant elicited by single-pulse TMS from the contralateral suprahyoid motor cortex.

Fig. 2. Effects of cTBS on suprahyoid motor cortex excitability. Changes in SMEP amplitudes from baseline following cTBS to the left suprahyoid motor cortex. Left SMEPs are activated immediately and for up to 30 min, whereas right SMEPs are inhibited immediately and for up to 30 min (* $P \leq 0.0083$, ** $P \leq 0.001$, post hoc tests). Thus, following the cTBS intervention, the left suprahyoid motor cortex excitability was suppressed, whereas the right suprahyoid motor cortex excitability was activated. Values represent means $\pm$ SEM.

Fig. 3. Effects of iTBS on suprahyoid motor cortex excitability. Changes in the SMEP amplitudes from baseline following iTBS to the left suprahyoid motor cortex. No effect was identified in the left SMEPs; the right SMEPs increased immediately and for up to 30 min (** $P \leq 0.001$, post hoc tests). Thus, following iTBS intervention, the left suprahyoid motor cortex excitability was increased, whereas the right suprahyoid motor cortex excitability was not affected. Values represent means $\pm$ SEM.

Fig. 4. Effects of cTBS followed by iTBS on suprahyoid motor cortex excitability. Changes in the SMEP amplitudes from baseline after cTBS to the left suprahyoid motor cortex followed by iTBS to the right suprahyoid motor cortex. The left SMEPs are increased immediately and for up to 30 min (** $P \leq 0.001$, post hoc tests). The

right SMEPs also exhibited a trend towards an increase; however, multiple comparisons indicated there was no difference between the amplitudes at different time points. Thus, after the cTBS followed by iTBS intervention, the left suprahyoid motor cortex excitabilities were increased, and the right suprahyoid motor cortex excitabilities tended to increase. Values represent means $\pm$ SEM.

Table 1. FDI AMT accessed in every protocol

	Protocol A	Protocol B	Protocol C	
	RFDI	RFDI	RFDI	LFDI
AMT	36.5 ± 1.37	37.4 ± 1.43	35.9 ± 1.48	36.7 ±
(%MSO)				1.41

RFDI: right FDI

LFDI: left FDI

Values indicate mean ± SEM.

Table 2. Stimulation intensity used in every protocol

	Protocol A	Protocol B	Protocol C	
	cTBS	iTBS	cTBS	iTBS
Stimulation				
intensity (% MSO)	29.2 ± 1.10	29.9 ± 1.14	28.7 ± 1.18	29.3 ± 1.12

Values indicate mean ± SEM.

Table 3. Amplitudes of the left and right SMEPs recorded
in every protocol (μV , $\bar{x} \pm s$)

	Baseline	0 min	15 min	30 min	<i>F</i>	<i>P</i>
Protocol A						
Left	292 $\pm$ 153	337 $\pm$ 165	344 $\pm$ 164	351 $\pm$ 161	10.930	<0.001*
Right	330 $\pm$ 170	273 $\pm$ 132	267 $\pm$ 143	272 $\pm$ 134	22.537	<0.001*
Protocol B						
Left	325 $\pm$ 137	316 $\pm$ 123	317 $\pm$ 124	319 $\pm$ 131	0.470	0.704
Right	307 $\pm$ 145	352 $\pm$ 137	355 $\pm$ 143	362 $\pm$ 155	12.384	<0.001*
Protocol C						
Left	321 $\pm$ 151	362 $\pm$ 153	367 $\pm$ 160	372 $\pm$ 162	21.211	<0.001*
Right	327 $\pm$ 168	341 $\pm$ 186	349 $\pm$ 181	355 $\pm$ 191	5.531	0.005*

Left: left suprahyoid muscles. Right: right suprahyoid muscles.

* $P < 0.05$

Table 4. Latencies of the left and right SMEPs recorded
in every protocol (ms, $\bar{x} \pm s$)

	Baseline	0 min	15 min	30 min	<i>F</i>	<i>P</i>
Protocol A						
Left	7.75 $\pm$ 1.46	7.75 $\pm$ 1.41	7.73 $\pm$ 1.45	7.72 $\pm$ 1.52	0.198	0.846
Right	7.70 $\pm$ 1.43	7.69 $\pm$ 1.42	7.68 $\pm$ 1.44	7.70 $\pm$ 1.37	0.180	0.910
Protocol B						
Left	7.76 $\pm$ 1.40	7.79 $\pm$ 1.40	7.78 $\pm$ 1.37	7.77 $\pm$ 1.39	0.309	0.819
Right	7.73 $\pm$ 1.39	7.71 $\pm$ 1.37	7.71 $\pm$ 1.35	7.67 $\pm$ 1.43	0.773	0.512
Protocol C						
Left	7.81 $\pm$ 1.35	7.76 $\pm$ 1.37	7.78 $\pm$ 1.36	7.76 $\pm$ 1.33	0.855	0.467
Right	7.69 $\pm$ 1.32	7.68 $\pm$ 1.29	7.68 $\pm$ 1.33	7.63 $\pm$ 1.29	1.374	0.259

Left: left suprahyoid muscles. Right: right suprahyoid muscles.