RUNNING HEAD: Working Memory and Multitasking

On the Relation of Working Memory and Multitasking: Memory Span and Synthetic Work
Performance

Thomas S. Redick

Purdue University

Corresponding author information:

Thomas S. Redick
Purdue University
Psychological Sciences
703 Third Street
West Lafayette, IN 47907
tredick@redick.edu
Phone: 765-494-5132

Word Count: 4,498 (main text only)

3 Tables

1 Figure

Abstract

Recent research has identified working memory as a critical component of multitasking ability. These studies showed that working memory accounted for multitasking variance over-and-above that predicted by other cognitive, personality, and experience-based variables. However, a limitation of these previous studies was that the tasks selected to measure working memory were dual-tasks themselves. The purpose of the current research was to determine if working memory measures must be dual-tasks to predict multitasking performance, or if other types of working memory measures that do not rely upon the dual-task methodology predict multitasking just as well, if not better. Three different serial order memory span tasks (one dual-task and two single-task) and one multitask were administered to a sample of healthy young adults. The results showed that single- and dual-task working memory measures predicted multitasking to a similar degree. The results indicate there is something fundamental about working memory's relationship with multitasking ability.

KEYWORDS: working memory; multitasking; applied

On the Relation of Working Memory and Multitasking: Memory Span and Synthetic Work Performance

Working memory functioning is important for many activities both in and out of the lab. While experimental work has shown effects of working memory load upon decision-making, attention, and memory search, individual differences research has demonstrated relations between an individual's working memory level and reading comprehension, fluid intelligence, and mathematics ability. As another example of individual differences research on working memory, one outside-of-the-lab activity that applied cognitive psychologists have frequently investigated is multitasking. Recent research has identified working memory as a critical component of multitasking ability (Ackerman & Beier, 2007; Bühner, König, Pick, & Krumm, 2006; Colom, Martínez-Molina, Shih, & Santacreu, 2010; Hambrick, Oswald, Darowski, Rench, & Brou, 2010; König, Bühner, & Murling, 2005; Morgan et al., 2013), and these studies showed that working memory predicted multitasking performance over-and-above other cognitive, personality, and experience-based variables. However, a limitation of many of these previous studies was that the tasks selected to measure working memory were multitasks themselves. The purpose of the current research was to determine if working memory measures must be dual-tasks to predict multitasking performance, or if other types of working memory measures that do not rely upon the dual-task methodology predict multitasking just as well, if not better. If single-task working memory measures predict multitasking performance, this implies that there is something fundamental about working memory's relationship with multitasking ability, irrespective of any overlap of the method used to measure both working memory and multitasking.

Memory Span Measures of Working Memory

All of the aforementioned studies included complex memory span tasks as working memory measures, in which participants must remember a sequence of items while also performing an interleaved distracting task. For example, in the reading span, examinees read sentences and make a decision about the veracity or sensibility of the statement, and are presented with an item to remember (e.g., letter). A number of these processing-and-storage stimuli are presented, before participants are asked to recall the to-be-remembered items in order. Tasks like reading span are often called processing-and-storage or complex span tasks in contrast to storage-only, simple span tasks such as letter span, in which participants are instructed to serially recall a sequence of letters in order without the additional interleaved distractor task. One current view is that individual differences in working memory, as measured by complex span tasks, reflect a person's ability to maintain a select number of items in active memory, retrieve information from inactive memory, and control attention to counteract interference and distraction (McVay & Kane, 2012; Shipstead, Lindsey, Marshall, & Engle, 2014; Unsworth, Fukuda, Awh, & Vogel, 2014).

Although earlier studies suggested a distinction between the working memory processes measured by complex versus simple span tasks (Daneman & Carpenter, 1980; Engle, Tuholski, Laughlin, & Conway, 1999), more recent research has indicated that simple and complex span tasks measure largely overlapping processes and account for similar variance in cognitive abilities, particularly when partial credit-scoring and longer simple-span list lengths are used (Colom, Rebollo, Abad, & Shih, 2006; Unsworth & Engle, 2007). In addition, studies using variants of a running span task, in which participants must report a specified number of items at the end of a sequence of stimuli (e.g., letters), produce strong correlations with complex span

tasks and also account for shared variance in cognitive abilities (Broadway & Engle, 2010; Cowan et al., 2005).

Multitasking

Multitasking is important for many aspects of human behavior, particularly in certain employment sectors (e.g., medical field, military, aviation) where multitasking is necessary for vocational success and multitasking failures pose serious safety-related consequences. In the current work, I adopted the definition of multitasking provided by Oswald, Hambrick, and Jones (2007), whereby multitasking requires: (a) performing multiple tasks; (b) consciously shifting from one task to another; and (c) performing the component tasks over a relatively short time span. There are numerous operational definitions possible for multitasking, but in the current study I used the SynWin multitask, an established measure of synthetic work (Elsmore, 1994; Hambrick et al., 2010, 2011; Proctor, Wang, & Pick, 1998; Salthouse, Hambrick, Lukas, & Dell, 1996). SynWin involves simultaneous performance of four unrelated tasks varying in self- and externally-paced timing, visual and auditory information processing, and verbal and numerical stimuli (Figure 1). SynWin is thus similar to other tasks used in applied multitasking research, including the SIMKAP (Bratfisch & Hagman, 2003) and Multi-Attribute Task Battery (MATB; Comstock & Arnegard, 1992).

This performance-based way of measuring multitasking stands in contrast with alternatives such as self-report of the frequency of engaging in different types of multitasking (e.g., Ophir, Nass, & Wagner, 2009; Sanbonmatsu, Strayer, Medeiros-Ward, & Watson, 2013). This distinction is critical, because whereas previous research has shown that complex span tasks are strongly positively correlated with SynWin (e.g., Hambrick et al., 2010), self-report measures

of multitasking are either unrelated (Minear, Brasher, McCurdy, Lewis, & Younggren, 2013) or weakly negatively correlated (Sanbonmatsu et al., 2013) with complex span tasks.

In addition to Hambrick et al. (2010), two other studies examining SynWin and working memory performance are relevant for the current research. First, Hambrick et al. (2011) administered a verbal (digit) n-back task and SynWin to a large sample of Navy sailors. N-back working memory tasks require the participant to monitor a sequence of stimuli and determine whether or not the current stimulus matches the item presented N items back. Hambrick et al. (2011) observed that n-back and SynWin performance were positively correlated, and n-back accounted for significant SynWin variance over-and-above ASVAB test scores. Thus, digit n-back, a single-task verbal working memory measure that only weakly correlates with complex span working memory measures (Jaeggi, Buschkuhl, Perrig, & Meier, 2010; Kane, Conway, Miura, & Colflesh, 2007; Redick & Lindsey, 2013) also accounts for substantial multitasking performance, providing preliminary information that the relation between working memory and multitasking is not dependent upon a dual-task method of measuring working memory. A second study my colleagues and I conducted examined the effect of working memory training on a variety of cognitive outcomes (Redick et al., 2013). For the purpose of the current work, I re-analyzed the data from the pretest session, before participants were assigned to training and control groups. Among the tasks administered in the pretest session were symmetry span, running letter span, and SynWin. In $N = 72$ participants, the symmetry span-SynWin and running letter span-Synwin correlations were identical (both r 's = .37, both p 's < .01).¹ Again, these results suggest that the working memory and multitasking relation is of a similar magnitude independent of the single- or dual-task nature of the span task, although these data are not

conclusive considering symmetry span and running letter span differed in both methodology (dual- vs. single-task) and to-be-remembered content (nonverbal vs. verbal stimuli).

Current Study

The advantage for the current research is that although reading span is itself a dual-task, letter span and running letter span are not dual-tasks, and thus interpretation of the relationship with a multitasking criterion is more straightforward (Cowan et al., 2005) than when using combined processing-and-storage scores to predict multitasking performance (e.g., Colom et al., 2010). In addition, the to-be-remembered information and task required by the participant is the same in each memory span task (viz., recall letters in serial order), facilitating a fair comparison among the working memory measures within the same participants (compare to Hambrick et al., 2011, and Redick et al., 2013, above). If verbal memory span tasks without a secondary-task component do predict multitasking as well as complex span tasks, one can convincingly conclude that working memory and multitasking relationships observed in previous studies are robust across methodology, and working memory is an important contributor to multitasking ability independent of the exact measures administered.

An additional advantage of using variations of span tasks is that it permits a calculation and comparison of the contribution of different list lengths in relation to multitasking performance. For example, with the complex span tasks such as reading span, previous research has shown that the shortest list lengths accounted for as much variance in fluid intelligence as did the longest list lengths (Salthouse & Pink, 2008; Unsworth & Engle, 2006). In contrast, Unsworth and Engle (2006) showed that on simple span tasks such as letter span, the shortest list lengths had weak/non-significant relationships with fluid intelligence, but the correlation magnitude increased as list length increased.

The explanation provided by Unsworth and Engle (2006) is that the span tasks reflect a combination of maintenance in primary memory and retrieval from secondary memory, but different list lengths on different span tasks tap into these abilities in distinct ways. Complex span tasks reflect a combination of both active maintenance and retrieval at all list lengths, as the distractor processing task displaces to-be-remembered items from primary memory into secondary memory. Thus, when probed to recall the items on the current list, the most recently presented item may still be in primary memory because no subsequent distractor task displaced it from primary memory, but the other items must be retrieved from secondary memory. This occurs on both short and long list lengths – according to this explanation, what varies across list lengths in complex span tasks is the number of items needed to be retrieved from secondary memory.

In contrast, the shortest list lengths on simple span tasks can be maintained entirely within primary memory, according to estimates from Cowan (2001) and Luck and Vogel (2013) that most adults can actively maintain 4 ± 1 items. However, on longer list lengths in simple span tasks, the number of to-be-remembered items exceeds primary memory capacity, meaning that some items must be recalled from secondary memory and accessed from primary memory.

Thus, because short list lengths on simple span tasks measure mostly primary memory capacity, whereas longer list lengths on simple span tasks and all list lengths on complex span tasks measure both primary memory and secondary memory functioning, the former correlations with higher-order cognition should be weaker than the latter correlations. While this list-length dependent pattern of correlations has been observed for the simple and complex span tasks in relation to fluid intelligence (Unsworth & Engle, 2006), it has not been examined for other abilities, including multitasking.

In addition, I conducted list-length analyses for the running letter span task. Here, there was no previous research to explicitly compare against, but the prediction is that running letter span as administered here should be similar to the longer list lengths of the simple span tasks. Specifically, with the task across all running span trials to recall the last 4 items in order, again the necessary to-be-remembered items can be held within primary memory capacity. However, as Broadway and Engle (2010) showed, in the version of the running span used here, participants seem to try to remember all items presented, and then select only the necessary items at the end of the list when prompted to recall. This behavior would suggest that participants use both primary and secondary memory to hold all items until recall, and as such, all list lengths on the running span as administered here should moderately and equally correlate with SynWin.

Method

Participants

Sixty-five participants were recruited from a campus in the Midwestern United States, and were compensated with course credit. After providing demographics information, participants performed a multitask (SynWin; Elsmore, 1994) and three different memory span tasks, each of which required remembering a sequence of unrelated letters that were visually presented. The protocol was approved by the university IRB, and all participants provided informed consent.

Materials

SynWin. In SynWin, four simultaneous sub-tasks were presented, with each sub-task in its own quadrant of the screen, surrounding the central display of the participants' current total score (Figure 1). The four sub-tasks include: (a) probe-recognition memory; (b) arithmetic; (c) visual monitoring; and (d) auditory monitoring. All task properties and scoring formulas were

identical to those used in the ‘baseline’ condition in Hambrick et al. (2010, 2011). In the probe-recognition sub-task, six letters were presented briefly (5 s) at the beginning of the task block. For the remaining time in the block, the six letters were no longer visible, requiring the participant to hold the list in memory, although the participant had the option to ‘peek’ at the list again by clicking on the empty memory set box for a penalty of 10 points. Throughout the rest of the task block, a probe letter was presented every 10 s, and the participant made a yes/no decision whether the probe was in the memory set. Ten points were awarded for correct answers, ten points were subtracted for incorrect answers, and ten points were subtracted for misses (not responding to the probe letter within 5 seconds). In the arithmetic sub-task, participants added two 3-digit numbers and provided the final solution. A new problem was shown only after an answer was submitted – this was the only sub-task that in which the stimuli were presented based on the participant’s pace. Twenty points were awarded for correct answers, and ten points were subtracted for incorrect answers. There was no explicit penalty for not responding to the arithmetic sub-task. In the visual monitoring sub-task, a display analogous to a fuel gauge continuously (interval = 200 ms) dropped gradually from 100 to 0 and had to be reset to 100 by clicking on the gauge. Points were earned for responding before the gauge reached 0, and points were subtracted if the gauge reached 0. The scoring algorithm is such that participants received more (up to 10) points if they clicked to refuel when the gauge was closer to 0, versus clicking refuel when the gauge was still close to 100 – however, the penalty for letting the gauge fall to 0 was severe. If the gauge reached 0, a distinct alarm tone sounded once, and the current score that was displayed decreased as points were subtracted at a rate of 10 points per second until the gauge was reset. In the auditory monitoring sub-task, high- and low-frequency tones (2000 Hz and 1000 Hz, respectively) were presented every 10 s, and participants were instructed to click

on the quadrant when the infrequent (probability = .20) high-frequency tone was presented. Ten points were earned for hits, and ten points were subtracted for misses and false alarms.

Participants received instructions about each sub-task first. For practice, participants first completed 1 minute of each sub-task by itself, and then 1 minute of all four sub-tasks simultaneously. After completing practice and ensuring participants understood the task instructions, participants then completed two 5-minute blocks of all four sub-tasks simultaneously. Participants' total scores were determined by a formula that combined the points earned across all four sub-tasks in the two experimental blocks. Although the main analyses used the total score averaged across the two blocks, supplemental analyses used the individual sub-task scores averaged across the two blocks.

Memory span tasks. In all three tasks, participants saw letters presented individually and used mouse clicks to recall the letters on each trial in serial order. In all tasks, set sizes were randomly intermixed, in contrast to increasing set-size presentation, because of the randomized method's demonstrated improved predictive validity (St. Clair-Thompson, 2012). In all working memory tasks, letters were scored as correct if they were in the correct serial position, regardless of whether the entire set of letters on a given trial was recalled correctly. The partial-credit scoring method was used because of its established psychometric properties compared to alternative scoring techniques (Conway et al., 2005; Redick et al., 2012). Note that the conclusions reported below are not different if the all-or-nothing scoring methods are used for the span tasks instead.

In letter span, participants saw two lists each of set sizes 3-9 letters, with a maximum possible score of 84. Before completing the experimental trials, participants completed 4 practice trials with either 2 or 3 letters presented on each trial. In reading span, participants combined the

serial recall aspect of the letter span task with a sentence verification task requiring a true/false judgment (Redick et al., 2012). First, participants completed one practice trial of set size 2. Then, participants completed 15 practice trials of the sentence verification task only. Performance on the single-task condition of the sentence verification task was used to set an upper time limit for subsequent dual-task trials, using the method reported by Unsworth, Redick, Heitz, Broadway, and Engle (2009). Participants practiced 3 dual-task trials at set size 2, before proceeding to the experimental block of dual-task trials. Participants saw three lists each of set sizes 3-7 sentences and letters, with a maximum possible score of 75. In running span, participants saw between 4-7 letters on each trial, but were instructed to only recall the last 4 letters in order that they saw. In the practice trials, there was 1 trial at each set size; in the experimental trials, there were 3 trials at each set size, with a maximum possible score of 48. For each span task, list-length analyses were conducted by first calculating proportion correct for each trial (e.g., for list-length 4, a score of .75 means that 3 of 4 items were recalled in the correct serial position for that trial), and then averaging across all trials of that particular list length.

Results

Ten participants were excluded for the following reasons: (a) age (5 participants were > 40 years old; see Salthouse et al., 1996, for age-related impairments on SynWin); (b) missing data on a specific task due to experimenter error (3 participants); and (c) extreme negative scores on block 1 or block 2 of SynWin that were primarily produced by an extended period of time in which the gauge in the visual monitoring task was left on empty, producing a score of -830 for that sub-task for that block (2 participants). Z-scores confirmed these two participants as outliers; their mean visual monitoring performance produced z -scores of -5.03 and -5.02 (the participant with the next lowest visual monitoring score $z = -1.22$). The percentage of participants removed

from the current sample due to extremely poor SynWin performance was thus 3.1%, and similar to the rate of participants (3.8%) excluded by Hambrick et al. (2010). For the final 55 participants, 22 were male and 33 female with a mean age of 22.7 years ($SD = 5.2$).²

Descriptive statistics are presented in Table 1. All variables had acceptable values for skewness, kurtosis, and reliability, with the exception of the visual monitoring sub-task within SynWin. Mean performance on reading span was comparable to published normative data for young adults (Redick et al., 2012).

Before examining the relationships between the memory span tasks and SynWin, I first examined the interrelationships among the three memory span tasks. First, as seen in Table 2, the zero-order correlations among the span tasks were all strong and significant. Next, an exploratory factor analysis (principal-axis factor extraction) on the three memory span tasks clearly revealed a one-factor solution (eigenvalue = 2.06, variance = 68.6%). These results suggest the variance among the three memory span tasks is mostly overlapping, consistent with previous research (Broadway & Engle, 2010; Colom et al., 2006; Cowan et al., 2005; Unsworth & Engle, 2007).

Correlations for the three memory span tasks with SynWin performance were all significant (Table 2). More germane to the current research question, the correlations between the three span tasks and SynWin were nearly identical (all r 's = .48 to .52), and did not differ as a function of the single- or dual-task nature of the memory span task, as confirmed via Hotelling-Williams t -tests for dependent-samples correlations: letter span versus reading span, $t(52) = .33$, $p = .74$; running letter span versus reading span, $t(52) = .25$, $p = .81$.

In addition, I used relative weights analyses (Tonidandel & LeBreton, 2015) to determine whether the dual- versus single-task memory span tasks account for a different proportion of

multitasking performance. The raw relative weights for each predictor indicate what proportion of the overall variance can be attributed to each predictor variable, even if the predictor variables are intercorrelated. Using bootstrapping techniques (10,000 replications), one can then calculate confidence intervals testing if two particular predictor variables differ in the amount of variance accounted for in the criterion variable. First, the regression analyses indicated that letter span, running letter span, and reading span accounted for a combined 37.3% of the variance in SynWin overall scores, $F(3, 51) = 10.13, p < .01$. The relative weight analyses indicated that the relative weights, that is, the proportion of SynWin variance accounted for, did not differ between the dual and non-dual memory span tasks. Comparing letter span versus reading span, the bootstrap-generated confidence interval included 0 (95% CI: -.29 to .08). Similarly, comparing running letter span versus reading span, the bootstrap-generated confidence interval included 0 (95% CI: -.27, .13).

Finally, I conducted list-length analyses for each span task as described above. Descriptive statistics and correlations with SynWin are reported in Table 3. For all three span tasks, memory accuracy significantly decreased as list length increased – letter span: $F(6, 324) = 175.21, p < .01, \eta_p^2 = .76$; running letter span: $F(3, 162) = 44.18, p < .01, \eta_p^2 = .45$; reading span: $F(4, 216) = 34.39, p < .01, \eta_p^2 = .39$. Table 3 also shows that for letter span, the magnitude of the correlations with SynWin increases as list length increases, whereas for running span and reading span, the correlations with SynWin are significant and equal in magnitude for both short and longer list lengths.

As supplementary analyses, correlations between the memory span tasks and the SynWin sub-tasks are shown in Table 3. Analyses will not focus on the visual monitoring sub-task, given its high values of skewness and kurtosis, along with the lack of a correlation with the SynWin

total score. For the probe recognition and auditory monitoring sub-tasks, all memory span correlations were significant. For the arithmetic sub-task, only reading span was significantly correlated at $p < .05$ (running letter span: $p = .06$; letter span: $p = .13$). Critically, for the SynWin sub-tasks, Hotelling-Williams t -tests confirmed that none of the correlations significantly differed as a function of the dual versus non-dual nature of the memory span tasks (all t 's < 1.59 , all p 's $> .12$).

Discussion

Multitasking is an important yet nebulous activity that is present in many daily activities outside of the lab. Here, I chose a particular manifestation of multitasking, synthetic work. Working memory is strongly related to one's ability to multitask, yet one limitation of many previous studies is that dual-task measures of working memory have frequently been used as predictors of multitasking outcomes. The current work builds on recent accounts that complex span measures of working memory are not the only way to measure individual differences in working memory capacity (Colom et al., 2006; Cowan et al., 2005; Schmiedek, Hildebrandt, Lövdén, Wilhelm, & Lindenberger, 2009; Unsworth & Engle, 2007). Using three variations of memory span tasks with common to-be-remembered stimuli, correlation and regression results indicated that the single-task working memory measures accounted for similar multitasking variance compared to the dual-task measures. The list-length analyses replicate previous research that used fluid intelligence as the criterion variable (Unsworth & Engle, 2006), and provide further support for the idea that what is important for working memory's contribution to higher-order cognition is not the dual- or single-task nature of the task, but instead the cognitive abilities involved in performance of those tasks. The current work extends previous research showing working memory to be a strong predictor of multitasking, via list-length analyses indicating that

active maintenance in primary memory and retrieval from secondary memory contribute to the shared variance amongst the span tasks and synthetic work. This outcome furthers our knowledge of a poorly understood construct, multitasking, and could aid in the formation of less time-consuming ways to predict or improve multitasking in applied contexts (e.g., driving, air-traffic control, military settings). For example, lengthy intelligence tests and/or military cognitive batteries such as the ASVAB are currently used in personnel selection situations where multitasking is needed for job performance. The current work demonstrates that memory span tasks can predict synthetic work performance quicker than these existing methods, particularly if recently-developed short-form versions of the span tasks are used (Foster et al., 2015; Oswald, McAbee, Redick, & Hambrick, 2015).

Limitations and Future Directions

Although this was the first study to directly compare how different types of memory span tasks predict multitasking performance, there are limitations to the current work that could motivate future work. The current work shows that working memory is related to multitasking, but a stronger case for the specificity of this relationship would have been possible had non-working memory tasks been administered, too. In addition, I administered only verbal memory span tasks, but future work should also examine if nonverbal memory span tasks behave similarly in relation to multitasking. Given previous work showing that symmetry span, a nonverbal complex span task, correlates significantly with SynWin (Hambrick et al., 2010; Redick et al., 2013), I do not expect the to-be-remembered content to be a critical factor, but this remains an empirical question at this juncture. In addition, although Synwin has been used as a multitasking outcome analogous to the type of multitasking an individual might experience in the workplace, using alternative operational definitions of multitasking could provide different

results than those observed here. Finally, one as yet unexplored aspect of multitasking is identifying variables that account for extreme multitasking failures. Although one might be most interested in using predictor variables to identify those who are highest in multitasking performance for personnel selection purposes, a somewhat contrasting approach would be to identify those individuals who may be particularly unfit for jobs requiring multitasking. For example, whereas a multitasking failure in a typical workplace environment might mean that an email reply is slower than intended, in an air-traffic-control setting, a multitasking failure could easily result in numerous fatalities. In the current study, a small subset of individuals showed lapses of varying amounts of time on either the probe recognition or visual monitoring sub-tasks, but their performance on the other concurrent sub-tasks suggested they were providing effort and simply failed to adequately monitor all four sub-tasks in that moment. Notably, exploratory analyses indicated that performance on the three memory span tasks did not reliably distinguish between individuals with and without these multitasking lapses. Future work could be directed at understanding who is prone to such multitasking failures, and what could be done to minimize such occurrences, particularly in multitasking contexts where such errors would be costly. This approach might dovetail with research investigating a small subset of individuals labeled as ‘supertaskers’, who show no multitasking deficits relative to single-task situations (Watson & Strayer 2010; Medeiros-Ward, Watson, & Strayer, 2015).

Conclusion

The current work replicates previous research showing a robust relationship between working memory and multitasking, independent of the exact method of the task used to assess working memory. The results indicate that the shared cognitive processes across the memory

span tasks, including active maintenance in primary memory and retrieval from secondary memory, are important for understanding why working memory and synthetic work are related.

Acknowledgments

While collecting the data described in this article, TSR was supported by the Office of Naval Research (Award #N000140910129). While writing this article, TSR was supported by the National Institutes of Health (Award # 2R01AA013650-11A1).

I thank Jenny Armand, Rodney Burton, Marlyssa Fillmore, Melanie Jackson, Shonna Lane, Pamela McCoy, Jonah Robertson, and Angela Seaver for their assistance in data collection and analyses.

References

- Ackerman, P. L., & Beier, M. E. (2007). Further explorations of perceptual speed abilities in the context of assessment methods, cognitive abilities, and individual differences during skill acquisition. *Journal of Experimental Psychology: Applied*, 13, 249-272.
doi:10.1037/1076-898X.13.4.249
- Bratfisch, O., & Hagman, E. (2003). *Simultankapazität/Multi-Tasking (SIMKAP) Version 24.00: Handanweisung* [Simultaneous capacity/Multi-tasking (SIMKAP) release 24.00: Manual]. Mödling, Austria: Schuhfried
- Broadway, J. M., & Engle, R. W. (2010). Validating running memory span: Measurement of working memory capacity and links with fluid intelligence. *Behavior Research Methods*, 42, 563-570. doi:10.3758/BRM.42.2.563
- Bühner, M., König, C., Pick, M., & Krumm, S. (2006). Working memory dimensions as differential predictors of the speed and error aspect of multitasking performance. *Human Performance*, 19, 253-275. doi:10.1207/s15327043hup1903_4
- Colom, R., Rebollo, I., Abad, F. J., & Shih, P. (2006). Complex span tasks, simple span tasks, and cognitive abilities: A reanalysis of key studies. *Memory & Cognition*, 34, 158-171.
doi:10.3758/BF03193395
- Colom, R., Martínez-Molina, A., Shih, P., & Santacreu, J. (2010). Intelligence, working memory, and multitasking performance. *Intelligence*, 38, 543-551. doi:10.1016/j.intell.2010.08.002
- Comstock, J. R., & Arnegard, R. J. (1992). *The multi-attribute task battery for human operator workload and strategic behaviour research* (Tech. Memorandum No. 104174). Hampton, VA: NASA Langley Research Center.

- Conway, A. R. A., Kane, M. J., Bunting, M. F., Hambrick, D. Z., Wilhelm, O., & Engle, R. W. (2005). Working memory span tasks: A methodological review and user's guide. *Psychonomic Bulletin & Review*, 12(5), 769-786. doi:10.3758/BF03196772
- Cowan, N. (2001). The magical number 4 in short-term memory: A reconsideration of mental storage capacity. *Behavioral and Brain Sciences*, 24, 97-185.
- Cowan, N., Elliott, E. M., Sauls, J., Morey, C. C., Mattox, S., Hismjatullina, A., & Conway, A. R. A. (2005). On the capacity of attention: Its estimation and its role in working memory and cognitive aptitudes. *Cognitive Psychology*, 51, 42-100. doi:10.1016/j.cogpsych.2004.12.001
- Daneman, M., & Carpenter, P. A. (1980). Individual differences in working memory and reading. *Journal of Verbal Learning & Verbal Behavior*, 19(4), 450-466. doi:10.1016/S0022-5371(80)90312-6
- Elsmore, T. F. (1994). SYNWORK: A PC-based tool for assessment of performance in a simulated work environment. *Behavior Research Methods, Instrumentation, and Computers*, 26, 421-426. doi:10.3758/BF03204659
- Engle, R. W., Tuholski, S. W., Laughlin, J. E., & Conway, A. R. A. (1999). Working memory, short-term memory and general fluid intelligence: A latent-variable approach. *Journal of Experimental Psychology: General*, 128, 309-331. doi:10.1037/0096-3445.128.3.309
- Foster, J. L., Shipstead, Z., Harrison, T. L., Hicks, K. L., Redick, T. S., & Engle, R. W. (2015). Shortened complex span tasks can reliably measure working memory capacity. *Memory & Cognition*, 43, 226-236. doi:10.3758/s13421-014-0461-7

- Hambrick, D. Z., Oswald, F. L., Darowski, E. S., Rench, T. A., & Brou, R. (2010). Predictors of multitasking performance in a synthetic work paradigm. *Applied Cognitive Psychology*, 24, 1149-1167. doi:10.1002/acp.1624
- Hambrick, D. Z., Rench, T. A., Poposki, E. M., Darowski, E. S., Roland, D., Bearden, R. M., ... Brou, R. (2011). The relationship between the ASVAB and multitasking in Navy sailors: A process-specific approach. *Military Psychology*, 23, 365-380. doi: 10.1037/h0094762
- Jaeggi, S. M., Buschkuhl, M., Perrig, W. J., & Meier, B. (2010). The concurrent validity of the n-back task as a working memory measure. *Memory*, 18, 394-412. doi:10.1080/09658211003702171
- Kane, M. J., Conway, A. R. A., Miura, T. K., & Colflesh, J. H. (2007). Working memory, attention control, and the n-back task: A question of construct validity. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 33, 615-632. doi:10.1037/0278-7393.33.3.615
- König, C. J., Bühner, M., & Mürling, G. (2005). Working memory, fluid intelligence, and attention are predictors of multitasking performance, but polychronicity and extraversion are not. *Human Performance*, 18, 243-266. doi: 10.1207/s15327043hup1803_3
- Luck, S. J., & Vogel, E. K. (2013). Visual working memory capacity: From psychophysics and neurobiology to individual differences. *Trends in Cognitive Sciences*, 17, 391-400. doi: 10.1016/j.tics.2013.06.006
- McVay, J. C., & Kane, M. J. (2012). Why does working memory capacity predict variation in reading comprehension? On the influence of mind wandering and executive attention. *Journal of Experimental Psychology: General*, 141, 302-320. doi:10.1037/a0025250

- Medeiros-Ward, N., Watson, J. M., & Strayer, D. L. (2015). On supertaskers and the neural basis of efficient multitasking. *Psychonomic Bulletin & Review*, 22, 876-883.
doi:10.3758/s13423-014-0713-3
- Minear, M., Brasher, F., McCurdy, M., Lewis, J., & Younggren, A. (2013). Working memory, fluid intelligence, and impulsiveness in heavy media multitaskers. *Psychonomic Bulletin & Review*, 20, 1274-1281. doi:10.3758/s13423-013-0456-6
- Morgan, B., D'Mello, S., Abbott, R., Radvansky, G., Haass, M., & Tamplin, A. (2013). Individual differences in multitasking ability and adaptability. *Human Factors*, 55, 776-788. doi: 10.1177/1071181311551191
- Ophir, E., Nass, C., & Wagner, A. D. (2009). Cognitive control in media multitaskers. *PNAS Proceedings of the National Academy of Sciences of the United States of America*, 106, 15583-15587. doi:10.1073/pnas.0903620106
- Oswald, F. L., Hambrick, D. Z., & Jones, L. A. (2007). Keeping all the plates spinning: Understanding and predicting multitasking performance. In D.H. Jonassen (Ed.), *Learning to solve complex scientific problems* (pp. 77-97). Mahwah, NJ: Erlbaum.
- Oswald, F. O., McAbee, S. T., Redick, T. S., & Hambrick, D. Z. (2015). The development of a short domain-general measure of working memory capacity. *Behavior Research Methods*, 47, 1343-1355. doi:10.3758/s13428-014-0543-2
- Proctor, R. W., Wang, D. Y., & Pick, D. F. (1998). An empirical evaluation of the SYNWORK1 multiple-task work environment. *Behavior Research Methods, Instruments & Computers*, 30, 287-305. doi: 10.3758/BF03200657
- Redick, T. S., Broadway, J. M., Meier, M. E., Kuriakose, P. S., Unsworth, N., Kane, M. J., & Engle, R. W. (2012). Measuring working memory capacity with automated complex span

- tasks. *European Journal of Psychological Assessment*, 28, 164-171. doi:10.1027/1015-5759/a000123
- Redick, T. S., & Lindsey, D. R. B. (2013). Complex span and n-back measures of working memory: A meta-analysis. *Psychonomic Bulletin & Review*, 20, 1102-1113. doi:10.3758/s13423-013-0453-9
- Redick, T. S., Shipstead, Z., Harrison, T. L., Hicks, K. L., Fried, D. E., Hambrick, D. Z., Kane, M. J., & Engle, R. W. (2013). No evidence of intelligence improvement after working memory training: A randomized, placebo-controlled study. *Journal of Experimental Psychology: General*, 142, 359-379. doi:10.1037/a0029082
- Salthouse, T. A., Hambrick, D. Z., Lukas, K. E., & Dell, T. C. (1996). Determinants of adult age differences on synthetic work performance. *Journal of Experimental Psychology: Applied*, 2, 305-329. doi:10.1037/1076-898X.2.4.305
- Salthouse, T. A., & Pink, J. E. (2008). Why is working memory related to fluid intelligence?. *Psychonomic Bulletin & Review*, 15, 364-371. doi:10.3758/PBR.15.2.364
- Sanbonmatsu, D. M., Strayer, D. L., Medeiros-Ward, N., & Watson, J. M. (2013). Who multi-tasks and why? Multi-tasking ability, perceived multi-tasking ability, impulsivity, and sensation seeking. *PLoS ONE* 8(1): e54402. doi:10.1371/journal.pone.0054402
- Schmiedek, F., Hildebrandt, A., Lövdén, M., Wilhelm, O., & Lindenberger, U. (2009). Complex span versus updating tasks of working memory: The gap is not that deep. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, 35, 1089-1096. doi:10.1037/a0015730

- Shipstead, Z., Lindsey, D. B., Marshall, R. L., & Engle, R. W. (2014). The mechanisms of working memory capacity: Primary memory, secondary memory, and attention control. *Journal of Memory and Language*, 72, 116-141. doi:10.1016/j.jml.2014.01.004
- St. Clair-Thompson, H. (2012). Ascending versus randomised list lengths in working memory span tasks. *Journal of Cognitive Psychology*, 24, 335-341. doi: 10.1080/20445911.2011.639760
- Tonidandel, S., & LeBreton, J. M. (2015). RWA Web: A free, comprehensive, web-based, and user-friendly tool for relative weight analyses. *Journal of Business and Psychology*, 30(2), 207-216. doi:10.1007/s10869-014-9351-z
- Unsworth, N., & Engle, R. W. (2006). Simple and complex memory spans and their relation to fluid abilities: Evidence from list-length effects. *Journal of Memory and Language*, 54, 68-80. doi:10.1016/j.jml.2005.06.003
- Unsworth, N., & Engle, R. W. (2007). On the division of short-term and working memory: An examination of simple and complex span and their relation to higher order abilities. *Psychological Bulletin*, 133, 1038-1066. doi:10.1037/0033-2909.133.6.1038
- Unsworth, N., Fukuda, K., Awh, E., & Vogel, E. K. (2014). Working memory and fluid intelligence: Capacity, attention control, and secondary memory retrieval. *Cognitive Psychology*, 71, 1-26. doi: 10.1016/j.cogpsych.2014.01.003
- Unsworth, N., Redick, T. S., Heitz, R. P., Broadway, J. M., & Engle, R. W. (2009). Complex working memory span tasks and higher-order cognition: A latent-variable analysis of the relationship between processing and storage. *Memory*, 17, 635-654. doi:10.1080/09658210902998047

Watson, J. M., & Strayer, D. L. (2010). Supertaskers: Profiles in extraordinary multitasking ability. *Psychonomic Bulletin & Review*, 17, 479-485. doi:10.3758/PBR.17.4.479

Footnotes

¹Redick et al. (2013) reported complete pretest data from $N = 73$ participants, but 1 participant was removed from the current analyses because of a SynWin score of -2040 ($z = -6.48$; all other z -scores between -2.01 and 1.57).

²Including these additional participants does not fundamentally change the results reported here. The psychometric properties for all dependent variables were improved by excluding these participants, in particular the two participants with extreme negative SynWin scores. In addition, the other major difference between the inclusion and exclusion of these participants is that the magnitude of the zero-order correlations between the memory span tasks and SynWin increased when excluding these participants, although the correlations among the memory span tasks were relatively unchanged.

Table 1
Descriptive Statistics and Reliability Estimates for All Measures

Measure	Mean	SD	Range	Skew	Kurtosis	Reliab
Letter Span	51.73	9.73	26-71	0.01	-0.13	.69 ^a
Running Span	34.11	8.92	16-48	-0.37	-0.84	.82 ^a
Reading Span	49.25	12.72	23-71	-0.29	-0.99	.83 ^a
Reading Span Errors	5.16	4.43	0-23	1.89	5.08	.77 ^b
Synwin Overall	519.36	164.75	90-883.5	-0.24	0.07	.73 ^c
Synwin Probe Recog.	165.27	100.73	-100-290	-0.67	-0.59	.68 ^c
Synwin Arithmetic	263.00	110.24	95-585	0.60	0.01	.80 ^c
Synwin V. Monitoring	59.64	22.02	-59-74.5	-3.53	15.55	-.03 ^c
Synwin A. Monitoring	31.45	29.78	-75-55	-2.28	4.95	.66 ^c

Note. ^a For $n = 1$ to x presentations of each list length, reliability calculated by combining the first presentation of each list length into a single score, and the second presentation into a single score, through the x presentation of each list length into a single score, and computing Cronbach's alpha across the x scores. ^b Reliability calculated by computing Cronbach's alpha across all 75 sentence trials. ^c Reliability calculated by correlating scores on the first block with the second block of the task. Probe Recog: Probe Recognition; V. Monitoring: Visual Monitoring; A. Monitoring: Auditory Monitoring.

Table 2
Correlation Matrix for All Measures (N = 55)

Variable	1	2	3	4	5	6	7	8
1. Letter Span	-							
2. Running Span	.71*	-						
3. Reading Span	.44*	.42*	-					
4. Synwin Overall	.48*	.49*	.52*	-				
5. Synwin Probe Recog.	.52*	.46*	.32*	.69*	-			
6. Synwin Arithmetic	.21	.25	.39*	.67*	-.04	-		
7. Synwin V. Monitoring	-.22	.13	-.19	.01	-.01	-.15	-	
8. Synwin A. Monitoring	.28*	.31*	.48*	.71*	.58*	.28*	-.08	-

Note. $N = 55$. * $p < .05$. All correlations $r \geq .27$ and $\leq -.27$ are significant. Probe Recog: Probe Recognition; V. Monitoring: Visual Monitoring; A. Monitoring: Auditory Monitoring.

Table 3

List Length Analyses for Span Tasks in Relation to SynWin Multitasking Performance

Variable	Mean	SD	SynWin r	p
Letter Span 3	1.00	.02	.11	.41
Letter Span 4	.96	.09	.17	.23
Letter Span 5	.86	.21	.22	.10
Letter Span 6	.77	.20	.31	.02
Letter Span 7	.57	.22	.42	<.01
Letter Span 8	.43	.21	.40	<.01
Letter Span 9	.30	.16	.23	.09
Running Span 4	.89	.16	.33	.01
Running Span 5	.78	.22	.47	<.01
Running Span 6	.61	.27	.33	.01
Running Span 7	.56	.28	.42	<.01
Reading Span 3	.85	.16	.22	.12
Reading Span 4	.79	.21	.34	.01
Reading Span 5	.67	.26	.47	<.01
Reading Span 6	.62	.25	.36	<.01
Reading Span 7	.52	.21	.45	<.01

Note. In the Variable column, the number indicates the list length on that particular task.

Figure 1. Example screenshot from SynWin.