The common genetic variants in *IL-1B* and *IL-1RN* may have no predisposition to alcoholic liver disease: a meta-analysis

Running title: *IL-1B* and *IL-1RN* polymorphisms and alcoholic liver disease

Mo Xiao, Rui Yang, Yu-Ying Zhao, Tao Zeng*

Institute of Toxicology, School of Public Health, Shandong University, Jinan, Shandong Province, China.

Correspondence should be addressed to Tao Zeng, Associate professor in Institute of Toxicology, School of Public Health, Shandong University, 44 Wenhua West Road, Shandong Province, Jinan City, 250012, China; Tel.: +86-531-8838-2132; Fax: +86-531-8838-2553; E-mail addresses: zengtao@sdu.edu.cn

List of abbreviations

ALD, alcoholic liver disease; AWLD, alcoholics without liver disease; ALC, alcoholic liver cirrhosis; CD14, cluster of differentiation 14; CIs, confidence intervals; HCs, healthy controls; IL-1 β , Interleukin-1 β ; IL1-Ra, interleukin-1 receptor antagonist; KCs, Kupffer cells; LPS, lipopolysaccharide; MD-2, myeloid differentiation factor 2; ORs, odds ratios; PCR-RELP, polymerase chain reaction-restriction fragment length polymorphism; ROS, reactive oxygen species; TLR-4, toll-like receptor 4; TNF- α , tumor necrosis factor α ; VNTR, variable number tandem repeat;

Abstract

Background: Interleukin-1 β (IL-1 β) has been demonstrated to play critical roles in the development of alcoholic liver disease (ALD). However, the association studies about functional polymorphisms of *interleukin-1 β* gene (*IL-1B*) and *interleukin-1 receptor antagonist* gene (*IL-1RN*) and individual susceptibility to ALD produced conflicting results. Thus, a systematic review and meta-analysis was performed to investigate the associations between *IL-1B* and *IL1RN* polymorphisms and ALD risks.

Methods: Relevant studies were retrieved from six database including PubMed, Web of Science, Scopus, China Biology Medical literature database (CBM), Database of Chinese Scientific and Technical Periodicals (VIP), and Wanfang Data. Pooled odds ratio (OR) and 95% confidence interval (95%CI) were calculated by random-effects model.

Results: A total of 10 studies (including 825 ALD patients, 470 alcoholics without ALD and 743 health controls) were identified and included in the meta-analysis. The combined results showed that there were no significant associations between *IL-1B* (-511C>T and +3953T>C) polymorphisms and ALD susceptibility, or between *IL-1RN VNTR* polymorphism and ALD susceptibility. Moderate to higher heterogeneity was detected in most comparisons. No obvious publication bias was observed in all the genetic models.

Conclusion: The common genetic variants in *IL-1B* and *IL-1RN* may have no predisposition to ALD. Large and well-designed epidemiological studies are needed to confirm these results.

Keywords: Interleukin-1 β ; Interleukin-1 receptor antagonist; Alcoholic liver disease; Kupffer cells; Meta-analysis

1. Introduction

Alcoholic liver disease (ALD) represents a progressively aggravated liver disease that range from fatty liver to hepatic inflammation, fibrosis, and cirrhosis (Gramenzi et al., 2006). Although the majority of long-term heavy drinkers develop fatty liver, only 10%-35% of them develop hepatitis and 8%-20% of them will progress to cirrhosis (O'Shea et al., 2010). Accumulating evidences indicate that genetic factors may play essential roles in the pathogenesis and progression of ALD (Zeng et al., 2013). In this regards, the common genetic variants in endotoxin/lipopolysaccharide (LPS) signaling pathway have gained great interest due to the important roles of endotoxin/LPS and Kupffer cells (KCs) in the pathogenesis of ALD (Gao, 2012).

Innate immunity response to ethanol has been demonstrated to play predominant roles in the pathogenesis of ALD (Gao, 2012; Zeng et al., 2016). Excessive alcohol consumption enhances the gut permeability, resulting in increased translocation of gut-sourced endotoxin/LPS to liver (Rao, 1998; Basuroy et al., 2005; Zeng et al., 2016). LPS combines with Toll-like receptor 4 (TLR-4), cluster of differentiation 14 (CD14), and myeloid differentiation factor 2 (MD-2), leading to the activation of KCs (Szabo and Bala, 2010; Zeng et al., 2016). The activated KCs can release a large amount of reactive oxygen species (ROS) and pro-inflammatory cytokines including interleukin 1 β (IL-1 β), which has been demonstrated to play critical roles in the pathogenesis of ALD (Ceni et al., 2014). The serum levels of IL-1 β in ALD patients and chronic ethanol-intoxicated rats were significantly increased (Tilg et al., 1992; Valles et al., 2003). Importantly, physiological doses of IL-1 β induced steatosis, inflammatory and pro-steatotic chemokine MCP-1 in hepatocytes, while recombinant interleukin-1 receptor antagonist (IL-1Ra) blocked IL-1 signaling and markedly attenuated ALD (Petrasek et al., 2012). Results of these studies suggest that IL-1 β

plays critical roles in the pathogenesis of ALD and inhibition of IL-1 signaling may be beneficial for various stage of ALD, including fatty liver, steatohepatitis, and fibrosis (Mathews and Gao, 2013).

IL-1 β and IL-1Ra are encoded by *IL-1B* and *IL-1RN* genes, respectively, which share the same region of chromosome 2q12-21(Mao et al., 2000). Several functionally relevant genetic variants in these genes have been identified. Three polymorphic loci in *IL-1B* gene, i.e. -511 C>T in the promoter, +3953 T>C in exon 5 and -31 C>T in the promoter, have been reported to regulate the secretion of IL-1 β (Pociot et al., 1992; Santtila et al., 1998). The *IL-1RN* gene contains an 86bp variable number tandem repeat (VNTR) polymorphism in intron 2. As previous epidemiological studies about the associations between these polymorphisms and ALD risks provided conflicting results, a systematic review and meta-analysis was performed.

2. Methods

2.1. Literature and search strategy

Relevant studies were retrieved from six databases including PubMed, ISI Web of Science, Scopus, China Biology Medical literature database (CBM), Database of Chinese Scientific and Technical Periodicals (VIP), and Wanfang Data. The following keywords were jointly used to identify the potentially relevant studies: “gene polymorphism or variation or mutation”, “alcoholic liver disease or alcoholic liver cirrhosis or alcoholic hepatitis or alcoholic liver fibrosis” and “cytokine or interleukin-1 β or IL-1 β , or interleukin-1 receptor antagonist, or IL-1Ra”. The reference lists and cited literatures of selected studies were manually examined for other potentially eligible studies. There was no restriction on time period, population, sample size, or language. The literature search was updated to 20 July 2015.

2.2. Inclusion and exclusion criteria

Studies **included** in this meta-analysis must meet the following criteria: (1) case-control design; (2) definite diagnose of ALD; (3) evaluating the associations between ALD susceptibility and *IL-1B* polymorphisms (-511C>T, and/or +3953T>C, and/or -31T>C), or between ALD susceptibility and *IL-1RN VNTR* polymorphism; (4) numbers of individuals with each genotype were reported; (5) if more than on article was published using the same case series, only the study with largest sample size was selected.

2.3. Data extraction

The following information was extracted from each study: (1) name of the first author; (2) publication year; (3) country of origin; (4) age and sex of enrolled participants; (5) genotyping method; (6) genotype and allele distribution; and (7) detailed definition of case and control in each study.

2.4. Assessment of the methodological quality of included studies

The methodological quality of included studies was assessed using the Newcastle-Ottawa Scale (NOS) as previously reported (Thelma Beatriz et al., 2014; Wu et al., 2015). The quality of each study was evaluated based on 8 items that assess patient selection, study comparability and exposure(Wells GA et al.). A score of 6 will be considered as the cut-off point to distinguish higher quality studies from lower ones (Thelma Beatriz et al., 2014).

2.5. Statistical analysis

Deviation from Hardy-Weinberg equilibrium (HWE) for genotype distribution in the control group was analyzed using χ^2 test. The associations between *IL-1B*, *IL-1RN* polymorphisms and ALD susceptibility were evaluated by the crude odds ratios (ORs) with 95% confidence intervals (CIs). The comparisons were made between ALD patients and alcoholics without liver disease (AWLD), and also between ALD patients

and healthy controls (HCs). Meta-analyses were not performed when there were less than 3 studies available. The pooled ORs were calculated using a random-effect model, and the statistical significance was determined by the Z-test. The statistical heterogeneity among studies was estimated by Cochran's Q test and the I^2 statistic. Sensitivity analysis was performed to assess the effects of individual study on pooled results by removing one single study once a time to check whether the pooled ORs were stable (Tobias, 1999). The publication bias was detected with Begg's funnel plot and Egger's linear regression method. Probability less than 0.05 was judged to be significant. All statistical analyses were carried out using Stata software (version 11.0; StataCorp 2P, College Station, Texas, USA).

3. Result

3.1. Characteristics of studies included in the meta-analysis

The flow chart of literature search was shown in Figure 1. A total of 489 potentially relevant articles were identified in the literature search, and 463 irrelevant papers were excluded by reading the titles and abstracts. Another 16 studies were excluded as they were review articles, not about ALD, etc. Finally, 10 eligible studies (8 about the **associations** between *IL-1B* polymorphisms and ALD susceptibility, 4 about *IL-1RN VNTR* polymorphism and ALD susceptibility) were included in the meta-analyses (Pastor et al., 2000; Takamatsu et al., 2000; Kim et al., 2004; Chen et al., 2005; Pastor et al., 2005; Gleeson et al., 2008; Petrasek et al., 2009; Roy et al., 2012; Dutta, 2013; Yang et al., 2014). For the studies about *IL-1B* polymorphisms, 6 studies investigated the associations between -511 C>T polymorphism and ALD susceptibility, while 3 **studies** and 1 **study** were between +3953 T>C polymorphism and ALD susceptibility and between -31 T>C polymorphism and ALD susceptibility, respectively. The genotype distributions in controls of all studies were consistent with

HWE. All ALD patients had definite diagnose and excluded other causes of liver disease. Most of included studies used peripheral blood for DNA extraction and polymerase chain reaction-restriction fragment length polymorphism (PCR-RFLP) methods for genotyping. The characteristics of selected studies were summarized in Table 1 and Table 2. The detailed criteria for the selection of cases and controls in the included studies were provided in supplementary Table 1s.

3.2. Quantitative data synthesis

Results of meta-analysis for the associations between *IL-1B* polymorphism and ALD susceptibility were shown in Table 3 and Figure 2. Overall, no significant association between *IL-1B* -511C>T polymorphism and ALD susceptibility was detected in the comparison between ALD patients and HCs (T vs. C, OR =0.980, 95%CI 0.719-1.335; TT vs. CC, OR=0.892, 95%CI 0.471-1.690), or in the comparison between ALD patients and alcoholics without liver disease (AWLD) (T vs. C, OR =1.126, 95%CI 0.815-1.556; TT vs. CC, OR=1.233, 95%CI 0.398-2.544). As ALD contains many types of histological forms, we further extracted clearly defined alcoholic liver cirrhosis (ALC) as subgroup to investigate whether its susceptibility was correlated with *IL-1B* -511C>T polymorphism. However, no significant association was detected either compared with HCs or with ALWD (supplementary Table 2s). For the associations between *IL-1B* +3953 T>C polymorphism and ALD susceptibility, combined results of 3 studies showed no significant association in the comparisons between ALD patients and AWLD (C vs. T: OR=1.086, 95%CI 0.564-2.092; CC vs. CT+TT: OR=1.223, 95%CI 0.600-2.494). In regard to the association between *IL-1RN* VNTR polymorphism and ALD susceptibility, no significant association was detected in the comparisons between ALD patients and HCs (allele 2 containing genotypes vs. other genotypes: OR=0.769, 95%CI

0.291-2.028; allele 2 vs. other alleles: OR=0.798, 95%CI 0.344-1.847). (Table 3)

3.3. Heterogeneity and sensitivity analysis

Moderate to higher heterogeneity was detected in most comparisons ($I^2 > 50\%$ and $P_{Qtest} < 0.10$) (Table 3). The heterogeneity in the subgroup analyses of ALC risks and *IL-1B -511 C>T* polymorphism was not significantly altered, which suggested that the heterogeneity might not be related with the histological types of ALD. A single study was removed from meta-analysis each time to determine the influence of its individual data sets to the pooled ORs, and the corresponding pooled ORs were not materially altered, which indicated that the results were relatively credible and stable.

3.4. Publication bias

Begg's funnel plot and Egger's test revealed there was no obvious publication bias. These results were also demonstrated by the shape of funnel plot (supplementary Fig.1s-Fig.4s).

Discussion

In the current meta-analysis, we combined 10 eligible studies with 825 cases, 470 alcoholics control and 743 health controls, to evaluate the associations between *IL-1B* and *IL-1RN VNTR* polymorphisms and ALD susceptibility. To the best of our knowledge, this is the first meta-analysis focused on the relationships between the *IL-1B* and *IL-1RN VNTR* polymorphisms and ALD susceptibility. The combined results showed no significant association between the *IL-1B -511C>T* polymorphism and ALD susceptibility in the allele model, recessive model, dominant model, or co-dominant model. Furthermore, the subgroup analyses in ALC patients also revealed no significant association. Hence, the *IL-1B -511C>T* polymorphism might be not the major risk factor in the individual susceptibility to ALD. Similarly, combined results also revealed that the *IL-1B +3953T>C* polymorphism and the

IL-1RN VNTR polymorphism may be not significantly associated with individual risks to ALD. However, as only 3 studies were included in the meta-analyses for the *IL-1B* +3953T>C polymorphism and *IL1RN VNTR* polymorphism, much more works are needed to confirm these results.

It should be emphasized that moderate to higher between-study heterogeneity was detected in this meta-analysis. ALD is a multifactorial disease affected by many factors including the types of alcohol, drinking habits/patterns, age, gender, obesity, iron overload, ethnicity, and genetic factors of enrolled subjects (Dohmen et al., 1996; Bellentani et al., 1997; Cederbaum, 2003; Douds et al., 2003; Stickel and Österreicher, 2006). In addition, ALD is a progressively aggravated liver disease with many pathological manifestations from simple steatosis to irreversible cirrhosis. In the current meta-analysis, although all ALD patients in included studies had definite diagnose and had similar **quantity** and **frequency** of alcohol consumption; however, other factors such as the ethnicity and the disease stage were not all consistent, which may explain the heterogeneity detected in the meta-analysis. The subgroup analyses in ALC patients showed that the heterogeneity might not be related with the histological types of ALD; however, subgroup analyses based on the gender, age, and ethnicity could not be performed due to the limited studies included in the meta-analysis. Therefore, although the current meta-analysis showed no significant associations between *IL-1B* -511C>T polymorphism and ALD susceptibility, it could not completely preclude its roles in individual susceptibility to ALD.

The previous study by Petrasek *et al.* provides convincing evidence that IL-1 signaling plays an important role in ethanol-induced hepatic steatosis, steatohepatitis, and fibrosis, and supports the potential roles of IL-1Ra in the treatment of ALD (Petrasek et al., 2012; Mathews and Gao, 2013). If the results of the current

meta-analysis are correct, the null association between *IL-1B* -511C>T polymorphism and ALD susceptibility may be related with several reasons. Firstly, the *IL-1B* -511C>T polymorphism may be not significantly associated with the production of IL-1 β in alcoholics. Although the mutant allele of *IL-1B* -511C>T polymorphism was associated with increased IL-1 β production *in vitro*, and serum IL-1 β levels were significantly increased in chronic ethanol-exposed mice compared with control mice; however, the association between *IL-1B* -511C>T polymorphism and the serum IL-1 β levels in alcoholics has not been proved. It is possible that *IL-1B* -511C>T polymorphism may not significantly affect the IL-1 β production in chronic alcoholics. Secondly, the study by Petrasek *et al.* demonstrated that IL-1 signaling were required for the pathogenesis of alcohol-induced inflammation, steatosis, liver damage, and fibrosis (Petrasek et al., 2012). However, most studies included in this meta-analysis enrolled ALC patients. Thus, it is interesting to further investigate the association between *L-1B* -511C>T polymorphism and the earlier phenotypes of ALD (steatosis, hepatitis, and fibrosis).

In summary, this meta-analysis did not detect significant associations between *IL-1B* (-511C>T and +3953C>T) and *IL1RN VNTR* polymorphisms and individual susceptibility to ALD. Considering the limited numbers of included studies and potential confounding factors in this study, large and well-designed epidemiological studies are needed to confirm these results.

Conflict of interest

The authors declare that they have no competing interests exist.

Acknowledgements

This work was supported by the National Science Foundation of China (Grant No. 81473004, 81102153) and the Young Scholars Program of Shandong University

(Grant No. 2015WLJH52).

References

- Basuroy, S., Sheth, P., Mansbach, C.M. and Rao, R.K., 2005. Acetaldehyde disrupts tight junctions and adherens junctions in human colonic mucosa: protection by EGF and L-glutamine. *Am J Physiol Gastrointest Liver Physiol* 289, G367-75.
- Bellentani, S., Saccoccio, G., Costa, G., Tiribelli, C., Manenti, F., Sodde, M., Saveria Croce, L., Sasso, F., Pozzato, G., Cristianini, G. and Brandi, G., 1997. Drinking habits as cofactors of risk for alcohol induced liver damage. The Dionysos Study Group. *Gut* 41, 845-50.
- Cederbaum, A.I., 2003. Iron and CYP2E1-dependent oxidative stress and toxicity. *Alcohol* 30, 115-20.
- Ceni, E., Mello, T. and Galli, A., 2014. Pathogenesis of alcoholic liver disease: role of oxidative metabolism. *World J Gastroenterol* 20, 17756-72.
- Chen, W.-X., Xu, G.-Y., Yu, C.-H., Wang, P., Yu, M., Xu, P., Liu, Y. and Li, Y., 2005. Correlation of polymorphism in the interleukin-1 receptor antagonist gene intron 2 with alcoholic liver disease. *Hepatobiliary Pancreat Dis Int* 4, 41-45.
- Dohmen, K., Baraona, E., Ishibashi, H., Pozzato, G., Moretti, M., Matsunaga, C., Fujimoto, K. and Lieber, C.S., 1996. Ethnic differences in gastric sigma-alcohol dehydrogenase activity and ethanol first-pass metabolism. *Alcohol Clin Exp Res* 20, 1569-76.
- Douds, A.C., Cox, M.A., Iqbal, T.H. and Cooper, B.T., 2003. Ethnic differences in cirrhosis of the liver in a British city: alcoholic cirrhosis in South Asian men.

- Alcohol Alcohol 38, 148-50.
- Dutta, A.K., 2013. Genetic factors affecting susceptibility to alcoholic liver disease in an Indian population. *Ann Hepatol* 12, 901-7.
- Gao, B., 2012. Hepatoprotective and anti-inflammatory cytokines in alcoholic liver disease. *J Gastroenterol Hepatol* 27 Suppl 2, 89-93.
- Gleeson, D., Bradley, M.P., Jones, J., Peck, R.J., Bond, S.K., Teare, M.D. and Duff, G.W., 2008. Cytokine gene polymorphisms in heavy drinkers with and without decompensated liver disease: a case-control study. *Am J Gastroenterol* 103, 3039-46.
- Gramenzi, A., Caputo, F., Biselli, M., Kuria, F., Loggi, E., Andreone, P. and Bernardi, M., 2006. Review article: alcoholic liver disease--pathophysiological aspects and risk factors. *Aliment Pharmacol Ther* 24, 1151-61.
- Kim, M.S., Lee, D.H., Kang, H.S., Park, H.S., Jung, S., Lee, J.W., Kwon, K.S., Kim, P.S., Kim, H.G., Shin, Y.W., Kim, Y.S., Baek, I. and Lee, M.S., 2004. [Genetic polymorphisms of alcohol-metabolizing enzymes and cytokines in patients with alcohol induced pancreatitis and alcoholic liver cirrhosis]. *Korean J Gastroenterol* 43, 355-63.
- Mao, X.Q., Kawai, M., Yamashita, T., Enomoto, T., Dake, Y., Sasaki, S., Kataoka, Y., Fukuzumi, T., Endo, K., Sano, H., Aoki, T., Kurimoto, F., Adra, C.N., Shirakawa, T. and Hopkin, J.M., 2000. Imbalance production between interleukin-1beta (IL-1beta) and IL-1 receptor antagonist (IL-1Ra) in bronchial asthma. *Biochem Biophys Res Commun* 276, 607-12.

- Mathews, S. and Gao, B., 2013. Therapeutic potential of interleukin 1 inhibitors in the treatment of alcoholic liver disease. *Hepatology* 57, 2078-80.
- O'Shea, R.S., Dasarathy, S. and McCullough, A.J., 2010. Alcoholic liver disease. *Hepatology* 51, 307-28.
- Pastor, I.J., Laso, F.J., Avila, J.J., Rodriguez, R.E. and Gonzalez-Sarmiento, R., 2000. Polymorphism in the interleukin-1 receptor antagonist gene is associated with alcoholism in Spanish men. *Alcohol Clin Exp Res* 24, 1479-82.
- Pastor, I.J., Laso, F.J., Romero, A. and Gonzalez-Sarmiento, R., 2005. Interleukin-1 gene cluster polymorphisms and alcoholism in Spanish men. *Alcohol Alcohol* 40, 181-6.
- Petrasek, J., Bala, S., Csak, T., Lippai, D., Kodys, K., Menashy, V., Barrieau, M., Min, S.Y., Kurt-Jones, E.A. and Szabo, G., 2012. IL-1 receptor antagonist ameliorates inflammasome-dependent alcoholic steatohepatitis in mice. *J Clin Invest* 122, 3476-89.
- Petrasek, J., Hubacek, J.A., Stickel, F., Sperl, J., Berg, T., Ruf, E., Wichmann, H.E., Pfeufer, A., Meitinger, T., Trunecka, P., Spicak, J. and Jirsa, M., 2009. Do common genetic variants in endotoxin signaling pathway contribute to predisposition to alcoholic liver cirrhosis? *Clin Chem Lab Med* 47, 398-404.
- Pociot, F., Molvig, J., Wogensen, L., Worsaae, H. and Nerup, J., 1992. A TaqI polymorphism in the human interleukin-1 beta (IL-1 beta) gene correlates with IL-1 beta secretion in vitro. *Eur J Clin Invest* 22, 396-402.
- Rao, R.K., 1998. Acetaldehyde-induced increase in paracellular permeability in

- Caco-2 cell monolayer. *Alcohol Clin Exp Res* 22, 1724-30.
- Roy, N., Mukhopadhyay, I., Das, K., Pandit, P., Majumder, P.P., Santra, A., Datta, S., Banerjee, S. and Chowdhury, A., 2012. Genetic variants of TNFalpha, IL10, IL1beta, CTLA4 and TGFbeta1 modulate the indices of alcohol-induced liver injury in East Indian population. *Gene* 509, 178-88.
- Santtila, S., Savinainen, K. and Hurme, M., 1998. Presence of the IL-1RA allele 2 (IL1RN*2) is associated with enhanced IL-1beta production in vitro. *Scand J Immunol* 47, 195-8.
- Stickel, F. and Österreicher, C.H., 2006. The role of genetic polymorphisms in alcoholic liver disease. *Alcohol Alcohol* 41, 209-224.
- Szabo, G. and Bala, S., 2010. Alcoholic liver disease and the gut-liver axis. *World J Gastroenterol* 16, 1321-9.
- Takamatsu, M., Yamauchi, M., Maezawa, Y., Saito, S., Maeyama, S. and Uchikoshi, T., 2000. Genetic polymorphisms of interleukin-1beta in association with the development of alcoholic liver disease in Japanese patients. *Am J Gastroenterol* 95, 1305-11.
- Thelma Beatriz, G.C., Isela, J.R., Alma, G., Maria Lilia, L.N. and Carlos Alfonso, T.Z., 2014. Association between HTR2C gene variants and suicidal behaviour: a protocol for the systematic review and meta-analysis of genetic studies. *BMJ Open* 4, e005423.
- Tilg, H., Wilmer, A., Vogel, W., Herold, M., Nolchen, B., Judmaier, G. and Huber, C., 1992. Serum levels of cytokines in chronic liver diseases. *Gastroenterology*

103, 264-74.

Tobias, A., 1999. Assessing the influence of a single study in the meta-analysis estimate. Stata Technical Bulletin 8.

Valles, S.L., Blanco, A.M., Azorin, I., Guasch, R., Pascual, M., Gomez-Lechon, M.J., Renau-Piqueras, J. and Guerri, C., 2003. Chronic ethanol consumption enhances interleukin-1-mediated signal transduction in rat liver and in cultured hepatocytes. Alcohol Clin Exp Res 27, 1979-86.

Wells GA, Shea B, O'Connell D, Peterson J, Welch V, Losos M and P, T. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses.

http://www.ohri.ca/programs/clinical_epidemiology/oxford.asp.

Wu, N., Zhang, X., Jia, P. and Jia, D., 2015. Lack of an Association between the SDF-1 rs1801157 Polymorphism and Coronary Heart Disease: A Meta-Analysis. Sci Rep 5, 11803.

Yang, A.M., Wen, L.L., Yang, C.S., Wang, S.C., Chen, C.S. and Bair, M.J., 2014. Interleukin 10 promoter haplotype is associated with alcoholic liver cirrhosis in Taiwanese patients. Kaohsiung J Med Sci 30, 291-8.

Zeng, T., Guo, F.F., Zhang, C.L., Song, F.Y., Zhao, X.L. and Xie, K.Q., 2013. Roles of cytochrome P4502E1 gene polymorphisms and the risks of alcoholic liver disease: a meta-analysis. PLoS One 8, e54188.

Zeng, T., Zhang, C.-L., Xiao, M., Yang, R. and Xie, K.-Q., 2016. Critical roles of Kupffer cells in the pathogenesis of alcoholic liver disease: from basic science

to clinical trials. *Frontiers in Immunology* 7, 538.

Figure legends

Figure 1 The flow chart of study selection based on the inclusion and exclusion criteria.

Figure 2 Forest plot for the association between *IL-1B -511C>T* polymorphism and individual susceptibility to alcoholic liver disease (TT vs. CC). 1) Patients with alcoholic liver disease (ALD) vs. healthy controls (HCs); 2) ALD patients vs. alcoholics without liver disease (AWLD). Each study was shown by a point estimate of the effect size (size inversely proportional to its variance) and its 95% confidence interval (95%CI) (horizontal lines). The white diamond denotes the pooled OR.

Table 1 Characteristics of individual studies included in the meta-analysis for the association between the *IL-1B* polymorphisms and risk of alcoholic liver disease

Study ^[Ref.]	Country	Age ^a	NOS score ^b	Type of ALD	SNP	Genotyping	HC			AWLD			ALD			<i>P</i> _{HWE} ^d
							CC	TC	TT	CC	TC	TT	CC	TC	TT	
Takamatsu, 2000 ^[36]	Japan	NA/52.4/55.0	7	AH, ALC	-511 C>T +3953 T>C	PCR-RFLP	68	109	41	16	12	2	38	86	18	>0.05
Kim, 2004 ^[35]	Korea	49.2/50.4/48.8	na ^c	ALC	-511 C>T	PCR-RFLP	201	17	0	24	6	0	133	9	0	
Pastor, 2005 ^[33]	Spain	matched	7	ALC	-511 C>T	PCR-RFLP	26	40	15	41	36	5	29	25	3	NA
					+3953 T>C		49	30	2	52	27	3	31	20	6	
Gleeson, 2008 ^[32]	UK	-/49.5/47.1	6	ALD	-511 C>T	Taqman fluorogenic 5' nuclease assay	-	-	-	71	76	14	97	97	29	>0.05
					+3953 T>C		-	-	-	96	60	6	136	70	16	
Petrasek, 2009 ^[31]	Czech Republic	50/-/50	7	ALC	-511 C>T	PCR-RFLP	75	82	23	-	-	-	41	45	14	>0.05
					-31 C >T		23	82	75	-	-	-	14	45	41	
Roy, 2012 ^[30]	India	-/41.0/43.9	7	ALC	-511 C>T	PCR-RFLP	-	-	-	36	49	23	77	83	21	>0.05
Dutta, 2013 ^[29]	India	32.9/38.0/49.5	7	ALC	-511 C>T	PCR-RFLP	12	60	28	11	27	22	6	27	27	>0.05
Yang, 2014 ^[28]	China	NA/NA/NA	4	ALC	-511 C>T	PCR-RFLP	43	17	4	-	-	-	23	12	5	>0.05

^a Mean age of each group;

^b The Newcastle-Ottawa Scale (NOS) was used to assess the quality of included case-control studies; AWLD was considered as the control group unless the study only included healthy subjects as controls.

^c The NOS of the study by Kim., 2004 can not be obtained, as the study was published in Korean.

^d *P* for Hardy–Weinberg equilibrium test in healthy controls (non-alcoholics);

HC, healthy control; AWLD, alcoholics without liver diseases; ALD, alcoholic liver disease; AH, alcoholic hepatitis; ALC, alcoholic liver cirrhosis; NA, not available; PCR-RFLP, polymerase chain reaction-restriction fragment length polymorphism.

Table 2 Characteristics of individual studies included in the meta-analysis about *IL-1RN intron2 VNTR* polymorphism

Study ^[Ref.]	Country	Age ^a	NOS score ^b	Type of ALD	Genotyping	HC					AWLD					ALD				
						genotypes ^c		alleles			genotypes ^c		alleles			genotypes ^c		alleles		
						1/1	others	1	2	other	1/1	others	1	2	others	1/1	others	1	2	others
Pastor, 2000 ^[37]	Spain	matched/50/50	7	ALC	PCR-RFLP	26	14	61	14	5	57	3	115	2	3	28	2	56	2	2
Chen, 2005 ^[34]	China	45.9/44.5/44.5	7	ALC	PCR-RFLP	39	26	100	22	8	39	4	82	4	0	20	11	49	11	2
Pastor, 2005 ^[33]	Spain	matched/50/50	7	ALC	PCR-RFLP	59	22				76	6				53	4			
Petrasek, 2009 ^[31]	Czech Republic	50/-/50	7	AIC	PCR-RFLP	82	98	249	104	7						42	58	132	67	1

^a Mean age of each group;

^b The Newcastle-Ottawa Scale (NOS) was used to assess the quality of included case-control studies; AWLD was considered as the control group unless the study only included healthy subjects as controls.

^c other genotypes including 2/2, 1/2, 1/3, 1/4, 3/3, 4/4, 2/4 were combined as the frequencies of these genotypes were very low;

HC, healthy controls; AWLD, alcoholics without liver diseases; ALD, alcoholic liver disease; AH, alcoholic hepatitis; ALC, alcoholic liver cirrhosis; NA, not available; PCR-RFLP, polymerase chain reaction-restriction fragment length polymorphism.

Table 3 Meta-analysis of the association between genetic polymorphisms and individual susceptibility to alcoholic liver disease

SNP	Comparisons	Number ^a	Test of associations				Heterogeneity		<i>P</i> _{Egger's test}
			OR	95%CI	M ^b	<i>P</i> _{OR}	<i>I</i> ² (%)	<i>P</i> _c	
<i>IL-1B-511 C>T</i>	ALD patients vs. HCs								
	T vs. C	6	0.980	0.719-1.335	R	0.896	60.6	0.027	0.904
	TT vs. CC	6	0.892	0.471-1.690	R	0.727	50.5	0.073	0.758
	TC vs. CC	6	1.038	0.785-1.373	R	0.791	0.0	0.444	0.426
	TT vs. TC+CC	6	0.929	0.491-1.757	R	0.820	63.2	0.018	0.658
	TT+TC vs. CC	6	0.981	0.703-1.369	R	0.908	30.0	0.210	0.717
	ALD patients vs. AWLD								
	T vs. C	6	1.126	0.815-1.556	R	0.471	61.5	0.024	0.297
	TT vs. CC	6	1.233	0.598-2.544	R	0.570	56.9	0.040	0.418
	TC vs. CC	6	1.181	0.802-1.738	R	0.400	42.4	0.122	0.134
	TT vs. TC+CC	6	1.069	0.634-1.084	R	0.802	39.7	0.141	0.671
	TT+TC vs. CC	6	1.220	0.789-1.885	R	0.371	57.9	0.037	0.183
<i>IL-1B+3953 T>C</i>	ALD patients vs. AWLD								
	C vs. T	3	1.086	0.564-2.092	R	0.805	72.4	0.027	0.607
	CC vs. CT+TT	3	1.223	0.600-2.494	R	0.580	68.2	0.043	0.654
<i>IL-1RN VNTR</i>	ALD patients vs. HCs								
	1/1 genotype vs. others	3	1.552	0.613-3.932	R	0.354	74.3	0.020	0.303
	Allele 2 containing genotype vs. others	3	0.769	0.291-2.028	R	0.595	65.5	0.055	0.222
	Allele 1 vs. others	3	1.382	0.640-2.988	R	0.410	72.0	0.028	0.260
	Allele 2 vs. others	3	0.798	0.344-1.847	R	0.598	69.6	0.037	0.301

CI, confidence interval; OR, odds ratio; ALD, alcoholic liver disease; HCs, healthy controls; AWLD, alcoholics without liver diseases.

^a Number of studies included in the meta-analysis;

^b Model of meta-analysis; Random-effect model (R) was used to calculate the pooled ORs;

^c *P* value for heterogeneity based on Q-test.

NOTE: Weights are from random effects analysis

NOTE: Weights are from random effects analysis