“Other” possibilities? Assessing regional and extra-regional dental affinities of populations in the Portuguese Estremadura to explore the roots of Iberia’s Late Neolithic-Copper Age

Joel D. Irish^a, Katina T. Lillios^b, Anna J. Waterman^c, and Ana M. Silva^d

Running Title: Dental affinities in the Portuguese Estremadura

Key Words: Bioarchaeology; biological affinity; nonmetric traits; social complexity; Portugal

Abstract: The relationship between the development of social complexity in the Iberian Peninsula during the 4th and 3rd millennia BCE (Late Neolithic and Copper Age) and population movement has been a longstanding question. Biological affinity analyses were used to explore Iberian demographic dynamics, and specifically, to discern whether there is evidence for migration and gene flow between northwest African, eastern Mediterranean, and Iberian populations. Affinities based on comparisons of nonmetric traits from the Arizona State University Dental Anthropology System were estimated among samples of burial populations from three key Late Neolithic-Copper Age sites in the Portuguese Estremadura: Cova da Moura (3700-2300 BCE), Bolores (2800-2600 BCE), and Pai Mogo I (2800-2600 BCE). Results indicate: 1) the possibility of genetic exchange with African and other Mediterranean peoples, 2) some measure of population continuity over time in the Estremadura, and 3) possible local isolation of populations, given distinctive patterning at the site of Pai Mogo, located 23 km north of Cova da Moura and Bolores.

1.1 Introduction

Understanding the relationship between social complexity and demographic dynamics is a central theme in several of the grand challenges recently identified for archaeology, especially those related to movement, mobility, and migration (Kintigh et al., 2014). In the Iberian Peninsula, the emergence of social complexity developed in the second half of the 4th and the 3rd millennia BCE – in the periods known as the Late Neolithic and Copper Age. During this time a number of fortified or ditched enclosure sites emerged as important social, political, and ritual centers, particularly in the fertile Guadiana and Guadalquivir river valleys of southern Iberia. Some sites extended over large areas, such as Valencina de la Concepción (Sevilla) (450 ha) (Costa et al., 2010), Marroquíes Bajos (Jaén) (113 ha) (Zafra et al., 1999), and La Pijotilla (Badajoz) (80 ha) in Spain (Hurtado, 1997), and Porto Torrão (Beja) (70 ha) (Valera and Filipe, 2004), Alcalar (20 ha) (Morán and Parreira, 2003) and Perdigões (Reguengos de Monsaraz) (>16 ha) (Valera et al., 2014) in Portugal. A diversity of tomb architecture was utilized, including megaliths, rockshelters, natural caves, hypogea, pits, and tholos (Boaventura, 2009; Boaventura et al., 2014; Silva, 2002, 2003). The manufacture of technically complex objects and personal adornments, some made from exotic raw materials such as ivory, amber, copper, gold, variscite, and rock crystal (Morgado Rodríguez et al., 2015; Müller et al., 2007; Murillo-Barroso and Martín-Torres, 2012; Murillo-Barroso et al., 2015; Odriozola et al., 2016; Schuhmacher et al., 2009; Sousa and Gonçalves, 2012; Thomas, 2011; Villalobos García and Odriozola, 2016; Valera et al., 2015), also attests to the formation of new identities and categories of social beings.

The relationship between the process of social differentiation during the 4th and 3rd millennia and population movement has been a longstanding question in Iberian archaeology.

The Orientalist explanatory model was adopted by the main Iberian archeologists of the late 19th and early 20th centuries, especially in the Spanish Southeast and Portuguese Estremadura. Since the pioneering studies of Siret (1913), and later work of Martin Almagro and A. Arribas at Los Millares, Afonso do Paço at Vila Nova de São Pedro, and E. Sangmeister and H. Schubart at Zambujal (Sangmeister and Schubart, 1981), the fortifications were associated with eastern Mediterranean colonists; the latter purportedly came to Iberia in search of metals and to establish outposts during the Iberian Copper Age (Siret, 1913). However, when archaeologists began applying radiocarbon dating in the 1970s, they increasingly recognized that material practices, such as megalith-building, predated their supposed exogenous models (Chapman, 1990). As a result, they turned to local social or economic factors to account for the changes between the 4th and 2nd millennia cal. BCE, and discredited “migrationism” and “diffusionism” as explanatory vehicles. Instead, they replaced their models with a “radical autochthonism” (Aranda Jimenez et al., 2015: 55). Demography took a back seat to economic and political dynamics (Gonçalves et al., 2013).

Through the development of analytical technologies to source raw materials and track the mobility of people, as well as new archaeological discoveries, the “Other” – both in the form of exotic materials and non-local individuals – is once again making its appearance in narratives of late prehistoric Iberian social history. For example, archaeologists had understood that ivory objects found in Copper and Bronze Age contexts pointed to long-distance exchange networks, presumably originating from sources (e.g., elephants, hippopotami) in North Africa and Egypt (Harrison and Gilman, 1977). However, recent studies by Schumacher et al. (2009) suggest the exchange was larger in scale than imagined; ivory from Asian elephants was sometimes used, as well as that from Pleistocene and African elephants. Furthermore, building on the work of Bosch Gimpera (1932) and Martínez Santa-

Ollala (1941), Manen et al. (2007) submit that African exchange extended as far back as the Early Neolithic (i.e., 6th to 5th millennia BCE). Given their geographic proximity, could genetic exchange have also occurred between the human populations of Iberia and northwest Africa, i.e., in the form of south-to-north gene flow and/or migration? And is there yet merit in the idea of eastern Mediterranean influence (Siret, 1913) that corresponds with the Iberian Late Neolithic-Copper Age?

To explore the viability of these possible scenarios based on archaeological evidence, we turned to the subfield of bioarchaeology. Specifically, we use biological affinity analyses to better understand the demographic dynamics of the time. Previous affinity studies of early Iberian populations focused on periods that, like across much of Europe, ushered in significant social and economic change, i.e., the Mesolithic-Neolithic (c. 5600 BCE) (e.g., Jackes et al., 2001) and Middle-Late Neolithic transitions (4th-3rd millennia BCE (Boaventura, 2009; Boaventura et al., 2014; Horwath et al., 2014). A primary goal of these studies was to assess whether the change was accompanied by population replacement or, minimally, discernable outside genetic input. Our research continues this diachronic emphasis.

Biological affinities based on comparisons of nonmetric traits from the Arizona State University Dental Anthropology System (Turner et al., 1991) were estimated among samples of burial populations from three key Late Neolithic-Copper Age sites in Portugal: Cova da Moura (3700-2300 BCE), Bolores (2800-2600 BCE), and Pai Mogo I (3000-2600 BCE). Dental trait data are ideal for this investigation because, like all local burial sites of this period, the remains are so fragmented and commingled that most skeletal elements can neither be measured (e.g., for craniometrics, geometric morphometric study, etc.) nor recombined into individuals. The three samples come from the Estremadura of west-central

Portugal (Fig. 1), an important region for understanding biological diversity given its many mortuary sites and favorable conditions for skeletal preservation (Boaventura et al., 2014; Silva, 2002). They, in turn, were compared to four extra-regional samples affiliated with northwest Africa and three from the eastern Mediterranean to address the two questions posed above. If gene flow and/or some degree of population movement from either region did occur – in potential correspondence with the 4th-3rd millennia increase in social differentiation – it may be evident in the Portuguese samples.

[FIGURE 1 HERE]

At a basic level, the African scenario may be tested statistically by phrasing it in the form of a null hypothesis, viz.: there is no significant difference between one or more samples from the Portuguese Estremadura and northwest Africa. By substituting the latter region, a second null hypothesis can be created to test for any Mediterranean input. For both, the mean measure of divergence distance statistic (Irish, 2010; Sjøvold, 1977) was applied to compare the dental data among samples. If one or both of these hypotheses cannot be rejected, discernable extra-regional genetic input may have occurred. Of course, such complex problems are unlikely to be resolved through simply reaching or failing to reach an arbitrary alpha level; however, by considering the weight of evidence across all inter-sample comparisons, along with the aid of supplementary quantitative and illustrative methods (below), some pertinent insights may be achieved. As usual, all results are contingent upon two assumptions (Irish 2005, 2006, 2016; Irish et al., 2014) that: 1) all samples are representative of their corresponding populations, and 2) phenetic similarity provides some indication of genetic relatedness (Larsen, 1997; Martínón-Torres et al., 2007; Rightmire, 1999; Scott et al., 1983; Scott and Turner, 1997). In sum, our goal is to complement and build upon the findings of prior archaeological and bioarchaeological researchers to promote a

better biocultural understanding of the ancient peoples of the Iberian Peninsula and, specifically, Portugal.

1.2 Materials

The three Portuguese samples are suitable for evaluating local biological/genetic variation during the Late Neolithic-Copper Age because: 1) Cova da Moura predates and temporally overlaps with Bolores, while the two sites are located just 2 km apart in the Rio Sizandro valley, 2) the Bolores and Pai Mogo I sites are largely contemporary, but the latter is located two river valleys to the north, 23 km distant near the Atlantic coast, and 3) Cova da Moura and Pai Mogo I also overlap in time and are spatially distinct. Dietary differences and differential, albeit low-level migration based on stable isotopes have also been reported (Guiry et al., 2016; Waterman et al., 2014, 2015). All samples were studied as part of a broader analysis of social change in the Sizandro/ Alcabrichel river valleys between the 4th-2nd millennia BCE, codirected by the second author and Michael Kunst (Lillios 2015, Waterman et al., 2016).

Cova da Moura is a natural cave site in the Sizandro river valley. It was discovered in 1930, with preliminary excavations undertaken in 1932 (Belo et al., 1961). Subsequent fieldwork was conducted in 1961 by Belo and Trindade (Gallay and Spindler, 1970; Spindler, 1981). Seven radiocarbon dates were obtained on the human remains, which indicate that the site was used between 3700 and 2300 BCE. Based on duplicate skeletal elements the MNI is 90 individuals, including 15 sub-adults (Silva, 2002, 2003). However, as noted, all remains are fragmented and comingled, so multiple maxillae, mandibles, and loose teeth were combined to yield “composite” individuals for analyses following standard procedure (Irish, 1998a). Thus, the Cova da Moura dental sample consists of 41 such individuals. In terms of material culture, Cova da Moura is by far the richest burial site in the region. Grave goods

include limestone and bone idols, green stone pendants, gold artifacts, engraved slate plaques, bone, ivory, and variscite rabbit figurines, and beads. As well, both pre-Beaker and Beaker ceramics were recovered (Belo et al., 1961; Gallay and Spindler, 1970; Spindler, 1981; Thomas, 2011). As such, some population, and thus sample, heterogeneity cannot be ruled out, and should be considered in the results.

Bolores is a semi-artificial cave located in the Sizandro river valley, about 15 km from the Atlantic coast. Following testing in 1986, four seasons of excavation were conducted by the second author and a team from the University of Iowa (Lillios et al., 2015). The site itself is about 3 x 5 m, subdivided by large stone blocks into three chambers. Approximately 75% of the site was excavated. The MNI is 36, which includes adults, adolescents, and children. Of these, 36 individual dentitions could be identified for analysis. Eleven AMS dates were obtained that put the use of the site between 2800-2600 BCE, with a short period of use near 1800 BCE. Material culture is relatively scarce, and consists of stone and shell beads, stone ‘idols’ of various forms, flint blades, and ceramic vessels.

Pai Mogo I (or Paimogo I) is a corbel-vaulted tomb on the Rio Grande, 1 km from the Atlantic coast. The site was first discovered in 1968, with excavations carried out in 1971. Four dates on human remains from the site indicate the site was in use from 3000 to 2600 BCE (Silva, 2002). Little information is available about the context of the human remains recovered from this burial (Gallay et al., 1973). The MNI is 413 (Silva 2002, 2003). Forty-nine composite dentitions were assembled for analysis. An extensive array of Late Neolithic and Copper Age artifacts was found, including decorated pre-Beaker and Beaker ceramics, groundstone and flaked stone tools, bone tools, limestone idols, other limestone objects, and copper implements (see Spindler and Gallay, 1973; Gallay et al., 1973).

Lastly, to assess potential outside genetic influence, the above were compared with seven extra-regional samples. Four are linked with northwestern Africa, i.e., Neolithic Capsian (n=24), precontact Canary Islanders (n=163), historic Kabyle (n=32) Berbers, and a fourth sample from an area once inhabited by Shawia Berbers (n=26). The remaining three samples derive from countries along the north shore of the Mediterranean: 1) Italy, which consists of 90 Roman and historic individuals, 2) Greece (n=77 individuals), dating from Classic and historic times, and 3) Turkey (n=40) that, along with individuals from nearby Cyprus, dates from Classic (Anatolian) to Ottoman times (Fig. 2). Although heterogeneous in composition, particularly the latter sample, they should at least yield some measure of characterization of eastern Mediterranean populations (re: Siret, 1913). All comparative samples, like the Portuguese, were recorded by the first author; thus, inter-observer error is not a concern and any intra-observer recording variation was found to be minimum and random. Background information has been presented elsewhere (Horwath 2012; Horwath et al., 2014; Irish, 1993, 1998b, 1998c, 2000). For quick reference, summary data for all 10 samples (n=578 total individuals) are listed in Table 1.

[FIGURE 2 HERE]

[TABLE 1 HERE]

1.3 Methods

1.3.1 Dental trait recording

Up to 125 nonmetric crown, root, and osseous oral traits were recorded in each dentition. For those with bilateral expression, both antimeres were recorded and, allowing for asymmetry, the side with the greatest expression was counted (Turner and Scott, 1977); the aim is to identify each trait's maximum genetic potential for each individual (Turner, 1985; Turner et al., 1991). Thirty-six of these traits (see list in Table 2) used by the first author in earlier

dental studies (e.g., Irish, 1993, 1997, 2005, 2006, etc.) were recorded for the following analyses. Except for UI1 midline diastema (Irish, 1993), all are part of the Arizona State University Dental Anthropology System (ASUDAS); the traits are recorded with the aid of 23 rank-scale reference plaques that help standardize scoring by providing representations of minimum, maximum, and intermediate expressions (Scott and Turner, 1997; Turner et al., 1991). The ASUDAS traits have a range of advantages (Irish, 1993, 1998b, 1998c 2005, 2006), but in brief they: 1) can be recorded despite slight attrition (though see below) or are unaffected by it in the cases of root and osseous traits, 2) have nominal inter- and intra-observer error rates when recording (as above), 3) are easy to identify, and 4) represent all dental morphogenetic fields. Of greater importance, the traits have a high genetic component in expression (Larsen, 1997; Martín-Torres et al., 2007; Rightmire, 1999; Scott, 1973; Scott and Turner, 1997), and they are evolutionarily conservative, making them excellent markers for biodistance analyses (Larsen, 1997). Another advantage is an overall lack of sexual dimorphism (Bermúdez de Castro, 1989; Hanihara, 1992; Irish, 1993; Scott, 1973, 1980; Smith and Shegev, 1988), to permit the pooling of sexes for maximizing sample size. A complete description of the ASUDAS is presented in Turner et al. (1991) and Scott and Turner (1997).

1.3.2 Quantitative analyses

All rank-scale ASUDAS traits were first dichotomized into categories of present or absent, based on their appraised morphological thresholds (Nichol, 1990; Scott, 1973) according to standard procedure (Irish, 1993; Turner, 1985, 1987). Dichotomization simplifies the tabulation of trait frequencies for presentation, and is necessary before these data can be compared with most distance statistics, including the mean measure of divergence (MMD) as used here (Berry and Berry, 1967; Green and Suchey, 1976; Harris and Sjøvold, 2004; Irish,

2010; Sjøvold, 1973, 1977). However, relative to prior studies (Irish, 1993, 1997, 2005, 2006, etc.) breakpoints that define “presence” were shifted upward for two important traits: UM1 Carabelli’s (by 3 grades to describe cusp expression only) and LP2 lingual cusp (1 grade, to better discern cusp number). This strategy minimizes the potential for grade shift and sampling biases when scoring traits in the present samples, which exhibit marked attrition differences among geographic regions; it also helps to maintain all-important sample size (Burnett et al., 2013; Burnett, 2016).

Next, the MMD was used to estimate among-sample phenetic affinities by calculating a measure of dissimilarity between each sample pair, i.e., high values indicate divergence and vice versa. Beyond holding several advantages over other distance measures (Irish, 2010) the MMD works with pooled sample data, to better address missing data in archaeological remains and the use of composite individuals. However, it is important to thoroughly edit these data. Traits with little or no contributory information should be deleted (Harris and Sjøvold, 2004). Those which are invariant can be recognized qualitatively, whereas traits that are the least, or conversely most, likely to drive inter-sample variation may be quantitatively identified, e.g., with correspondence analysis (Irish, 2005, 2006) or, like in the present study, principal components analysis (PCA) (Irish, 2016; Irish and Guatelli-Steinberg, 2003). The MMD distances should also be based on as many traits as possible, although none should be highly inter-correlated – which could render the results inaccurate (Sjøvold, 1977). To identify such traits the rank-scale data were submitted to Kendall’s tau-*b* correlation coefficient.

The MMD formula contains the Freeman and Tukey angular transformation to correct for low ($\leq .05$) or high ($\geq .95$) trait frequencies and small sample sizes ($n \geq 10$) (Green and Suchey, 1976; Sjøvold, 1973, 1977). To determine if two samples differ significantly, the

MMD value is compared with its standard deviation (SD). If the $MMD > 2 \times SD$, the null hypothesis of $P_1 = P_2$ (where P = sample population) is rejected at the 0.025 level. The MMD and standard deviation formulae, rationale for assessing significance, and other information can be accessed elsewhere (Irish, 2010; Sjøvold, 1977). Beyond presenting an MMD distance matrix, interval-level multi-dimensional scaling (MDS) (Kruskal and Wish, 1978) in SPSS 23.0 Procedure Alscal was used to create 3D spatial representations of the sample variation to aid interpretation.

Finally, the correlation between MMD and geographic distances in km was calculated with a two-tailed Mantel test (Smouse et al., 1986), and both distances were used as coordinates to plot pertinent sample pair relationships in 2D. These methods help explore if any Portuguese samples appear more similar to an extra-regional group than expected, under the assumption that genetic (and phenetic) relatedness among populations decreases exponentially as spatial distance increases (Relethford, 2004). Gene flow, the causative agent with isolation-by-distance (Wright, 1943), cannot pertain directly to the present diachronic samples; nevertheless, some indication of potential African or Mediterranean influence may be obtained to supplement the MMD results. The Geographic Distance Matrix Generator (vers. 1.2.3) (Ersts, 2014) was employed to calculate inter-sample straight line distances. The latter do not reflect reality on the landscape; however, because all comparative sample locations (especially eastern Mediterranean) are approximations, as would be potential migration routes toward Iberia, linear distances (though underestimates) should be less biased for analytical purposes.

1.4 Results

Percents of individuals that express each trait and the total number scored are listed in Table 2. The ASUDAS present/absent dichotomies are provided under each trait name. Samples

with <10 trait observations adversely impact several traits, especially CAP (see table and figure captions for the key to three-letter abbreviations). The BOL and PAI samples are affected similarly for palatine torus, midline diastema, and rocker jaw. Such data are unlikely to be characteristic of the parent populations; results using them should be interpreted with caution and/or they should be addressed during the trait editing process.

[TABLE 2 HERE]

That said, the MMD is a very robust statistic, so to obtain a preliminary indication of affinities among samples all 36 traits were first compared. The distance matrix is presented in Table 3, and the MDS solution from it (Fig. 3) provides a good representation, where the r^2 value is 0.897 and Kruskal's stress formula 1 is 0.113; for the latter value <0.10 is considered excellent and ≥ 0.15 unacceptable (Borgatti, 1997). Many significant MMD distances ($p \leq 0.025$) reflect the trait variation evident across samples (Table 2) including, among others, interruption groove UI1 [range of 0-46% (or 60% in the small CAP sample)], distal accessory ridge UC (8-74 %), rocker jaw (0-30%), and torsomolar angle LM3 (0-31%). However, the loose association of samples by geographic region in the figure (CDM/BOL/PAI, GRK/ITY/TRK, and CAN/CAP/KAB/SHA) is indicative of relative among-sample trait uniformity from those regions (again see Table 2).

[TABLE 3 and FIGURE 3 HERE]

Trait editing was then conducted as noted above. Non-contributory traits odontome P1-P2 (0-3% across samples), midline diastema (0-5%), mandibular torus (0-4%), and protostylid LM1 (0%) were removed. Percent data in the remaining 32 traits were submitted to PCA to identify less evident non-contributors in nine of the 10 total samples; CAP was not included after judging that many of its small and likely non-representative trait frequencies artificially increased several resulting loadings.

Eight components (eigenvalues >1.0) account for 100% of the total variance, though the PCA scree plot (not shown) indicates that the first three (58% variance) are most important; their un-rotated loadings are provided in Table 4. Those traits having very strong positive or negative loadings (i.e., >|0.5|) drive most of the variation as depicted in a graph of the group component scores in Figure 4. For Comp 1, loadings between 0.555 and 0.891 for tuberculum dentale UI2, distal accessory ridge UC, cusp 5 UM1, enamel extension UM1, root number UM2, anterior fovea LM1, deflecting wrinkle LM1, and cusp 7 LM1 push samples with high percents of these traits nearer the positive end of the x-axis (BOL, PAI, and to a lesser extent, CDM and ITY). Negative loadings of -0.589 to -0.855 for interruption groove UI2, congenital absence UM3, rocker jaw, cusp number LM2, and torsomolar angle LM3 typify samples toward the x-axis' opposite end (e.g., TRK, GRK, SHA). Traits of greater importance are also evident on the y- (Comp 2) and z-axes (Comp 3). As such, palatine torus, shoveling UI1, double shoveling UI1, hypocone UM2, and lingual cusp LP2 (loadings <|0.5| on all axes) were dropped because they are mostly noncontributory. Varimax rotation, which maximizes differences between large and small loadings, provided similar results (not shown).

Kendall's tau-*b* correlations among the remaining traits revealed high values (>|0.5|) for molar cusp number LM1 with: molar cusp number LM2, deflecting wrinkle LM1, and cusp 7 LM1 ($\tau_b = 0.573, 0.670, \text{ and } 0.818$, respectively). Similarly high values were found between c1-c2 crest LM1 and anterior fovea LM1 and, again, deflecting wrinkle LM1 with cusp 7 LM1 ($\tau_b = 0.657, 0.787, \text{ and } 0.706$). Therefore, together with lower loadings than other non-correlated traits (Table 4), molar cusp number LM1 and mandibular torus were dropped. As a result 25 traits, as denoted by asterisks, in Table 4 were used in the final MMD comparison.

[TABLE 4 and FIGURE 4 HERE]

This MMD distance matrix is presented in Table 5. Heterogeneity is again evident by the number of significantly different MMD distances. However, with a focus on the three Portuguese samples, the greater emphasis on divergence after deleting invariant and other traits that are non-contributory served to make CDM, BOL, and PAI, more distinct – from the other samples and one another (i.e., note the increase in MMD distances). The MDS solution again provides a good representation of the matrix, with the r^2 and Kruskal's stress formula 1 values improving to 0.919 and 0.109, respectively. Greater separation among the Portuguese samples is evident, though the new configuration (Fig. 5) otherwise looks similar in patterning to the 36-trait MDS graph (Fig. 3) in the association of samples by geographic region. To quantify this apparent correspondence, the 25-trait MMD distances were compared to geographic distances among sites and/or regions (Table 6); the Mantel correlation between matrices, $r=0.247$ ($p=0.102$), is positive though weak (per Cohen, 1988).

[TABLES 5-6 and FIGURE 5 HERE]

Lastly, CDM, BOL, and PAI were plotted individually with the rest of the samples using pairwise geographic and MMD distances as coordinates on the x- and y-axes in Figures 6-8. In each plot, a linear equation reference line (solid black) with a slope (b) of 1 and y-intercept (a) of 0 is provided (i.e., $y = 0 + 1x$, where $y = a + bx$) to simply illustrate where the other Portuguese and extra-regional samples would be located if a 1:1 correspondence existed between spatial and phenetic distances. The real sample locations identify those which are phenetically closer to the respective Portuguese sample than expected (i.e., below reference line), and vice versa (above line), relative to geographic separation. By way of comparison, the actual regression coefficient is also provided as the line of best fit (dashed) through the data cloud in each figure. Assuming that phenetic affinity is a function of spatial separation

(as above), coefficients of determination were calculated via the linear regression procedure in SPSS 23.0 for CDM ($r^2=0.095$, $r=0.308$, $p=0.420$), BOL ($r^2=0.290$, $r=0.539$, $p=0.135$), and PAI ($r^2=0.764$, $r=0.874$, $p=0.002$). Thus for CDM, 9.5% of the variability in MMD distances is associated with, or explained by, variability in the geographic distances. The corresponding values for BOL and PAI are 29.0% and 76.4%.

[FIGURES 6-8 HERE]

1.5 Discussion

Our overarching goal in this analysis was to ascertain if there is evidence of genetic input from northwestern African and/or eastern Mediterranean populations in late prehistoric peoples of the Portuguese Estremadura. Several high-frequency African traits (Irish, 1993, 1997, 1998a, 1998b, 1998c) are present in CDM, BOL, and/or PAI, including: UI1 labial curvature, Bushman canine UC, and LM1 cusp 7 (Table 2; also Table 4 loadings). In accordance, the 36- and, particularly, 25-trait MMD distances (Tables 3 and 5) and MDS scatterplots (Figs. 3 and 5; also see Fig. 4 PCA plot) indicate that the KAB and, to a lesser extent, SHA Berber samples, share very low and mostly insignificant phenetic distances with CDM and BOL (25-trait MMD between 0.002 and 0.070), though not with PAI (0.125-0.128). These findings support Cunha's (2012) dental study of Guadiana basin samples, from southwest Portugal, which found some traits suggestive of African links. And, as mentioned, archaeological evidence of African-sourced raw materials and artifacts is documented in southern Iberia (Harrison and Gilman, 1977; Manen et al., 2007), including ivory from North Africa at the site of Zambujal, a Copper Age settlement just 2 km south of Cova da Moura and Bolores (Sangmeister and Schubart, 1981). Perhaps an African contribution to the Estremadura gene pool did take place. However, ubiquitous traits in eastern Mediterranean populations, like UM2 hypocone reduction (grades <3) and two-rooted

LC (Scott and Turner, 1997), also occur in the Portuguese. The same MMD matrices and inter-sample plots identify a west-east affinity, most notably BOL (25-trait MMD=0.031) and, to a lesser degree, CDM (0.043) with ITY. The PAI sample (0.108), from farther north in the Estremadura, is again an exception (discussed below). Imported artifacts and materials from the east are also known (Schumacher et al., 2009). So genetic exchange with eastern Mediterranean peoples may too have occurred.

Of course, these extra-regional dental similarities may simply be a function of Portugal's proximity to northwest Africa and Italy relative to more geographically divergent samples (per Relethford, 2004; Wright, 1943), as in describing a vast, circum-Mediterranean cline; a positive though weak correlation does exist between the MMD and geographic straight-line distances. Going a step further, these affinities may reflect local mobility, in the form of interaction with other Iberian peoples who, in turn, had genetic contact with outside populations (Cunha, 2012). Otherwise, the resemblance of CDM and BOL to KAB, SHA, and ITY may be indicative of direct contact (*à la* Siret, 1913), as in actual migration and/or gene flow into southern Iberia, specifically the Estremadura. Based on isotopic evidence, Waterman et al. (2014) did find that four of 12 skeletons sampled at CDM were non-local – though two were simply from other parts of Iberia. In support of external influence, only 9.5% of the variability in MMD distances for CDM and 29% for BOL, as noted, is associated with the variability in geographic distances with the remaining samples. These results imply that at least some of the latter are not plotted where they “should” be (Figs. 6-7), assuming phenetic affinity is directly related to spatial separation. Focusing only on extra-regional comparisons, CAP and, to a lesser extent, CAN are plotted above the solid black reference line, meaning they are more divergent phenetically from CDM than anticipated based on geographic location. Below this line the opposite is true for KAB and ITY, which share low

MMD distances with CDM (Table 5), and SHA, GRK, and TRK, which differ significantly from this Portuguese sample. With some variation, inter-sample patterning for BOL is similar (Fig. 7). Therefore, based on the weight of the evidence, namely: 1) extra-regional traits in the Portuguese, 2) pertinent low and insignificant MMD distances, and 3) closer-than-expected plots of KAB and ITY, among others, neither of the hypotheses in the introduction can be rejected – though with qualification. Specifically, there is no significant difference between one or more samples from the south-central Portuguese Estremadura (CDM and BOL) and northwest Africa (KAB and SHA), nor between CDM and BOL and the eastern Mediterranean (ITY). The PAI sample does not exhibit overt extra-regional input based on these analyses. It differs significantly from all comparative samples (Table 5), and any similarities do indeed appear to be mostly a function of geographic distance ($r^2=0.764$, $p=0.002$) (Fig. 8).

Within the Estremadura, dental heterogeneity is evident. The average intra-region 25-trait MMD among CDM, BOL, and PAI, is 0.058 (calculated from Table 5); despite issues affecting the composition of several comparative samples (above), the corresponding value for northwest Africa (CAN, CAP, KAB, and SHA) is 0.025, whereas that for the eastern Mediterranean (ITY, GRK, and TRK) is 0.015. The phenetic similarity between CDM and BOL is suggestive of short term continuity in the Rio Sizandro valley, but the MMD of 0.036 ($p<0.025$) is high considering a separation of 2 km between sites (Figs. 6-7). The uniqueness of PAI is clear (Fig. 8). Though most akin to CDM (MMD=0.058) and BOL (0.079), it is significantly different ($p>0.025$) from both; Pai Mogo 1 is situated 23 km north from Cova da Moura and Bolores yet, among other possible factors, this relatively small distance interspersed by two river valleys was seemingly a contributor to substantial biocultural isolation. As mentioned, differences in diet, artifacts, and architecture were also reported

between areas (Gallay et al., 1973; Guiry et al., 2016; Spindler and Gallay, 1973; Waterman et al., 2014, 2015). This heterogeneity, the divergence of PAI in particular (at least compared to the samples in this study), reminds us that our understanding of the Iberian Late Neolithic/Copper Age must include recognition of variation – both regional and community-wide. Inhabitants of communities at this time may have been diverse in biological origin and, likely, the experiences of their worlds. Such diversity may help identify possible tensions within society that generated profound social changes by the Early Bronze Age, c. 2200 BCE, including the abandonment of settlements and emergence of new, individualized mortuary practices.

Lastly, relationships among all 10 samples are instructive. The demonstrable distinction by geographic region (e.g., Figs. 3 and 5) and relative intra-region similarity (above), quantified by the Mantel correlation ($r=0.247$), support the viability of these dental nonmetric analyses. In fact, when northwest African-associated CAN is dropped from study, the r -value between the 25-trait MMD and geographic distances increases substantially to 0.05 ($p=0.002$), a strong positive correlation (Cohen, 1988). Canary Islanders are considered by researchers to be something of an outlier. Though descendants of early (potentially Neolithic) Berbers, the remote island location led to genetic isolation in these non-seagoing peoples, with only subsequent one-way gene flow – mostly from Europeans beginning in the 15th century CE (Bermúdez de Castro, 1989; Guatelli-Steinberg et al., 2001; Irish, 1993; Mercer, 1980; Schwidetsky, 1963). The low and insignificant MMD distances of CAN with both Berber samples and Neolithic CAP, plus comparable values with geographically distant but younger ITY, GRK, and TRK, though not older CDM, BOL, or PAI, are supportive of these known and suspected biological relationships. Other links between the northwest African and eastern Mediterranean samples are obvious (Tables 3 and 5, Figs. 3 and 5). As

well, high-frequency African traits are present to varying degrees in ITY, GRK, and TRK, and vice versa (Table 2). These findings are suggestive of more than just indirect contact among circum-Mediterranean populations, contra the strict anti-migrationism and diffusionism interpretations of some researchers.

In sum, our study has attempted to better understand the cultural and demographic dynamics of human communities within the Portuguese Estremadura, using three samples dated between 3700-2300 BCE. Results indicate intra-regional heterogeneity relative to samples from other Mediterranean regions, though with some measure of population continuity, particularly in the Sizandro River valley. That said, long-range genetic input from more distant, yet neighboring regions, such as northwestern Africa and the eastern Mediterranean, cannot be ruled out. These data parallel findings of low-level migration at Cova da Moura and elsewhere in the Late Neolithic and Copper Age, based upon isotopic analyses of human remains in collective burials (Waterman et al., 2014). This correspondence may imply that exotic materials recovered from settlement and burials of this time are indicative of down-the-line exchange but, again, actual movement of peoples may be possible based on the present quantitative analyses. As usual, it was likely a combination of influences that led to the emergence of social complexity during the Late Neolithic/Copper Age in the 4th and 3rd millennia BCE, not mutually exclusive migrationism versus autochthonism. Whatever the case, all explanatory models should remain in play until fully discredited. Further dental nonmetric and (planned) genetic analyses of human remains in burials dating to earlier and contemporary periods in Portugal, and greater Iberia, should help to clarify these findings.

Acknowledgements

Our appreciation is extended to Ana Catarina Sousa for the valuable comments about the history of Portuguese prehistory, and Isabel Luna from the Museu Municipal Leonel Trindade – Torres Vedras (Portugal).

Funding: This work was supported by the National Science Foundation, to JDI (BNS-9013942, BNS-0104731) and KTL (BCS-1153568).

References

- Aranda Jimenez, G., Montón-Subías, S., Sánchez Romero, R., 2015. The Archaeology of Bronze Age Iberia. Argaric Societies. Routledge, New York.
- Belo, A.R., Trindade, L., Ferreira, O. da Veiga., 1961. A gruta da Cova da Moura (Torres Vedras). Comunicações dos Serviços Geológicos de Portugal 45, 391-418.
- Bermúdez de Castro, J.M., 1989. The Carabelli trait in human prehistoric populations of the Canary Islands. Hum Biol. 61, 117–131.
- Berry, A.C., Berry, R.J. 1967. Epigenetic variation in the human cranium. J Anat. 101, 361-379.
- Boaventura, R., 2009. As antas e o Megalitismo da região de Lisboa. Doctoral Thesis. University of Lisbon, Lisbon.
- Boaventura, R., Ferreira, M.T., Neves, M.J., Silva, A.M., 2014. Funerary practices and anthropology during the middle-late Neolithic (4th and 3rd Millennia BCE) in Portugal: old bones, new insights. Anthropologie LII, 183-205.
- Borgatti, S.P. 1997. Multidimensional scaling. Available at: www.analytictech.com/borgatti/mds.htm accessed on August 15, 2016.
- Bosch Gimpera, P. 1932. Etnologia de la Península Ibérica. Editorial Alpha, Barcelona.
- Burnett, S.E., 2016. Crown wear: Identification and categorization, in: Irish, J.D., Scott, G.R. (Eds.), A Companion to Dental Anthropology. Wiley Blackwell, London, pp. 415–432.
- Burnett, S.E., Irish, J.D., Fong, M.R. 2013. Wears the problem? Examining the effect of

- dental wear on studies of crown morphology, in: Scott, G.R., Irish, J.D. (Eds.), *Anthropological Perspectives on Tooth Morphology: Genetics, Evolution, Variation*. Cambridge University Press, Cambridge, pp. 535–553.
- Chapman, R., 1990. *Emerging Complexity*. Cambridge University Press, Cambridge.
- Costa, M.E., Díaz-Zorita, M., García Sanjuán, L., Wheatley, D., 2010. The Copper Age settlement of Valencina de la Concepción (Seville, Spain): Demography, metallurgy and spatial organization. *Trabajos de Prehistoria* 67, 85–117. doi:10.3989/tp.2010.10032.
- Cunha, C. 2012. Caracterização da morfologia dentária no Médio Guadiana no Neolítico Final – Calcolítico. Fundamentação para o mapeamento morfológico das populações locais na Pré-História Recente. *Apontamentos de Arqueologia e Património*. 8. Lisboa. *Era Arqueologia*. 49-55
- Ersts, P.J. 2014. Geographic Distance Matrix Generator (version 1.2.3). American Museum of Natural History, Center for Biodiversity and Conservation. Available at: http://biodiversityinformatics.armh.org/open_source/gdrrg, accessed on September 7, 2016.
- Gallay, G., Spindler, K., 1970. *Archaeologische und Anthropologische Betrachtungen zu den Neolithisch-Kupferzeitlichen Funden aus der Cova da Moura /Portugal*. Madrider Mitteilungen 11. Heidelberg F.H. Kerle Verlag.
- Gallay, G., Spindler, K., Trindade, L. Ferreira, O. da Veiga, 1973: O monumento préhistórico de Pai Mogo (Lourinhã). *Associação dos Arqueólogos Portugueses*, Lisbon.
- Green, R., Suchey, J. 1976. The use of inverse sine transformation in the analysis of non-metrical data. *Am J Phys Anthropol*. 45, 61–68.

Gonçalves, V., Sousa, A.C., Costeira, C., 2013. Walls, gates and towers. Fortified settlements in the south and centre of Portugal: some notes about violence and walls in the 3rd millennium BCE. *Cadernos de Prehistoria y Arqueologia de la Universidade de Granada* 23, 35-97.

Guatelli-Steinberg, D., Irish, J.D., Lukacs, J. 2001. Canary Island - North African population affinities: Measures of divergence based on dental morphology. *Homo* 52, 173-188.

Guiry, E.J., Hillier, M., Boaventura, R., Silva, A.M., Oosterbeek, L., Tomé, T., Valera, A. Cardoso, J.L., Hepburn, J.C., Richards, M.P., 2016. The transition to agriculture in south-western Europe: new isotopic insights from Portugal's Atlantic coast. *Antiquity* 90, 604-616 (doi:10.15184/aqy.2016.34).

Hanihara, T. 1992. Dental and cranial affinities among populations of East Asia and the Pacific: the basic populations in East Asia, IV. *Am J Phys Anthropol.* 88, 163-182.

Harris, E.F., Sjøvold, T. 2004. Calculation of Smith's mean measure of divergence for intergroup comparisons using nonmetric data. *Dent Anthropol.* 17, 83–93.

Harrison, R.J., Gilman, A., 1977. Trade in the second and third millennia BC between the Maghreb and Iberia, in: Markotic, V. (Ed.), *Ancient Europe and the Mediterranean*. Aris and Phillips, Warminster, pp. 89-104.

Horwath, B. 2012. Diet and Affinity from the Middle Neolithic to Early Bronze Age, Estremadura, Portugal: A Comparison of Human Dental Remains from Feteira II and Bolores. M.A. Thesis. University of Alaska, Fairbanks.

- Horwath, B., Waterman, A.J., Lillios, K.T., Irish, J., 2014. Assessing change in diet and biological affinity between the 4th and 3rd millennia cal BCE in the Portuguese Estremadura: A preliminary dental comparison of Feteira II and Bolores. *Homo* 65, 87-100.
- Hurtado, V.M., 1997. The dynamics of the occupation of the middle basin of the river Guadiana between the fourth and second millennia BC, in: Diaz-Andreu, M. Keay, S. (Eds.), *The Archaeology of Iberia. The Dynamics of Change*. Routledge, London. pp. 98-127.
- Irish, J.D. 1993. Biological Affinities of Late Pleistocene through Modern African Aboriginal Populations: The Dental Evidence. Ph.D. Dissertation. Arizona State University, Tempe.
- Irish, J.D. 1997. Characteristic high- and low-frequency dental traits in sub-Saharan African populations. *Am J Phys Anthropol.* 102, 455–467.
- Irish, J.D. 1998a. Ancestral dental traits in recent Sub-Saharan Africans and the origins of modern humans. *J Hum Evol.* 34, 81–98.
- Irish, J.D. 1998b. Dental morphological affinities of late Pleistocene through recent sub-Saharan and North African peoples. *Bull Mem Soc Anthropol Paris* 10, 237–272.
- Irish, J.D. 1998c. Diachronic and synchronic dental trait affinities of Late and post-Pleistocene peoples from North Africa. *Homo* 49, 138–155.
- Irish JD. 2000. The Iberomaurusian enigma: North African progenitor or dead end? *J Hum Evol.* 39, 393–410.
- Irish, J.D. 2005. Population continuity versus discontinuity revisited: dental affinities among Late Paleolithic through Christian era Nubians. *Am J Phys Anthropol.* 128, 520–535.

Irish, J.D. 2006. Who were the ancient Egyptians? Dental affinities among Neolithic through post-dynastic peoples. *Am J Phys Anthropol.* 129, 529–543.

Irish, J.D. 2010. The mean measure of divergence (MMD): its utility in model-free and model-bound analyses relative to the Mahalanobis D2 distance for nonmetric traits. *Am J Hum Biol.* 22, 378–395.

Irish, J.D. 2016. Who were they really? Model-free and model-bound dental nonmetric analyses to affirm documented population affiliations of seven South African “Bantu” samples, *Am J Phys Anthropol.* 159, 655-670.

Irish, J.D., Guatelli-Steinberg, D. 2003. Ancient teeth and modern human origins: an expanded comparison of African Plio-Pleistocene and recent world dental samples. *J Hum Evol.* 45, 113-144.

Irish, J.D., Black, W., Sealy, J., Ackermann, R. 2014. Questions of Khoesan continuity: dental affinities among the indigenous Holocene peoples of South Africa. *Am J Phys Anthropol.* 155, 33–44.

Jackes, M., Silva, A.M., Irish, J., 2001. Dental morphology: a valuable contribution to our understanding of prehistory. *J. of Iberian Archaeology* 3, 97-119.

Kintigh, K.W., Altschul, J.H., Beaudry, M.C., Drennan, R.D., Kinzig, A.P., Kohler, T.A., Limp, W.G., Maschner, H.D.G., Michener, W.K., Pauketat, T.R., Peregrine, P., Sabloff, J.A., Wilkinson, T.J., Wright, H.T., Zeder, M.A., 2014. Grand challenges for archaeology. *PNAS* 111, 879-880; doi:10.1073/pnas.1324000111.

Kruskal, J.B., Wish, M. 1978. *Multidimensional Scaling*. Sage Publications, Beverly Hills.

- Larsen, C.S. 1997. *Bioarchaeology*. Cambridge University Press, Cambridge.
- Lillios, K.T., 2015. Practice, process, and social change in third millennium BC Europe: A view from the Sizandro Valley, Portugal. *European Journal of Archaeology* 77, 245-258.
- Lillios, K.T., Waterman, A.J., Mack, J.M., Artz, J.A, Nilsson-Stutz, L., 2015. In Praise of Small Things: Excavations at the Late Neolithic-Early Bronze Age Burial of Bolores (Torres Vedras), Portugal. *British Archaeological Reports*, Oxford.
- Manen C., Marchand G., Carvalho A. F., 2007. Le Néolithique ancien de la péninsule ibérique: vers une nouvelle évaluation du mirage africain, in: *Congrès du Centenaire (2007). Un siècle de construction du discours scientifique en Préhistoire*, Société Préhistorique Française. Vol. 3, pp 133-135.
- Martínez Santa-Olalla, J., 1941. Esquema paleotnológico de la Península hispánica, in: *Martínez Santa-Olalla, J. (Ed.), Corona de Estudios que la Sociedad Española de Antropología, Etnología y Prehistoria dedica a sus mártires. Consejo superior Investigación Científica de Madrid*, pp. 144-166.
- Martinón-Torres, M., Bermúdez de Castro, J., Gomez-Robles, A., Arsuaga, J., Carbonell, E., Lordkipanidze, D., Manzi, G., Margvelashvili, A. 2007. Dental evidence on the hominin dispersals during the Pleistocene. *Proc Natl Acad Sci*. 104, 13279–13282.
- Mercer, J. 1980. *The Canary Islanders: Their Prehistory, Conquest and Survival*. Rex Collings, London.

Morán, E., Parreira, R., 2003. O povoado calcolítico de Alcalar (Portimão) na paisagem cultural do Alvor no III milénio antes da nossa Era. Recintos murados da pré-história. Porto/Coimbra, FLUP/CEAUCP: 307-327.

Morgado Rodríguez, A., Lozano Rodríguez, J.A., García Sanjuán, L., Odriozola Lloret, C., and Lucíañez Triviño, M., 2015. The allure of rock crystal in southern Iberian Copper Age: Technical mastery and distinguished objects at Valencina de la Concepción (Seville, Spain). *Quaternary International*. doi:10.1016/j.quaint.2015.08.004.

Müller, R., Goldenberg, G., Kunst, M., Bartelheim, M., Pernicka, E., 2007. Zambujal and the beginnings of metallurgy in southern Portugal, in: La Niece, S., Hook, D., Craddock, P.T. (Eds.), *Metals and Mines. Studies in Archaeometallurgy. Papers from the Conference Metallurgy: a Touchstone for Cross-cultural Interaction* (British Museum), London 28-30, April 2005, pp. 15-26.

Murillo-Barroso, M., Costa Caramé, M.E., Díaz-Guardamino Uribe, M., García Sanjuán, L., Mora Molina, C., 2015. A reappraisal of Iberian Copper Age goldwork: Craftmanship, symbolism and art in a non-funerary gold sheet from Valencina de la Concepción. *Cambridge Archaeological Journal* 25, 565-596.

Murillo-Barroso, M.; Martínón-Torres, M., 2012. Amber source and trade in the Prehistory of the Iberian Peninsula. *European Journal of Archaeology* 15, 187 – 216.

Nichol, C.R. 1990. *Dental Genetics and Biological Relationships of the Pima Indians of Arizona*. Ph.D. Dissertation. Arizona State University, Tempe.

- Odriozola, C., Villalobos García, R., Burdidge, C., Boaventura, R., Sousa, A.C., Rodríguez-Ariza, O., Parilla-Giraldez, R., Prudêncio, M.I., Dias, M.I. 2016. Distribution and chronological framework for Iberian variscite mining and consumption at Pico Centeno, Encinasola, Spain. *Quaternary Research* 85, 159-176.
- Rightmire, G.P. 1999. Dental variation and human history. *Rev Archaeol.* 20, 1–3.
- Sangmeister, E., Schubart, H., 1981. Zambujal die Gräbungen 1964 bis 1967. Philipp von Zabern Mainz. *Madriider Beiträge*, Band 5.
- Schuhmacher, T.X., Cardoso, J.L., Banerjee, A., 2009. Sourcing African ivory in Chalcolithic Portugal. *Antiquity* 83, 983-997.
- Schwidetsky, I. 1963. La Población Prehispánica de las Islas Canarias. Museo Arqueológico de Tenerife, Santa Cruz de Tenerife.
- Scott, G.R. 1973. Dental Morphology: A Genetic Study of American White Families and Variation in Living Southwest Indians. Ph.D. Dissertation. Arizona State University, Tempe.
- Scott, G.R. 1980. Population variation of Carabelli's trait. *Hum Biol.* 52, 63–78.
- Scott, G.R., Turner, C.G. II., 1997. *The Anthropology of Modern Human Teeth: Dental Morphology and its Variation in Recent Human Populations*. Cambridge University Press, Cambridge.
- Scott, G.R., Yap Potter, R.H., Noss, J.F., Dahlberg, A.A., Dahlberg, T. 1983. The dental morphology of Pima Indians. *Am J Phys Anthropol.* 61, 13–31.

Silva, A.M., 2002. Antropologia funerária e Paleobiologia das populações portuguesas (litorais) do Neolítico final/Calcolítico. Doctoral thesis. Department of Anthropology, University of Coimbra, Coimbra (Portugal).

Silva, A.M., 2003: Portuguese populations of Late Neolithic and Chalcolithic periods exhumed from collective burials: an overview. *Anthropologie* XLI/1-2, 55-64.

Siret, L. 1913. Questions de chronologie et d'ethnographie ibériques. I. De la fin du quaternaire a la fin du bronze. P. Geuthner, Paris.

Sjøvold, T. 1973. Occurrence of minor non-metrical variants in the skeleton and their quantitative treatment for population comparisons. *Homo* 24, 204–233.

Sjøvold, T. 1977. Non-metrical divergence between skeletal populations: The theoretical foundation and biological importance of C.A.B. Smith's mean measure of divergence. *Ossa* 4 Suppl. 1, 1–133.

Smith, P., Shegev, M. 1988. The dentition of Nubians from Wadi Halfa, Sudan: an evolutionary perspective. *J Dent Assoc S Afr.* 43, 539–541.

Smouse, P.E., Long, J.C., Sokal, R.R. 1986. Multiple regression and correlation extensions of the Mantel test of matrix correspondence. *Syst Zool.* 35, 627–632.

Spindler, K., 1981. Cova da Moura-Die Besiedlung des Atlantischen Küstengebietes Mittelportugals vom Neolithikum bis an das Ende der Bronzezeit. *Madriider Beiträge* 7,

Zabern-Verlag. Mainz.

Spindler, K., Gallay, G., 1973. Kupferzeitliche Siedlung und Begraebnisstaetten von Matacões in Portugal. Mainz am Rhein, Verlag Philipp von Zabern.

Sousa, A.C., Gonçalves, V., 2012. In and out, Tecnologias, símbolos e cultura material. Interacções e identiades regionais no centro e sul de Portugal no 3º milénio A.N.E. Congrès International Xarxes al Neolític. Neolithic Networks. Rubricatum 5, 383-392.

Thomas, J.T., 2011. Fashioning identities, forging inequality: Late Neolithic/Copper Age personal ornaments of the Portuguese Estremadura. European Journal of Archaeology 14, 29-39.

Turner, C.G., II. 1985. Expression count: a method for calculating morphological dental trait frequencies by using adjustable weighting coefficients. Am J Phys Anthropol. 68, 263-268.

Turner, C.G., II. 1987. Late Pleistocene and Holocene population history of East Asia based on dental variation. Am J Phys Anthropol. 73, 305–322.

Turner, C.G., II, Nichol, C.R., Scott, G.R. 1991. Scoring procedures for key morphological traits of the permanent dentition: the Arizona State University dental anthropology system, in: Kelley, M.A., Larsen, C.S. (Eds.), Advances in Dental Anthropology. Wiley-Liss, New York, pp. 13–32.

Turner, C.G., II, Scott, G.R. 1977. Dentition of Easter Islanders, in: Dahlberg, A.A., Graber, T.M. (Eds.), Orofacial Growth and Development. Mouton, The Hague, pp. 229–249.

Valera, A.C., Filipe, I., 2004. O povoado do Porto Torrão (Ferreira do Alentejo): novos dados

e novas problemáticas no contexto da calcolitização do Sudoeste peninsular. *ERA Arqueologia*, 6, 28–61.

Valera, A.C., Schumacher, T., Banerjee, A., 2015. Ivory in the Chalcolithic enclosure of Perdigões (South Portugal): the role of an exotic raw material. *World Archaeology* 47, 390-413.

Valera, A.C., Silva, A.M., Marquez Romero, J., 2014. La temporalidad del recinto de fosos de Perdigões: cronología absoluta de estruturas y praticas sociales. *SPAE* 23, 11-26.

Villalobos García, R., Odriozola, C.P., 2016. Organizing the production of variscite personal ornaments in later prehistoric Iberia: The mines of Aliste and the production sites of Quiruelas de Vidriales (Zamora, Spain). *European Journal of Archaeology* 1-21.

Waterman, A.J., Lillios, K.T., Tykot, R.H., Kunst, M., 2016. Environmental change and economic practices between the third and second millennia BC using isotope analyses of ovicaprid remains from the archaeological site of Zambujal (Torres Vedras), Portugal. *Journal of Archaeological Science: Reports* 5, 181-189.

Waterman, A.J., Peate, D.W., Silva, A.M., Thomas, J.T., 2014. In search of homelands: Using strontium isotopes to identify biological markers of mobility in late prehistoric Portugal. *Journal of Archaeological Science* 42C, 119-127.

Waterman, A.J., Tykot, R.H., Silva, A.M., 2015. Stable isotope analysis of diet-based social differentiation at late prehistoric collective burials in southwestern Portugal. *Archaeometry* 58, 131–151.

Zafra de la Torre, N., Hornos Mata, F., Castro López, M., 1999. Una macro-aldea en el origen del modo de vida campesino: Marroquíes Bajos (Jaén) c. 2500-2000 cal. ANE. Trabajos de Prehistoria 56, 77-102.

Figure legends

Figure 1. Origin locations of the three Portuguese Late Neolithic-Copper Age samples.

Figure 2. Origin locations of the Portuguese and comparative samples. See Table 1 and text Section 1.2 for sample compositions.

Figure 3. Three-dimensional MDS of 36-trait MMD distances among the three Portuguese and seven comparative samples. The three-letter sample abbreviations are defined in Tables 1-3 and 5-6.

Figure 4. Three-dimensional scatterplot of the first three components among the three Portuguese and six comparative samples (without CAP; see text) for 32 dental traits from Table 4. Accounts for 58.14% of the total variance (23.32% on x-axis, 19.47% on y-axis, and 15.34% on z-axis). Sample abbreviations defined in Tables 1-3 and 5-6.

Figure 5. Three-dimensional MDS of 25-trait MMD distances among the three Portuguese and seven comparative samples. The three-letter sample abbreviations are defined in Tables 1-3 and 5-6.

Figure 6. Two-dimensional scatterplot of the CDM sample relative to the other Portuguese and comparative samples based on geographic (x-axis) vs. phenetic (y-axis) distances. Solid black linear equation reference line with slope (b) of 1 and y-intercept (a) of 0 provided (i.e., $y = 0 + 1x$, where $y = a + bx$) to illustrate where the other samples would be if 1:1 correspondence existed between the distances. The actual regression coefficient is also provided as the line of best fit (dashed) through the data cloud in each figure (with equation provided). See text Section 1.4 for details.

Figure 7. Two-dimensional scatterplot of the BOL sample relative to the other Portuguese and comparative samples based on geographic (x-axis) vs. phenetic (y-axis) distances. Solid black linear equation reference line with slope (b) of 1 and y-intercept (a) of 0 provided (i.e., $y = 0 + 1x$, where $y = a + bx$) to illustrate where the other samples would be if 1:1 correspondence existed between the distances. The actual regression coefficient is also provided as the line of best fit (dashed) through the data cloud in each figure (with equation provided). See text Section 1.4 for details.

Figure 8. Two-dimensional scatterplot of the PAI sample relative to the other Portuguese and comparative samples based on geographic (x-axis) vs. phenetic (y-axis) distances. Solid black linear equation reference line with slope (b) of 1 and y-intercept (a) of 0 provided (i.e., $y = 0 + 1x$, where $y = a + bx$) to illustrate where the other samples would be if 1:1 correspondence existed between the distances. The actual regression coefficient is also provided as the line of best fit (dashed) through the data cloud in each figure (with equation provided). See text Section 1.4 for details.

Region

- ▼ Northern Mediterranean
- Northwest Africa
- Portugal

Region

- ▼ Northern Mediterranean
- Northwest Africa
- Portugal

TABLE 6. Symmetrical geographic straight-line distance matrix (km) among the Portuguese and comparative samples (using actual or approximated center of each site or geographic region)

	BOL	CDM	PAI	CAN	CAP	KAB	SHA	GRK	ITY	TRK
BOL	0	2.0	19.9	1367.3	1578.6	1359.9	1560.8	2750.5	1982.3	3745.5
CDM	2.0	0	19.7	1367.8	1578.1	1359.4	1560.3	2749.9	1981.7	3744.9
PAI	22.9	22.7	0	1385.1	1580.1	1362.4	1565.3	2748.2	1976.9	3742.8
CAN	1367.3	1367.8	1385.1	0	2451.9	2226.8	2303.5	3674.6	3072.9	4647.8
CAP	1578.6	1578.1	1580.1	2451.9	0	232.1	241.9	1230.8	722.7	2222.2
KAB	1359.9	1359.4	1362.4	2226.8	232.1	0	249.1	1461.9	894.7	2454.2
SHA	1560.8	1560.3	1565.3	2303.5	241.9	249.1	0	1374.3	958.2	2345.5
GRK	2750.5	2749.9	2748.2	3674.6	1230.8	1461.9	1374.3	0	851.4	997.7
ITY	1982.3	1981.7	1976.9	3072.9	722.7	894.7	958.2	851.4	0	1799.9
TRK	3745.5	3744.9	3742.8	4647.8	2222.2	2454.2	2345.5	997.7	1799.9	0

BOL=Bolores, CDM=Cova da Moura, PAI=Pai Mogo, CAN=Canary Islanders, CAP=NW African Capsian, KAB=NW African Kabyle Berbers, SHA=NW African Shawia Berbers, GRK=Greece, ITY=Italy, TRK=Turkey (see text Section 1.2 for details).

TABLE 5. Symmetrical MMD distance matrix for Portuguese and comparative samples based on 25 dental traits

	BOL	CDM	PAI	CAN	CAP	KAB	SHA	GRK	ITY	TRK
BOL	0	0.036	<u>0.079</u>	<u>0.073</u>	<u>0.114</u>	0.021	0.060	<u>0.123</u>	0.031	<u>0.118</u>
CDM	0.036	0	<u>0.058</u>	<u>0.111</u>	<u>0.159</u>	0.002	<u>0.070</u>	<u>0.110</u>	<u>0.043</u>	<u>0.081</u>
PAI	<u>0.079</u>	<u>0.058</u>	0	<u>0.185</u>	<u>0.170</u>	<u>0.125</u>	<u>0.128</u>	<u>0.187</u>	<u>0.108</u>	<u>0.248</u>
CAN	<u>0.073</u>	<u>0.111</u>	<u>0.185</u>	0	0.067	0.025	0.000	0.021	0.015	0.043
CAP	<u>0.114</u>	<u>0.159</u>	<u>0.170</u>	0.067	0	0.020	0.040	0.090	<u>0.081</u>	<u>0.132</u>
KAB	0.021	0.002	<u>0.125</u>	0.025	0.020	0	0.000	0.034	0.000	0.002
SHA	0.060	<u>0.070</u>	<u>0.128</u>	0.000	0.040	0.000	0	0.000	0.000	0.000
GRK	<u>0.123</u>	<u>0.110</u>	<u>0.187</u>	0.021	0.090	0.034	0.000	0	0.031	0.002
ITY	0.031	<u>0.043</u>	<u>0.108</u>	0.015	<u>0.081</u>	0.000	0.000	0.031	0	0.013
TRK	<u>0.118</u>	<u>0.081</u>	<u>0.248</u>	0.043	<u>0.132</u>	0.002	0.000	0.002	0.013	0

BOL=Bolores, CDM=Cova da Moura, PAI=Pai Mogo, CAN=Canary Islanders, CAP=NW African Capsian, KAB=NW African Kabyle Berbers, SHA=NW African Shawia Berbers, GRK=Greece, ITY=Italy, TRK=Turkey (see text Section 1.2 for details).

Underlined MMD distances indicate significant difference at the 0.025 level.

TABLE 4. Component loadings, eigenvalues, and variance explained for 32 traits in the three Portuguese and six of seven comparative samples

Trait	Comp 1	Comp 2	Comp 3
Winging UI1*	0.418	0.555	0.212
Labial Curvature UI1*	-0.465	-0.255	-0.745
Palatine Torus	-0.120	0.448	0.301
Shoveling UI1	0.245	-0.020	-0.183
Double Shoveling UI1	-0.034	-0.485	0.192
Interruption Groove UI2*	-0.589¹	0.131	0.432
Tuberculum Dentale UI2*	0.665	-0.140	-0.459
Bushman Canine UC*	-0.081	0.657	0.204
Distal Acc. Ridge UC*	0.678	-0.609	0.350
Hypocone UM2	-0.092	0.069	0.260
Cusp 5 UM1*	0.891	0.376	-0.060
Carabelli's Trait UM1*	-0.043	-0.772	-0.039
Parastyle UM3*	-0.031	-0.271	0.660
Enamel Extension UM1*	0.555	0.476	-0.226
Root Number UP1*	-0.202	-0.550	0.187
Root Number UM2*	0.706	0.374	0.283
Peg-Reduced UI2*	0.363	0.486	-0.693
Congenital Abs. UM3*	-0.721	0.235	0.317
Lingual Cusp LP2	-0.361	-0.271	0.167
Anterior Fovea LM1*	0.699	-0.555	-0.073
Groove Pattern LM2*	0.284	-0.029	0.816
Rocker Jaw*	-0.711	0.496	0.186
Cusp Number LM1	-0.018	-0.511	-0.234
Cusp Number LM2*	-0.591	0.222	0.471
Deflecting Wrinkle LM1*	0.668	-0.323	0.405
C1-C2 Crest LM1	0.077	0.706	0.085
Cusp 7 LM1*	0.605	0.546	0.141
Tomes' Root LP1*	0.436	-0.557	0.599
Root Number LC*	0.122	-0.408	-0.538
Root Number LM1*	0.150	0.587	-0.427
Root Number LM2*	0.327	0.381	0.519

Torsomolar Angle LM3*	-0.855	0.004	-0.179
Eigenvalue	7.463	6.230	4.910
Variance (%)	23.323	19.468	15.344
Total Variance	23.323	42.791	58.135

¹Values in bold-face indicate strong loadings ($>|0.5|$) as detailed in text.

*Denotes the 25 traits used in the final MMD comparison.

TABLE 3. Symmetrical MMD distance matrix for Portuguese and comparative samples based on 36 dental traits

	BOL	CDM	PAI	CAN	CAP	KAB	SHA	GRK	ITY	TRK
BOL	0	0.011	0.033	<u>0.055</u>	<u>0.084</u>	0.024	0.038	0.050	0.014	0.052
CDM	0.011	0	0.033	<u>0.086</u>	<u>0.118</u>	0.002	0.041	<u>0.057</u>	<u>0.047</u>	<u>0.049</u>
PAI	0.033	0.033	0	<u>0.140</u>	<u>0.110</u>	<u>0.072</u>	<u>0.080</u>	<u>0.113</u>	<u>0.079</u>	<u>0.158</u>
CAN	<u>0.055</u>	<u>0.086</u>	<u>0.140</u>	0	0.049	0.031	0.015	0.019	<u>0.031</u>	<u>0.058</u>
CAP	<u>0.084</u>	<u>0.118</u>	<u>0.110</u>	0.049	0	0.006	0.018	0.071	<u>0.090</u>	<u>0.098</u>
KAB	0.024	0.002	<u>0.072</u>	0.031	0.006	0	0.000	0.030	0.017	0.027
SHA	0.038	0.041	<u>0.080</u>	0.015	0.018	0.000	0	0.000	<u>0.042</u>	0.000
GRK	0.050	<u>0.057</u>	<u>0.113</u>	0.019	0.071	0.030	0.000	0	0.023	0.000
ITY	0.014	<u>0.047</u>	<u>0.079</u>	<u>0.031</u>	<u>0.090</u>	0.017	<u>0.042</u>	0.023	0	0.019
TRK	0.052	<u>0.049</u>	<u>0.158</u>	<u>0.058</u>	<u>0.098</u>	0.027	0.000	0.000	0.019	0

BOL=Bolores, CDM=Cova da Moura, PAI=Pai Mogo, CAN=Canary Islanders, CAP=NW African Capsian, KAB=NW African Kabyle Berbers, SHA=NW African Shawia Berbers, GRK=Greece, ITY=Italy, TRK=Turkey (see text Section 1.2 for details).

Underlined MMD distances indicate significant difference at the 0.025 level.

TABLE 2. Dental trait percentages (%) and number of individuals scored (n) for the Portuguese and comparative samples

Sample		BOL	CDM	PAI	CAN	CAP	KAB	SHA	GRK	ITY	TRK
Incisor UI1 (SU 1)	%	7.1	3.2	3.6	11.5	0.0	0.0	0.0	1.5	4.00	0.0
	n	14	31	28	130	5	29	26	68	76	36
Mandibular Curvature UI1 (SU 2-4)	%	19.4	23.1	10.4	18.2	25.0	50.0	28.6	20.0	24.1	40.0
	n	31	39	48	22	4	8	7	5	29	10
Mandibular Torus (SU 2-3)	%	0.0	0.0	3.7	23.9	0.0	3.5	0.0	4.3	10.4	0.0
	n	3	10	27	130	10	29	25	70	77	35
Mandibular Folding UI1 (SU 2-6)	%	4.6	13.2	6.4	14.3	0.0	14.3	0.0	0.0	26.9	0.0
	n	22	38	47	14	5	7	7	5	26	10
Mandibular Shoveling UI1 (SU 2-6)	%	3.5	5.1	8.9	0.0	0.0	12.5	25.0	0.0	0.0	0.0
	n	29	39	45	17	5	8	8	5	27	10
Mandibular Cupation Groove UI2 (SU +)	%	7.7	0.0	14.3	20.8	60.0	21.4	46.2	35.0	13.3	15.4
	n	26	19	49	24	5	14	13	20	30	13

Variable	Unit	Mean	SD	Median	Q1	Q3	Min	Max	Skewness	Kurtosis	Shapiro-Wilk
Periculum Dentale UI2	%	47.8	38.9	29.2	33.3	60.0	50.0	25.0	5.3	36.7	15.4
(n=ASU 2-6)	n	23	18	48	18	5	12	12	19	30	13
Human Canine UC	%	4.0	0.0	0.0	2.7	22.2	0.0	0.0	8.7	2.6	0.0
(n=ASU 1-3)	n	25	39	49	37	9	16	14	23	39	19
1 Acc. Ridge UC	%	30.0	51.5	73.9	21.7	42.9	27.3	22.2	8.3	19.2	6.3
(n=ASU 2-5)	n	20	33	46	23	7	11	9	12	26	16
Pyrocone UM2	%	57.1	56.0	63.8	77.8	100.0	63.6	68.4	50.0	59.7	
(n=ASU 3-5)	n	14	25	47	99	10	22	19	54	72	
usp 5 UM1	%	29.4	8.7	17.0	13.0	30.0	11.8	10.0	5.7	17.5	
(n=ASU 2-5)	n	17	23	47	77	10	17	20	53	63	
Carabelli's Trait UM1	%	6.3	29.6	26.5	17.4	83.3	26.3	11.1	16.7	21.0	
(n=ASU 5-7)	n	16	27	49	69	6	19	18	48	62	
Parastyle UM3	%	0.00	0.0	4.8	1.3	0.0	0.0	7.7	0.0	0.0	
(n=ASU 2-5)	n	15	22	42	76	9	22	13	33	41	
Chamel Extension UM1	%	18.2	3.7	2.3	1.7	0.0	0.0	4.8	3.7	1.5	

Phenotypic trait	Unit	ASU 1	ASU 2	ASU 3	ASU 4	ASU 5	ASU 6	ASU 7	ASU 8	ASU 9
ASU 1-3)	n	11	27	44	119	13	23	21	54	69
ASU 1-3)	%	45.5	70.0	69.8	52.5	33.3	52.2	52.2	61.9	59.3
ASU 2+)	n	11	40	43	118	12	23	23	63	59
ASU 2+)	%	92.3	57.1	89.8	84.8	85.7	68.4	72.2	58.3	76.9
ASU 3+)	n	13	21	49	79	7	19	18	36	39
ASU 3+)	%	10.3	2.9	0.0	5.3	0.0	6.3	0.0	0.0	9.6
ASU P or R)	n	29	34	49	38	10	16	13	73	83
ASU P or R)	%	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	2.7
ASU +)	n	35	40	44	85	12	22	23	44	74
ASU +)	%	0.0	6.9	8.3	16.5	16.7	3.5	23.1	17.7	23.5
ASU -)	n	18	29	48	133	12	29	26	68	81
ASU -)	%	0.0	0.0	0.0	7.0	0.0	5.0	0.0	3.03	4.9
ASU 0.5 mm)	n	14	20	3	131	5	20	23	66	82
ASU 0.5 mm)	%	30.8	63.3	31.7	64.0	53.9	53.9	61.5	50.0	23.3
ASU 3-9)	n	13	30	41	50	13	13	13	10	43

Table 1: Prevalence of facial wrinkles and folds in different age groups and genders										
Feature	Gender	Age Group								
		18-24	25-34	35-44	45-54	55-64	65-74	75-84	85+	Total
Upper Face										
Forehead Wrinkle LM1	F	12.3	15.6	18.9	22.1	25.4	28.7	32.0	35.3	21.4
Forehead Wrinkle LM2	F	8.7	10.2	12.5	14.8	17.1	19.4	21.7	24.0	14.5
Glabella Wrinkle LM1	F	5.4	6.8	8.2	9.6	11.0	12.4	13.8	15.2	9.3
Glabella Wrinkle LM2	F	3.2	4.1	5.0	5.9	6.8	7.7	8.6	9.5	5.6
Upper Lip Wrinkle LM1	F	7.1	8.5	9.9	11.3	12.7	14.1	15.5	16.9	10.8
Upper Lip Wrinkle LM2	F	4.5	5.4	6.3	7.2	8.1	9.0	9.9	10.8	6.7
Lower Face										
Marionette Line LM1	F	6.8	8.2	9.6	11.0	12.4	13.8	15.2	16.6	10.2
Marionette Line LM2	F	3.9	4.8	5.7	6.6	7.5	8.4	9.3	10.2	6.3
Chin Wrinkle LM1	F	2.1	2.9	3.7	4.5	5.3	6.1	6.9	7.7	4.4
Chin Wrinkle LM2	F	1.3	1.7	2.1	2.5	2.9	3.3	3.7	4.1	2.5
Lower Lip Wrinkle LM1	F	5.6	6.5	7.4	8.3	9.2	10.1	11.0	11.9	7.8
Lower Lip Wrinkle LM2	F	3.4	4.0	4.6	5.2	5.8	6.4	7.0	7.6	4.9
Mid Face										
Cheek Wrinkle LM1	F	4.2	5.1	6.0	6.9	7.8	8.7	9.6	10.5	6.5
Cheek Wrinkle LM2	F	2.5	3.0	3.5	4.0	4.5	5.0	5.5	6.0	3.8
Nasolabial Fold LM1	F	9.5	11.2	12.9	14.6	16.3	18.0	19.7	21.4	13.9
Nasolabial Fold LM2	F	5.8	6.9	8.0	9.1	10.2	11.3	12.4	13.5	8.6
Smile Line LM1	F	3.7	4.5	5.3	6.1	6.9	7.7	8.5	9.3	5.9
Smile Line LM2	F	2.2	2.7	3.2	3.7	4.2	4.7	5.2	5.7	3.5
General Features										
Deep Frown LM1	M	18.5	22.1	25.7	29.3	32.9	36.5	40.1	43.7	28.8
Deep Frown LM2	M	10.2	12.5	14.8	17.1	19.4	21.7	24.0	26.3	16.4
Wrinkle Pattern LM1	M	7.8	9.3	10.8	12.3	13.8	15.3	16.8	18.3	11.5
Wrinkle Pattern LM2	M	4.9	5.9	6.9	7.9	8.9	9.9	10.9	11.9	7.4
Smile Line LM1	M	6.3	7.4	8.5	9.6	10.7	11.8	12.9	14.0	9.1
Smile Line LM2	M	3.8	4.5	5.2	5.9	6.6	7.3	8.0	8.7	5.4
Total Prevalence										
Forehead Wrinkle LM1	All	10.5	12.8	15.1	17.4	19.7	22.0	24.3	26.6	17.2
Forehead Wrinkle LM2	All	6.5	7.8	9.1	10.4	11.7	13.0	14.3	15.6	10.0
Glabella Wrinkle LM1	All	4.5	5.5	6.5	7.5	8.5	9.5	10.5	11.5	7.0
Glabella Wrinkle LM2	All	2.8	3.4	4.0	4.6	5.2	5.8	6.4	7.0	4.3
Upper Lip Wrinkle LM1	All	7.5	8.8	10.1	11.4	12.7	14.0	15.3	16.6	10.5
Upper Lip Wrinkle LM2	All	4.7	5.6	6.5	7.4	8.3	9.2	10.1	11.0	6.9
Marionette Line LM1	All	6.5	7.8	9.1	10.4	11.7	13.0	14.3	15.6	9.8
Marionette Line LM2	All	3.7	4.5	5.3	6.1	6.9	7.7	8.5	9.3	5.9
Chin Wrinkle LM1	All	2.5	3.0	3.5	4.0	4.5	5.0	5.5	6.0	3.8
Chin Wrinkle LM2	All	1.5	1.8	2.1	2.4	2.7	3.0	3.3	3.6	2.2
Lower Lip Wrinkle LM1	All	6.5	7.6	8.7	9.8	10.9	12.0	13.1	14.2	9.3
Lower Lip Wrinkle LM2	All	4.0	4.7	5.4	6.1	6.8	7.5	8.2	8.9	5.6
Cheek Wrinkle LM1	All	5.0	5.9	6.8	7.7	8.6	9.5	10.4	11.3	7.4
Cheek Wrinkle LM2	All	3.0	3.6	4.2	4.8	5.4	6.0	6.6	7.2	4.5
Nasolabial Fold LM1	All	11.5	13.6	15.7	17.8	19.9	22.0	24.1	26.2	17.5
Nasolabial Fold LM2	All	6.8	8.1	9.4	10.7	12.0	13.3	14.6	15.9	10.2
Smile Line LM1	All	5.0	5.9	6.8	7.7	8.6	9.5	10.4	11.3	7.2
Smile Line LM2	All	3.0	3.6	4.2	4.8	5.4	6.0	6.6	7.2	4.5
Gender Comparison										
Forehead Wrinkle LM1	F vs M	10.5	12.8	15.1	17.4	19.7	22.0	24.3	26.6	17.2
Forehead Wrinkle LM2	F vs M	6.5	7.8	9.1	10.4	11.7	13.0	14.3	15.6	10.0
Glabella Wrinkle LM1	F vs M	4.5	5.5	6.5	7.5	8.5	9.5	10.5	11.5	7.0
Glabella Wrinkle LM2	F vs M	2.8	3.4	4.0	4.6	5.2	5.8	6.4	7.0	4.3
Upper Lip Wrinkle LM1	F vs M	7.5	8.8	10.1	11.4	12.7	14.0	15.3	16.6	10.5
Upper Lip Wrinkle LM2	F vs M	4.7	5.6	6.5	7.4	8.3	9.2	10.1	11.0	6.9
Marionette Line LM1	F vs M	6.5	7.8	9.1	10.4	11.7	13.0	14.3	15.6	9.8
Marionette Line LM2	F vs M	3.7	4.5	5.3	6.1	6.9	7.7	8.5	9.3	5.9
Chin Wrinkle LM1	F vs M	2.5	3.0	3.5	4.0	4.5	5.0	5.5	6.0	3.8
Chin Wrinkle LM2	F vs M	1.5	1.8	2.1	2.4	2.7	3.0	3.3	3.6	2.2
Lower Lip Wrinkle LM1	F vs M	6.5	7.6	8.7	9.8	10.9	12.0	13.1	14.2	9.3
Lower Lip Wrinkle LM2	F vs M	4.0	4.7	5.4	6.1	6.8	7.5	8.2	8.9	5.6
Cheek Wrinkle LM1	F vs M	5.0	5.9	6.8	7.7	8.6	9.5	10.4	11.3	7.4
Cheek Wrinkle LM2	F vs M	3.0	3.6	4.2	4.8	5.4	6.0	6.6	7.2	4.5
Nasolabial Fold LM1	F vs M	11.5	13.6	15.7	17.8	19.9	22.0	24.1	26.2	17.5
Nasolabial Fold LM2	F vs M	6.8	8.1	9.4	10.7	12.0	13.3	14.6	15.9	10.2
Smile Line LM1	F vs M	5.0	5.9	6.8	7.7	8.6	9.5	10.4	11.3	7.2
Smile Line LM2	F vs M	3.0	3.6	4.2	4.8	5.4	6.0	6.6	7.2	4.5
Age Group Comparison										
Forehead Wrinkle LM1	18-24 vs 85+	10.5	12.8	15.1	17.4	19.7	22.0	24.3	26.6	17.2
Forehead Wrinkle LM2	18-24 vs 85+	6.5	7.8	9.1	10.4	11.7	13.0	14.3	15.6	10.0
Glabella Wrinkle LM1	18-24 vs 85+	4.5	5.5	6.5	7.5	8.5	9.5	10.5	11.5	7.0
Glabella Wrinkle LM2	18-24 vs 85+	2.8	3.4	4.0	4.6	5.2	5.8	6.4	7.0	4.3
Upper Lip Wrinkle LM1	18-24 vs 85+	7.5	8.8	10.1	11.4	12.7	14.0	15.3	16.6	10.5
Upper Lip Wrinkle LM2	18-24 vs 85+	4.7	5.6	6.5	7.4	8.3	9.2	10.1	11.0	6.9
Marionette Line LM1	18-24 vs 85+	6.5	7.8	9.1	10.4	11.7	13.0	14.3	15.6	9.8
Marionette Line LM2	18-24 vs 85+	3.7	4.5	5.3	6.1	6.9	7.7	8.5	9.3	5.9
Chin Wrinkle LM1	18-24 vs 85+	2.5	3.0	3.5	4.0	4.5	5.0	5.5	6.0	3.8
Chin Wrinkle LM2	18-24 vs 85+	1.5	1.8	2.1	2.4	2.7	3.0	3.3	3.6	2.2
Lower Lip Wrinkle LM1	18-24 vs 85+	6.5	7.6	8.7	9.8	10.9	12.0	13.1	14.2	9.3
Lower Lip Wrinkle LM2	18-24 vs 85+	4.0	4.7	5.4	6.1	6.8	7.5	8.2	8.9	5.6
Cheek Wrinkle LM1	18-24 vs 85+	5.0	5.9	6.8	7.7	8.6	9.5	10.4	11.3	7.4
Cheek Wrinkle LM2	18-24 vs 85+	3.0	3.6	4.2	4.8	5.4	6.0	6.6	7.2	4.5
Nasolabial Fold LM1	18-24 vs 85+	11.5	13.6	15.7	17.8	19.9	22.0	24.1	26.2	17.5
Nasolabial Fold LM2	18-24 vs 85+	6.8	8.1	9.4	10.7	12.0	13.3	14.6	15.9	10.2
Smile Line LM1	18-24 vs 85+	5.0	5.9	6.8	7.7	8.6	9.5	10.4	11.3	7.2
Smile Line LM2	18-24 vs 85+	3.0	3.6	4.2	4.8	5.4	6.0	6.6	7.2	4.5
Total Prevalence by Age Group										
Forehead Wrinkle LM1	18-24	10.5	12.8	15.1	17.4	19.7	22.0	24.3	26.6	17.2
Forehead Wrinkle LM2	18-24	6.5	7.8	9.1	10.4	11.7	13.0	14.3	15.6	10.0
Glabella Wrinkle LM1	18-24	4.5	5.5	6.5	7.5	8.5	9.5	10.5	11.5	7.0
Glabella Wrinkle LM2	18-24	2.8	3.4	4.0	4.6	5.2	5.8	6.4	7.0	4.3
Upper Lip Wrinkle LM1	18-24	7.5	8.8	10.1	11.4	12.7	14.0	15.3	16.6	10.5
Upper Lip Wrinkle LM2	18-24	4.7	5.6	6.5	7.4	8.3	9.2	10.1	11.0	6.9
Marionette Line LM1	18-24	6.5	7.8	9.1	10.4	11.7	13.0	14.3	15.6	9.8
Marionette Line LM2	18-24	3.7	4.5	5.3	6.1	6.9	7.7	8.5	9.3	5.9
Chin Wrinkle LM1	18-24	2.5	3.0	3.5	4.0	4.5	5.0	5.5	6.0	3.8
Chin Wrinkle LM2	18-24	1.5	1.8	2.1	2.4	2.7	3.0	3.3	3.6	2.2
Lower Lip Wrinkle LM1	18-24	6.5	7.6	8.7	9.8	10.9	12.0	13.1	14.2	9.3
Lower Lip Wrinkle LM2	18-24	4.0	4.7	5.4	6.1	6.8	7.5	8.2	8.9	5.6
Cheek Wrinkle LM1	18-24	5.0	5.9	6.8	7.7	8.6	9.5	10.4	11.3	7.4
Cheek Wrinkle LM2	18-24	3.0	3.6	4.2	4.8	5.4	6.0	6.6	7.2	4.5
Nasolabial Fold LM1	18-24	11.5	13.6	15.7	17.8	19.9	22.0	24.1	26.2	17.5
Nasolabial Fold LM2	18-24	6.8	8.1	9.4	10.7	12.0	13.3	14.6	15.9	10.2

Number LM2	%	95.0	82.5	100.0	93.8	85.7	88.9	95.5	91.3	100.0
SU 2+)	n	20	40	43	64	14	18	22	23	57
molar Angle LM3	%	0.0	8.8	5.6	23.4	23.1	21.4	23.5	13.0	16.3
SU +)	n	13	34	36	77	13	14	17	23	43

lores, CDM=Cova da Moura, PAI=Pai Mogo, CAN=Canary Islanders, CAP=NW African Capsian, KAB=NW African Kabyle, SHA=NW African Shawia Berbers, GRK=Greece, ITY=Italy, TRK=Turkey (see text Section 1.2 for details).

nk-scale trait breakpoints from Irish (1993, 1997, 1998a, 1998b, 2005, 2006) and Scott and Turner (1997), with modification

The three Portuguese and seven comparative dental samples.

	Geographic Origin	Affiliation	Dates	n	Institution*
BOL)	Estremadura	Neolithic/Copper Age	2800-2600 BCE	36	MMLT
Moura	Estremadura	Neolithic/Copper Age	3700-2300 BCE	41	MMLT
I (PAI)	Estremadura	Neolithic/Copper Age	3000-2600 BCE	49	MMLT
a					
landers	Canary Islands	Precontact Guanche	400-900(?) CE	163	NMNH, MH
CAP)	Algeria and Tunisia	Early to Middle Neolithic	6500–3000 BCE	24	UM
KAB)	Northern Algeria	Berber	1800–1900s CE	32	MH
SHA)	Southern Algeria	Berber	1800–1900s CE	26	MH
GRK)	Greece	Classic to historic	475–300 BCE to historic	77	AMNH
)	Italy	Roman and historic	30 BC–AD 395 to historic	90	NHM
RK)	Anatolia/Turkey and Cyprus	Classic to Ottoman	>300 BCE to ~14 th Cent CE	40	AMNH

ns in which the samples are curated: MMLT = Museu Municipal Leonel Trindade - Torres Vedras; NMNH = National Museum of Natural History, Washington, DC; MH = Musée de l'Homme, Paris; UM = University of Minnesota, Minneapolis; AMNH = American Museum of Natural History, New York; NHM = Natural History Museum, London.